

Idaho Reported Sexually Transmitted Disease 2010



Idaho Department of Health and Welfare
Office of Epidemiology, Food Protection,
and Immunization
450 W. State St., 4th Floor
Boise, ID 83720

Acknowledgements

Jared Bartschi, MHE, IDHW Office of Epidemiology, Food Protection, and Immunization

Christine Hahn, MD, IDHW Office of Epidemiology, Food Protection, and Immunization

Rhonda Jones, IDHW Office of Epidemiology, Food Protection, and Immunization

Cathy Ruby, IDHW Office of Epidemiology, Food Protection, and Immunization

District Health Departments

Contents

Acknowlegements	i
Listing of Tables and Figures.....	iv
Methods	v
Historical trends	1
Maps.....	3
Statewide Totals	4
Chancroid	5
Chlamydia.....	7
Gonorrhea	9
Syphilis	11
HIV	15
AIDS	16
District Tables	17
District 1	17
District 2	18
District 3	19
District 4	20
District 5	21
District 6	22
District 7	23

Listing of Tables and Figures

Tables

Table 1. Sexually Transmitted Disease (STD)—Idaho, 2010	4
Table 2. STD by health district of residence—Idaho, 2010.....	4
Table 3. Chancroid by sex, race/ethnicity, and age group—Idaho, 2010	5
Table 4. Chlamydia by sex, race/ethnicity, and age group—Idaho, 2010.....	7
Table 5. Chlamydia by sex and county of residence—Idaho, 2010	8
Table 6. Gonorrhea by sex, race/ethnicity, and age group—Idaho, 2010.....	9
Table 7. Gonorrhea by sex and county of residence—Idaho, 2010	10
Table 8. Early syphilis (<1 yr duration) by sex, race/ethnicity, and age group—Idaho, 2010	11
Table 9. Early syphilis (<1 yr) by sex and county of residence—Idaho, 2010	12
Table 10. Syphilis (all stages) by sex, race/ethnicity, and age group—Idaho, 2010.....	13
Table 11. Syphilis (all stages) by sex and county of residence—Idaho, 2010.....	14
Table 12. HIV by sex, race/ethnicity, and age group—Idaho, 2010.....	15
Table 13. AIDS by sex, race/ethnicity, and age group—Idaho, 2010	16
Table 14. STD, HIV, and AIDS by diagnosis and age group, Health District 1, 2010.....	17
Table 15. STD, HIV, and AIDS by year of report, Health District 1, 2001–2010	17
Table 16. STD, HIV, and AIDS by diagnosis and age group, Health District 2, 2010.....	18
Table 17. STD, HIV, and AIDS by year of report, Health District 2, 2001–2010.....	18
Table 18. STD, HIV, and AIDS by diagnosis and age group, Health District 3, 2010.....	19
Table 19. STD, HIV, and AIDS by year of report, Health District 3, 2001–2010.....	19
Table 20. STD, HIV, and AIDS by diagnosis and age group, Health District 4, 2010.....	20
Table 21. STD, HIV, and AIDS by year of report, Health District 4, 2001–2010	20
Table 22. STD, HIV, and AIDS by diagnosis and age group, Health District 5, 2010.....	21
Table 23. STD, HIV, and AIDS by year of report, Health District 5, 2001–2010.....	21
Table 24. STD, HIV, and AIDS by diagnosis and age group, Health District 6, 2010.....	22
Table 25. STD, HIV, and AIDS by year of report, Health District 6, 2001–2010	22
Table 26. STD, HIV, and AIDS by diagnosis and age group, Health District 7, 2010.....	23
Table 27. STD, HIV, and AIDS by year of report, Health District 7, 2001–2010	23

Figures

Figure 1. Idaho counties and health districts	v
Figure 2. Chancroid by year of report—Idaho, 1991–2010	1
Figure 3. Chlamydia by year of report—Idaho, 1991–2010	1
Figure 4. Gonorrhea by year of report—Idaho, 1991–2010	1
Figure 5. Syphilis by category and year of report—Idaho, 1991–2010	2
Figure 6. HIV (not AIDS) by year of report—Idaho, 1991–2010	2
Figure 7. AIDS by year of report—Idaho, 1991–2010	2
Figure 8. Reported chlamydia incidence rate by county—Idaho, 2010.....	3
Figure 9. Reported gonorrhea incidence rate by county—Idaho, 2010	3

Methods

Chlamydia, gonorrhea, syphilis, HIV/AIDS, and chancroid are sexually transmitted diseases (STDs) which are required to be reported by name. Providers who diagnose these STDs report to district health departments or the Idaho Department of Health and Welfare. CDC case definitions are used:

<http://www.cdc.gov/ncphi/disss/nndss/casedef/>

STD reports are mandated to include patient-level information, including sex, race, ethnicity, address, diagnosis, and diagnosing provider. This report contains aggregate data from reported cases of STD in individuals residing in Idaho.


“Early syphilis” presented in this report is meant to show syphilis that is recently acquired. This includes cases meeting the primary, secondary, and early latent syphilis case definitions. Cases categorized as late latent syphilis, latent syphilis of unknown duration, and neurosyphilis have no evidence of recent syphilis infection and may have long-standing syphilis infection. Because these may be entirely different populations than those involved in recent transmission, it is helpful to separate them. They are presented in tables and graphs as “Syphilis, late/unk/neuro.” Congenital syphilis is also presented separately.

This report presents data from HIV and AIDS among residents of Idaho at first HIV or AIDS diagnosis. HIV/AIDS Surveillance guidelines state, “Cases of HIV and AIDS are reported based on the place of residence at diagnosis of HIV and/or AIDS regardless of where the exposure may have occurred.” This applies even when the residence state is a state that did not require HIV infection reporting at the time of diagnosis or uses a non-name identifier, if the diagnosis can be verified through patient records. Ongoing nationwide HIV/AIDS case de-duplication processes can result in identification of cases reported in Idaho which were reported previously outside of Idaho. This publication supersedes prior publications. For more detailed analysis of HIV/AIDS in Idaho, consult the latest Epidemiologic Profile of HIV/AIDS in Idaho (<http://www.healthandwelfare.idaho.gov>).

Historical trends are presented for the past 20 years through 2010 or from the year the disease/condition was made reportable in Idaho, if fewer than 20 years.

Race is collected independent from ethnicity. A person with Hispanic ethnicity may be of any race. For practical purposes, all reports of infection in individuals with Hispanic ethnicity are categorized as “Hispanic” in this report; all other race/ethnic categories exclude Hispanic ethnicity. Case reports can include more than one race, categorized in this document as “Other race”.

Figure 1. Idaho counties and health districts



Rates in this report are raw rates per 100,000 population for the geographic area calculated. U.S. Census Bureau estimates for 2010 ("Intercensal Estimates of the Resident Population for Counties of Idaho: April 1, 2000 to July 1, 2010" [release date September 2011]) were used for county population denominators. Rates are not adjusted for age using a standardized population.

Historical trends

Figure 2. Chancroid by year of report—Idaho, 1991–2010




Figure 3. Chlamydia by year of report—Idaho, 1991–2010




Figure 4. Gonorrhea by year of report—Idaho, 1991–2010






Figure 5. Syphilis by category¹ and year of report—Idaho, 1991–2010




1-See methods section for an explanation of the presented syphilis categories.

Figure 6. HIV (not AIDS) by year of report—Idaho, 1991–2010¹



1. HIV became reportable in Idaho in 1986. This graph presents data by year of report, not year of diagnosis. For a more accurate depiction of HIV/AIDS trends using diagnosis year, see the latest Idaho HIV/AIDS Epidemiologic Profile at: www.healthandwelfare.idaho.gov.

Figure 7. AIDS by year of report—Idaho, 1991–2010¹



1. AIDS became reportable in Idaho in 1984. This graph presents data by year of report, not year of diagnosis, and may include AIDS progressions from previously-reported HIV. For a more accurate depiction of HIV/AIDS trends using diagnosis year, see the latest Idaho HIV/AIDS Epidemiologic Profile at: www.healthandwelfare.idaho.gov.

Maps

Figure 8. Reported chlamydia incidence rate by county—Idaho, 2010





Figure 9. Reported gonorrhea incidence rate by county—Idaho, 2010



Statewide Totals

Table 1. Sexually Transmitted Disease (STD)—Idaho, 2010

Condition	Sex			TOTAL
	Male	Female	Unknown	
Chancroid	0	0	0	0
Chlamydia	1,183	3,014	11	4,208
Gonorrhea	78	68	1	147
Syphilis, early	11	1	0	12
Syphilis, late/unk/neuro	6	3	0	9
Syphilis, congenital	0	1	0	1
HIV (not AIDS)	28	4	0	32
AIDS	42	9	0	51
TOTAL	1,348	3,100	12	4,460

Table 2. STD by health district of residence—Idaho, 2010

Condition	DISTRICT							TOTAL
	1	2	3	4	5	6	7	
Chancroid	0	0	0	0	0	0	0	0
Chlamydia	628	312	815	1,344	406	411	292	4,208
Gonorrhea	13	24	34	52	8	5	11	147
Syphilis, early	0	0	4	7	0	1	0	12
Syphilis, late/unk/neuro	1	0	2	4	0	2	0	9
Syphilis, congenital	0	0	1	0	0	0	0	1
HIV (not AIDS)	3	3	2	14	1	3	6	32
AIDS	5	3	11	21	2	3	6	51
TOTAL	650	342	869	1,442	417	425	315	4,460

Chancroid

Table 3. Chancroid by sex, race/ethnicity, and age group—Idaho, 2010

No chancroid reported during 2010

Page intentionally left blank

Chlamydia

Table 4. Chlamydia by sex, race/ethnicity, and age group—Idaho, 2010

Age at Diagnosis	Sex										TOTAL		
	Male					Female							
	Race					Race							
	American Indian/Alaska Native	Asian/Pacific Islander	Black (not Hispanic)	White (not Hispanic)	Hispanic	Total Males	Percent of Males	American Indian/Alaska Native	Asian/Pacific Islander	Black (not Hispanic)	Total Females	Percent of Females	
0- 4 yrs	0	0	0	1	0	0	0%	0	0	0	8	0%	
5- 9 yrs	0	0	0	0	0	0	0%	0	0	0	1	0%	
10-14 yrs	0	0	0	1	0	2	3	0%	3	0	0	21	1%
15-19 yrs	5	0	6	98	34	79	222	19%	26	9	9	1,060	35%
20-24 yrs	13	7	14	246	64	165	509	43%	35	11	11	1,206	40%
25-29 yrs	2	0	12	130	41	68	253	21%	21	5	3	420	14%
30-34 yrs	0	1	2	44	9	43	99	8%	5	1	0	181	6%
35-39 yrs	0	0	5	14	6	23	48	4%	5	0	0	66	2%
40-44 yrs	1	0	0	10	2	12	25	2%	0	0	0	31	1%
45-54 yrs	0	0	0	10	2	9	21	2%	1	0	0	14	0%
55-64 yrs	0	0	0	0	0	1	1	0%	0	0	0	3	0%
65+ yrs	0	0	0	0	0	1	1	0%	0	0	0	2	0%
Unknown	0	0	0	0	0	0	0	0%	0	0	0	1	0%
TOTAL	21	8	39	554	158	403	1,183	100%	96	26	23	3,014	100%
												4,208	100%

Total race/ethnicity	Number	Percent
American Indian/Alaska Native	117	3%
Asian/Pacific Islander	34	1%
Black (not Hispanic)	62	1%
White (not Hispanic)	2,025	48%
Hispanic	510	12%
Unknown	1,460	35%
TOTAL	4,208	100%

1. Total includes 11 cases of unknown sex: 1 Hispanic/Latino, and 10 Unknown race/ethnicity.

Note: 0-4 year age group cases are Ophthalmia Neonatorum, chlamydial infection of the infant conjunctiva, which occurs during some births to chlamydia-infected mothers. One case in the 5-9 year age group was chlamydia conjunctivitis.

Table 5. Chlamydia by sex and county of residence—Idaho, 2010

SEX						
County	Male	Female	Unknown	TOTAL	Population	Rate per 100,000
Ada	373	825	5	1,203	388,624	309.6
Adams	4	0	0	4	3,403	117.5
Bannock	61	225	0	286	83,548	342.3
Bear Lake	0	3	0	3	5,721	52.4
Benewah	7	15	0	22	9,270	237.3
Bingham	19	65	0	84	45,371	185.1
Blaine	18	24	0	42	22,166	189.5
Boise	1	7	0	8	7,536	106.2
Bonner	26	40	0	66	41,550	158.8
Bonneville	52	167	2	221	102,738	215.1
Boundary	3	10	0	13	11,147	116.6
Butte	1	2	0	3	2,820	106.4
Camas	0	0	0	0	1,078	0.0
Canyon	177	483	1	661	188,584	350.5
Caribou	1	4	0	5	6,936	72.1
Cassia	3	39	0	42	22,230	188.9
Clark	2	1	0	3	991	302.7
Clearwater	4	3	0	7	8,017	87.3
Custer	1	7	0	8	4,222	189.5
Elmore	48	74	1	123	28,606	430.0
Franklin	4	9	0	13	12,697	102.4
Fremont	7	8	0	15	12,693	118.2
Gem	18	34	0	52	16,188	321.2
Gooding	5	19	0	24	14,592	164.5
Idaho	4	22	0	26	15,628	166.4
Jefferson	9	11	0	20	25,126	79.6
Jerome	15	46	0	61	21,789	280.0
Kootenai	149	356	1	506	140,679	359.7
Latah	36	83	0	119	38,590	308.4
Lemhi	1	3	0	4	7,946	50.3
Lewis	3	7	0	10	3,675	272.1
Lincoln	4	11	0	15	4,679	320.6
Madison	0	15	0	15	38,550	38.9
Minidoka	14	44	0	58	19,438	298.4
Nez Perce	43	107	0	150	39,499	379.8
Oneida	2	3	0	5	4,228	118.3
Owyhee	4	19	0	23	11,129	206.7
Payette	11	43	0	54	23,140	233.4
Power	4	8	0	12	8,045	149.2
Shoshone	7	14	0	21	12,566	167.1
Teton	1	5	0	6	9,406	63.8
Twin Falls	35	128	1	164	76,386	214.7
Valley	2	8	0	10	8,416	118.8
Washington	4	17	0	21	10,153	206.8
STATE	1,183	3,014	11	4,208	1,559,796	269.8

Gonorrhea

Table 6. Gonorrhea by sex, race/ethnicity, and age group—Idaho, 2010

Age at Diagnosis	Sex										TOTAL							
	Male					Female												
	Race					Race												
	American Indian/Alaska Native	Asian/Pacific Islander	Black (not Hispanic)	White (not Hispanic)	Hispanic	Other/Unknown	Total Males	Percent of Males	American Indian/Alaska Native	Asian/Pacific Islander	Black (not Hispanic)	White (not Hispanic)	Hispanic	Other/Unknown	Total Females	Percent of Females	Total Cases ¹	Percent of Total
0- 4 yrs	0	0	0	0	0	0	0	0%	0	0	0	0	0	0	0	0%	0	0%
5- 9 yrs	0	0	0	0	0	0	0	0%	0	0	0	0	0	0	0	0%	0	0%
10-14 yrs	0	0	0	0	0	0	0	0%	0	0	0	1	0	0	1	1%	1	1%
15-19 yrs	0	0	0	3	1	1	5	6%	2	0	2	7	0	5	16	24%	21	14%
20-24 yrs	2	0	2	12	4	19	39	50%	2	0	1	11	3	12	29	43%	68	46%
25-29 yrs	0	0	2	9	2	8	21	27%	0	0	1	6	1	4	12	18%	34	23%
30-34 yrs	0	0	0	0	0	5	5	6%	0	0	1	3	0	3	7	10%	12	8%
35-39 yrs	0	0	0	2	0	1	3	4%	0	0	0	1	0	1	2	3%	5	3%
40-44 yrs	0	0	0	1	0	3	4	5%	0	0	0	0	1	0	1	1%	5	3%
45-54 yrs	0	0	0	0	0	0	0	0%	0	0	0	0	0	0	0	0%	0	0%
55-64 yrs	0	0	0	0	0	1	1	1%	0	0	0	0	0	0	0	0%	1	1%
65+ yrs	0	0	0	0	0	0	0	0%	0	0	0	0	0	0	0	0%	0	0%
Unknown	0	0	0	0	0	0	0	0%	0	0	0	0	0	0	0	0%	0	0%
TOTAL	2	0	4	27	7	38	78	100%	4	0	5	29	5	25	68	100%	147	100%

Total race/ethnicity	Number	Percent
American Indian/Alaska Native	6	4%
Asian/Pacific Islander	0	0%
Black (not Hispanic)	9	6%
White (not Hispanic)	56	38%
Hispanic	12	8%
Unknown	64	44%
TOTAL	147	100%

1. Total includes 1 case of unknown sex and unknown race/ethnicity.

Table 7. Gonorrhea by sex and county of residence—Idaho, 2010

County	SEX			TOTAL	Population	Rate per 100,000
	Male	Female	Unknown			
Ada	33	16	0	49	388,624	12.6
Adams	2	0	0	2	3,403	58.8
Bannock	1	3	0	4	83,548	4.8
Bear Lake	0	0	0	0	5,721	0.0
Benewah	2	2	0	4	9,270	43.1
Bingham	0	1	0	1	45,371	2.2
Blaine	1	0	0	1	22,166	4.5
Boise	0	0	0	0	7,536	0.0
Bonner	0	2	0	2	41,550	4.8
Bonneville	1	6	1	8	102,738	7.8
Boundary	0	0	0	0	11,147	0.0
Butte	0	0	0	0	2,820	0.0
Camas	0	0	0	0	1,078	0.0
Canyon	11	13	0	24	188,584	12.7
Caribou	0	0	0	0	6,936	0.0
Cassia	0	0	0	0	22,230	0.0
Clark	0	0	0	0	991	0.0
Clearwater	0	0	0	0	8,017	0.0
Custer	1	0	0	1	4,222	23.7
Elmore	2	1	0	3	28,606	10.5
Franklin	0	0	0	0	12,697	0.0
Fremont	0	1	0	1	12,693	7.9
Gem	0	2	0	2	16,188	12.4
Gooding	0	0	0	0	14,592	0.0
Idaho	1	0	0	1	15,628	6.4
Jefferson	0	0	0	0	25,126	0.0
Jerome	2	0	0	2	21,789	9.2
Kootenai	4	2	0	6	140,679	4.3
Latah	6	4	0	10	38,590	25.9
Lemhi	0	0	0	0	7,946	0.0
Lewis	0	0	0	0	3,675	0.0
Lincoln	0	0	0	0	4,679	0.0
Madison	0	1	0	1	38,550	2.6
Minidoka	0	0	0	0	19,438	0.0
Nez Perce	3	10	0	13	39,499	32.9
Oneida	0	0	0	0	4,228	0.0
Owyhee	0	0	0	0	11,129	0.0
Payette	3	3	0	6	23,140	25.9
Power	0	0	0	0	8,045	0.0
Shoshone	1	0	0	1	12,566	8.0
Teton	0	0	0	0	9,406	0.0
Twin Falls	4	1	0	5	76,386	6.5
Valley	0	0	0	0	8,416	0.0
Washington	0	0	0	0	10,153	0.0
STATE	78	68	1	147	1,559,796	9.4

Syphilis

Table 8. Early syphilis¹ (<1 yr duration) by sex, race/ethnicity, and age group—Idaho, 2010

Age at Diagnosis	Sex										TOTAL								
	Male					Female													
	Race	American Indian/Alaska Native	Asian/Pacific Islander	Black (not Hispanic)	White (not Hispanic)	Hispanic	Other/Unknown	Total Males	Percent of Males	American Indian/Alaska Native	Asian/Pacific Islander	Black (not Hispanic)	White (not Hispanic)	Hispanic	Other/Unknown	Total Females	Percent of Females		
0- 4 yrs	0	0	0	0	0	0	0	0	0%	0	0	0	0	0	0	0	0%	0	0%
5- 9 yrs	0	0	0	0	0	0	0	0	0%	0	0	0	0	0	0	0	0%	0	0%
10-14 yrs	0	0	0	0	0	0	0	0	0%	0	0	0	0	0	0	0	0%	0	0%
15-19 yrs	0	0	0	0	0	0	0	0	0%	0	0	0	0	0	0	0	0%	0	0%
20-24 yrs	0	0	0	0	2	0	2	2	18%	0	0	0	0	0	0	0	0%	2	17%
25-29 yrs	0	0	0	1	1	0	2	2	18%	0	0	0	0	0	0	0	0%	2	17%
30-34 yrs	0	0	0	1	1	0	2	2	18%	0	0	0	0	0	0	0	0%	2	17%
35-39 yrs	0	0	0	1	0	0	1	1	9%	0	0	0	0	1	0	1	100%	2	17%
40-44 yrs	0	0	0	2	0	0	2	2	18%	0	0	0	0	0	0	0	0%	2	17%
45-54 yrs	0	0	0	1	0	0	1	1	9%	0	0	0	0	0	0	0	0%	1	8%
55-64 yrs	0	0	0	0	0	1	1	1	9%	0	0	0	0	0	0	0	0%	1	8%
65+ yrs	0	0	0	0	0	0	0	0	0%	0	0	0	0	0	0	0	0%	0	0%
Unknown	0	0	0	0	0	0	0	0	0%	0	0	0	0	0	0	0	0%	0	0%
TOTAL	0	0	0	6	4	1	11	11	100%	0	0	0	0	1	0	1	100%	12	100%

Total race/ethnicity	Number	Percent
American Indian/Alaska Native	0	0%
Asian/Pacific Islander	0	0%
Black (not Hispanic)	0	0%
White (not Hispanic)	6	50%
Hispanic	5	42%
Unknown	1	8%
TOTAL	12	100%

1-Early syphilis includes primary, secondary, and early latent stages.

Table 9. Early syphilis (<1 yr) by sex and county of residence—Idaho, 2010

County	SEX			TOTAL	Population	Rate per 100,000
	Male	Female	Unknown			
Ada	7	0	0	7	388,624	1.8
Adams	0	0	0	0	3,403	0.0
Bannock	0	0	0	0	83,548	0.0
Bear Lake	0	0	0	0	5,721	0.0
Benewah	0	0	0	0	9,270	0.0
Bingham	1	0	0	1	45,371	2.2
Blaine	0	0	0	0	22,166	0.0
Boise	0	0	0	0	7,536	0.0
Bonner	0	0	0	0	41,550	0.0
Bonneville	0	0	0	0	102,738	0.0
Boundary	0	0	0	0	11,147	0.0
Butte	0	0	0	0	2,820	0.0
Camas	0	0	0	0	1,078	0.0
Canyon	2	1	0	3	188,584	1.6
Caribou	0	0	0	0	6,936	0.0
Cassia	0	0	0	0	22,230	0.0
Clark	0	0	0	0	991	0.0
Clearwater	0	0	0	0	8,017	0.0
Custer	0	0	0	0	4,222	0.0
Elmore	0	0	0	0	28,606	0.0
Franklin	0	0	0	0	12,697	0.0
Fremont	0	0	0	0	12,693	0.0
Gem	0	0	0	0	16,188	0.0
Gooding	0	0	0	0	14,592	0.0
Idaho	0	0	0	0	15,628	0.0
Jefferson	0	0	0	0	25,126	0.0
Jerome	0	0	0	0	21,789	0.0
Kootenai	0	0	0	0	140,679	0.0
Latah	0	0	0	0	38,590	0.0
Lemhi	0	0	0	0	7,946	0.0
Lewis	0	0	0	0	3,675	0.0
Lincoln	0	0	0	0	4,679	0.0
Madison	0	0	0	0	38,550	0.0
Minidoka	0	0	0	0	19,438	0.0
Nez Perce	0	0	0	0	39,499	0.0
Oneida	0	0	0	0	4,228	0.0
Owyhee	1	0	0	1	11,129	9.0
Payette	0	0	0	0	23,140	0.0
Power	0	0	0	0	8,045	0.0
Shoshone	0	0	0	0	12,566	0.0
Teton	0	0	0	0	9,406	0.0
Twin Falls	0	0	0	0	76,386	0.0
Valley	0	0	0	0	8,416	0.0
Washington	0	0	0	0	10,153	0.0
STATE	11	1	0	12	1,559,796	0.8

Table 10. Syphilis (all stages) by sex, race/ethnicity, and age group—Idaho, 2010

Age at Diagnosis	Sex										TOTAL
	Male					Female					
	Race				Other/Unknown	Total Males	Race				TOTAL
	American Indian/Alaska Native	Asian/Pacific Islander	Black (not Hispanic)	White (not Hispanic)			American Indian/Alaska Native	Asian/Pacific Islander	Black (not Hispanic)	White (not Hispanic)	
0- 4 yrs ¹	0	0	0	0	0	0	0	0	0	1	20%
5- 9 yrs	0	0	0	0	0	0	0	0	0	0	0%
10-14 yrs	0	0	0	0	0	0	0	0	0	0	0%
15-19 yrs	0	0	0	0	0	0	0	0	0	0	0%
20-24 yrs	0	0	0	0	2	2	0	0	0	0	0%
25-29 yrs	0	0	0	1	1	2	12%	0	0	2	40%
30-34 yrs	0	0	0	1	1	2	12%	0	0	0	0%
35-39 yrs	0	0	0	1	0	2	12%	0	0	1	20%
40-44 yrs	0	0	0	2	2	4	24%	0	0	0	0%
45-54 yrs	0	0	0	2	2	4	24%	0	0	0	0%
55-64 yrs	0	0	0	0	1	1	6%	0	0	1	20%
65+ yrs	0	0	0	0	0	0	0%	0	0	0	0%
Unknown	0	0	0	0	0	0	0%	0	0	0	0%
TOTAL	0	0	0	7	8	17	100%	0	0	2	100%
										5	100%
										22	100%

1. The 1 case in the 0-4 yrs age group was congenital syphilis, syphilis acquired by an infant from the infected mother.

Total race/ethnicity	Number	Percent
American Indian/Alaska Native	0	0%
Asian/Pacific Islander	0	0%
Black (not Hispanic)	0	0%
White (not Hispanic)	9	41%
Hispanic	10	45%
Unknown	3	14%
TOTAL	22	100%

Table 11. Syphilis (all stages) by sex and county of residence—Idaho, 2010

County	SEX			TOTAL	Population	Rate per 100,000
	Male	Female	Unknown			
Ada	11	0	0	11	388,624	2.8
Adams	0	0	0	0	3,403	0.0
Bannock	0	2	0	2	83,548	2.4
Bear Lake	0	0	0	0	5,721	0.0
Benewah	0	0	0	0	9,270	0.0
Bingham	1	0	0	1	45,371	2.2
Blaine	0	0	0	0	22,166	0.0
Boise	0	0	0	0	7,536	0.0
Bonner	0	0	0	0	41,550	0.0
Bonneville	0	0	0	0	102,738	0.0
Boundary	0	0	0	0	11,147	0.0
Butte	0	0	0	0	2,820	0.0
Camas	0	0	0	0	1,078	0.0
Canyon	3	3	0	6	188,584	3.2
Caribou	0	0	0	0	6,936	0.0
Cassia	0	0	0	0	22,230	0.0
Clark	0	0	0	0	991	0.0
Clearwater	0	0	0	0	8,017	0.0
Custer	0	0	0	0	4,222	0.0
Elmore	0	0	0	0	28,606	0.0
Franklin	0	0	0	0	12,697	0.0
Fremont	0	0	0	0	12,693	0.0
Gem	0	0	0	0	16,188	0.0
Gooding	0	0	0	0	14,592	0.0
Idaho	0	0	0	0	15,628	0.0
Jefferson	0	0	0	0	25,126	0.0
Jerome	0	0	0	0	21,789	0.0
Kootenai	1	0	0	1	140,679	0.7
Latah	0	0	0	0	38,590	0.0
Lemhi	0	0	0	0	7,946	0.0
Lewis	0	0	0	0	3,675	0.0
Lincoln	0	0	0	0	4,679	0.0
Madison	0	0	0	0	38,550	0.0
Minidoka	0	0	0	0	19,438	0.0
Nez Perce	0	0	0	0	39,499	0.0
Oneida	0	0	0	0	4,228	0.0
Owyhee	1	0	0	1	11,129	9.0
Payette	0	0	0	0	23,140	0.0
Power	0	0	0	0	8,045	0.0
Shoshone	0	0	0	0	12,566	0.0
Teton	0	0	0	0	9,406	0.0
Twin Falls	0	0	0	0	76,386	0.0
Valley	0	0	0	0	8,416	0.0
Washington	0	0	0	0	10,153	0.0
STATE	17	5	0	22	1,559,796	1.4

HIV

Table 12. HIV by sex, race/ethnicity, and age group—Idaho, 2010

Age at Diagnosis	SEX										TOTAL	
	Male					Female						
	Race/ethnicity					Race/ethnicity						
American Indian/Alaska Native	Asian/Pacific Islander	Black (not Hispanic)	White (not Hispanic)	Hispanic	Unknown	Total Males	Percent of Males	American Indian/Alaska Native	Asian/Pacific Islander	Black (not Hispanic)	Total Females	
0-4 yrs	0	0	0	0	0	0	0%	0	0	0	0	
5-9 yrs	0	0	0	0	0	0	0%	0	0	0	0	
10-14 yrs	0	0	0	0	0	0	0%	0	0	0	0	
15-19 yrs	0	0	0	2	0	2	7%	0	0	0	0	
20-24 yrs	0	0	0	1	1	2	7%	0	0	0	2	
25-29 yrs	0	0	0	4	0	4	14%	0	0	0	0	
30-34 yrs	0	0	0	3	0	3	11%	0	0	0	0	
35-39 yrs	2	0	0	3	0	6	21%	0	0	0	0	
40-44 yrs	0	0	0	3	0	3	11%	0	0	1	0	
45-54 yrs	0	0	0	6	0	8	29%	0	0	1	0	
55-64 yrs	0	0	0	0	0	0	0%	0	0	0	0	
65+ yrs	0	0	0	0	0	0	0%	0	0	0	0	
Unknown	0	0	0	0	0	0	0%	0	0	0	0	
TOTAL	2	0	0	22	1	28	100%	0	0	0	4	
											100%	
											32 100%	

Total race/ethnicity	Number	Percent
American Indian/Alaska Native	2	6%
Asian/Pacific Islander	0	0%
Black (not Hispanic)	0	0%
White (not Hispanic)	26	81%
Hispanic	1	3%
Unknown	3	9%
TOTAL	32	100%

AIDS

Table 13. AIDS by sex, race/ethnicity, and age group—Idaho, 2010

Age at Diagnosis	SEX										TOTAL					
	Male					Female										
	Race/ethnicity					Race/ethnicity										
American Indian/Alaska Native	Asian/Pacific Islander	Black (not Hispanic)	White (not Hispanic)	Hispanic	Unknown	Total Males	Percent of Males	American Indian/Alaska Native	Asian/Pacific Islander	Black (not Hispanic)	White (not Hispanic)	Total Females	Percent of Females	Total Cases	Percent of Total	
0-4 yrs	0	0	0	0	0	0	0%	0	0	0	0	1	11%	1	2%	
5-9 yrs	0	0	0	0	0	0	0%	0	0	0	0	0	0	0%	0	0%
10-14 yrs	0	0	1	0	0	0	2%	0	0	0	0	0	0	0%	1	2%
15-19 yrs	0	0	0	0	1	0	2%	0	0	0	0	0	0	0%	1	2%
20-24 yrs	0	0	0	1	2	0	7%	0	0	0	0	0	0	0%	3	6%
25-29 yrs	0	0	0	3	0	1	10%	0	0	0	0	0	0	0%	4	8%
30-34 yrs	0	0	1	4	0	0	12%	0	0	1	0	0	0	11%	6	12%
35-39 yrs	0	0	0	6	3	0	21%	0	1	0	1	1	0	33%	12	24%
40-44 yrs	0	0	0	5	1	1	17%	0	0	0	1	0	0	11%	8	16%
45-54 yrs	0	0	1	6	0	1	19%	0	0	0	3	0	0	33%	11	22%
55-64 yrs	0	0	0	3	0	1	10%	0	0	0	0	0	0	0%	4	8%
65+ yrs	0	0	0	0	0	0	0%	0	0	0	0	0	0	0%	0	0%
Unknown	0	0	0	0	0	0	0%	0	0	0	0	0	0	0%	0	0%
TOTAL	0	0	3	28	7	4	100%	0	1	1	5	2	0	100%	51	100%

Total race/ethnicity	Number	Percent
American Indian/Alaska Native	0	0%
Asian/Pacific Islander	1	2%
Black (not Hispanic)	4	8%
White (not Hispanic)	33	65%
Hispanic	9	18%
Unknown	4	8%
TOTAL	51	100%

District tables

District 1

Table 14. STD, HIV, and AIDS by diagnosis and age group, Health District 1, 2010

Age group at diagnosis	Disease / Condition								
	Chlamydia	Gonorrhea	Early syphilis	Syphilis, Late/Unk/ Neuro	Congenital syphilis	HIV (not AIDS)	AIDS	TOTAL	Percent
0- 4 yrs	1	0	0	0	0	0	0	1	0%
5- 9 yrs	1	0	0	0	0	0	0	1	0%
10-14 yrs	4	0	0	0	0	0	0	4	1%
15-19 yrs	222	3	0	0	0	0	0	225	35%
20-24 yrs	238	7	0	0	0	3	0	248	38%
25-29 yrs	94	3	0	0	0	0	1	98	15%
30-34 yrs	44	0	0	0	0	0	0	44	7%
35-39 yrs	13	0	0	0	0	0	1	14	2%
40-44 yrs	9	0	0	0	0	0	1	10	2%
45-54 yrs	2	0	0	1	0	0	2	5	1%
55-64 yrs	0	0	0	0	0	0	0	0	0%
65+ yrs	0	0	0	0	0	0	0	0	0%
Unknown	0	0	0	0	0	0	0	0	0%
TOTAL	628	13	0	1	0	3	5	650	100%

Table 15. STD, HIV, and AIDS by year of report, Health District 1, 2001–2010

Year of report	Chancroid	Chlamydia	Gonorrhea	Early syphilis	Syphilis, Late/Unk/ Neuro	Congenital syphilis	HIV (not AIDS)	AIDS
2001	0	276	15	0	0	0	2	0
2002	0	292	10	0	1	0	3	6
2003	0	320	11	0	1	0	1	6
2004	1	348	11	0	3	0	2	3
2005	0	294	8	0	1	0	0	4
2006	0	417	9	1	0	0	4	1
2007	0	479	40	1	3	0	2	3
2008	0	608	39	1	0	0	0	3
2009	0	604	22	0	1	0	3	2
2010	0	628	13	0	1	0	3	5

District 2

Table 16. STD, HIV, and AIDS by diagnosis and age group, Health District 2, 2010

Age group at diagnosis	Disease / Condition								TOTAL	Percent
	Chlamydia	Gonorrhea	Early syphilis	Syphilis, Late/Unk/ Neuro	Congenital syphilis	HIV (not AIDS)	AIDS			
0- 4 yrs	1	0	0	0	0	0	0	0	1	0%
5- 9 yrs	0	0	0	0	0	0	0	0	0	0%
10-14 yrs	3	0	0	0	0	0	0	0	3	1%
15-19 yrs	94	5	0	0	0	0	0	0	99	29%
20-24 yrs	134	7	0	0	0	0	0	0	141	41%
25-29 yrs	49	4	0	0	0	0	0	0	53	15%
30-34 yrs	17	4	0	0	0	0	0	1	22	6%
35-39 yrs	12	3	0	0	0	0	0	0	15	4%
40-44 yrs	1	1	0	0	0	1	2	0	5	1%
45-54 yrs	1	0	0	0	0	2	0	0	3	1%
55-64 yrs	0	0	0	0	0	0	0	0	0	0%
65+ yrs	0	0	0	0	0	0	0	0	0	0%
Unknown	0	0	0	0	0	0	0	0	0	0%
TOTAL	312	24	0	0	0	3	3	342	100%	

Table 17. STD, HIV, and AIDS by year of report, Health District 2, 2001–2010

Year of report	Chancroid	Chlamydia	Gonorrhea	Early syphilis	Syphilis, Late/Unk/ Neuro	Congenital syphilis	HIV (not AIDS)	AIDS
2001	0	88	2	0	0	0	0	4
2002	0	196	13	1	0	0	2	2
2003	0	203	11	1	0	0	3	1
2004	0	269	7	0	3	0	0	1
2005	0	202	3	0	1	0	3	1
2006	0	183	13	0	3	0	3	2
2007	0	196	14	0	1	0	0	1
2008	0	261	20	1	0	0	1	2
2009	0	271	7	0	2	0	2	5
2010	0	312	24	0	0	0	3	3

District 3

Table 18. STD, HIV, and AIDS by diagnosis and age group, Health District 3, 2010

Age group at diagnosis	Disease / Condition								
	Chlamydia	Gonorrhea	Early syphilis	Syphilis, Late/Unk/ Neuro	Congenital syphilis	HIV (not AIDS)	AIDS	TOTAL	Percent
0- 4 yrs	2	0	0	0	1	0	0	3	0%
5- 9 yrs	0	0	0	0	0	0	0	0	0%
10-14 yrs	7	0	0	0	0	0	0	7	1%
15-19 yrs	257	6	0	0	0	0	0	263	30%
20-24 yrs	344	21	2	0	0	0	0	367	42%
25-29 yrs	122	4	0	1	0	0	1	128	15%
30-34 yrs	48	2	0	0	0	0	0	50	6%
35-39 yrs	19	0	2	0	0	0	4	25	3%
40-44 yrs	7	1	0	0	0	1	2	11	1%
45-54 yrs	6	0	0	1	0	1	2	10	1%
55-64 yrs	2	0	0	0	0	0	2	4	0%
65+ yrs	1	0	0	0	0	0	0	1	0%
Unknown	0	0	0	0	0	0	0	0	0%
TOTAL	815	34	4	2	1	2	11	869	100%

Table 19. STD, HIV, and AIDS by year of report, Health District 3, 2001–2010

Year of report	Chancroid	Chlamydia	Gonorrhea	Early syphilis	Syphilis, Late/Unk/ Neuro	Congenital syphilis	HIV (not AIDS)	AIDS
2001	0	302	14	0	1	0	3	0
2002	0	468	22	6	3	0	1	4
2003	0	433	11	16	4	4	5	2
2004	0	481	20	34	16	2	4	4
2005	0	617	35	24	9	1	1	4
2006	0	693	43	4	1	0	4	10
2007	0	829	59	0	1	0	0	0
2008	0	787	30	2	1	0	5	2
2009	0	814	20	1	6	1	3	3
2010	0	815	34	4	2	1	2	11

District 4

Table 20. STD, HIV, and AIDS by diagnosis and age group, Health District 4, 2010

Age group at diagnosis	Disease / Condition								
	Chlamydia	Gonorrhea	Early syphilis	Syphilis, Late/Unk/Neuro	Congenital syphilis	HIV (not AIDS)	AIDS	TOTAL	Percent
0- 4 yrs	1	0	0	0	0	0	0	1	0%
5- 9 yrs	0	0	0	0	0	0	0	0	0%
10-14 yrs	5	0	0	0	0	0	1	6	0%
15-19 yrs	364	4	0	0	0	2	0	370	26%
20-24 yrs	578	21	0	0	0	1	2	602	42%
25-29 yrs	216	18	1	0	0	2	2	239	17%
30-34 yrs	94	5	2	0	0	2	4	107	7%
35-39 yrs	45	2	0	1	0	1	2	51	4%
40-44 yrs	24	1	2	2	0	1	2	32	2%
45-54 yrs	15	0	1	1	0	5	6	28	2%
55-64 yrs	1	1	1	0	0	0	2	5	0%
65+ yrs	1	0	0	0	0	0	0	1	0%
Unknown	0	0	0	0	0	0	0	0	0%
TOTAL	1,344	52	7	4	0	14	21	1,442	100%

Table 21. STD, HIV, and AIDS by year of report, Health District 4, 2001–2010

Year of report	Chancroid	Chlamydia	Gonorrhea	Early syphilis	Syphilis, Late/Unk/Neuro	Congenital syphilis	HIV (not AIDS)	AIDS
2001	0	813	28	1	5	0	10	10
2002	0	940	26	3	4	0	11	7
2003	0	814	23	1	7	0	8	4
2004	0	968	39	12	1	0	6	4
2005	0	1,057	53	3	4	0	19	11
2006	0	1,264	107	0	0	0	11	9
2007	0	1,389	109	0	4	0	14	5
2008	0	1,528	55	4	7	0	7	9
2009	0	1,221	30	3	12	0	17	15
2010	0	1,344	52	7	4	0	14	21

District 5

Table 22. STD, HIV, and AIDS by diagnosis and age group, Health District 5, 2010

Age group at diagnosis	Disease / Condition								
	Chlamydia	Gonorrhea	Early syphilis	Syphilis, Late/Unk/ Neuro	Congenital syphilis	HIV (not AIDS)	AIDS	TOTAL	Percent
0- 4 yrs	2	0	0	0	0	0	0	2	0%
5- 9 yrs	0	0	0	0	0	0	0	0	0%
10-14 yrs	3	0	0	0	0	0	0	3	1%
15-19 yrs	135	1	0	0	0	0	0	136	33%
20-24 yrs	152	5	0	0	0	0	0	157	38%
25-29 yrs	74	1	0	0	0	0	0	75	18%
30-34 yrs	19	0	0	0	0	1	0	20	5%
35-39 yrs	9	0	0	0	0	0	1	10	2%
40-44 yrs	6	1	0	0	0	0	0	7	2%
45-54 yrs	5	0	0	0	0	0	1	6	1%
55-64 yrs	0	0	0	0	0	0	0	0	0%
65+ yrs	1	0	0	0	0	0	0	1	0%
Unknown	0	0	0	0	0	0	0	0	0%
TOTAL	406	8	0	0	0	1	2	417	100%

Table 23. STD, HIV, and AIDS by year of report, Health District 5, 2001–2010

Year of report	Chancroid	Chlamydia	Gonorrhea	Early syphilis	Syphilis, Late/Unk/ Neuro	Congenital syphilis	HIV (not AIDS)	AIDS
2001	0	227	6	0	4	0	0	2
2002	0	287	5	0	3	0	3	3
2003	0	252	7	3	4	0	1	3
2004	0	314	10	2	0	0	4	0
2005	0	278	8	5	5	0	2	1
2006	0	323	25	1	0	0	0	1
2007	0	353	38	1	0	0	2	0
2008	0	461	14	3	0	0	0	7
2009	0	346	14	1	0	0	3	4
2010	0	406	8	0	0	0	1	2

District 6

Table 24. STD, HIV, and AIDS by diagnosis and age group, Health District 6, 2010

Age group at diagnosis	Disease / Condition								
	Chlamydia	Gonorrhea	Early syphilis	Syphilis, Late/Unk/ Neuro	Congenital syphilis	HIV (not AIDS)	AIDS	TOTAL	Percent
0- 4 yrs	2	0	0	0	0	0	0	2	0%
5- 9 yrs	0	0	0	0	0	0	0	0	0%
10-14 yrs	1	1	0	0	0	0	0	2	0%
15-19 yrs	125	0	0	0	0	0	0	125	29%
20-24 yrs	149	0	0	0	0	0	1	150	35%
25-29 yrs	77	2	1	1	0	0	0	81	19%
30-34 yrs	36	1	0	0	0	0	0	37	9%
35-39 yrs	9	0	0	0	0	2	2	13	3%
40-44 yrs	8	1	0	0	0	0	0	9	2%
45-54 yrs	4	0	0	0	0	1	0	5	1%
55-64 yrs	0	0	0	1	0	0	0	1	0%
65+ yrs	0	0	0	0	0	0	0	0	0%
Unknown	0	0	0	0	0	0	0	0	0%
TOTAL	411	5	1	2	0	3	3	425	100%

Table 25. STD, HIV, and AIDS by year of report, Health District 6, 2001–2010

Year of report	Chancroid	Chlamydia	Gonorrhea	Early syphilis	Syphilis, Late/Unk/ Neuro	Congenital syphilis	HIV (not AIDS)	AIDS
2001	0	175	5	0	0	0	3	0
2002	0	180	10	2	0	0	4	2
2003	0	184	2	1	1	0	1	2
2004	1	275	14	1	0	0	1	2
2005	0	241	8	1	1	0	3	1
2006	0	299	5	0	0	0	1	5
2007	0	305	3	1	0	0	4	3
2008	0	317	22	2	5	0	18	2
2009	0	348	10	1	1	0	8	2
2010	0	411	5	1	2	0	3	3

District 7

Table 26. STD, HIV, and AIDS by diagnosis and age group, Health District 7, 2010

Age group at diagnosis	Disease / Condition								
	Chlamydia	Gonorrhea	Early syphilis	Syphilis, Late/Unk/ Neuro	Congenital syphilis	HIV (not AIDS)	AIDS	TOTAL	Percent
0- 4 yrs	0	0	0	0	0	0	1	1	0%
5- 9 yrs	0	0	0	0	0	0	0	0	0%
10-14 yrs	1	0	0	0	0	0	0	1	0%
15-19 yrs	89	2	0	0	0	0	1	92	29%
20-24 yrs	124	7	0	0	0	0	0	131	42%
25-29 yrs	42	2	0	0	0	2	0	46	15%
30-34 yrs	24	0	0	0	0	0	1	25	8%
35-39 yrs	7	0	0	0	0	3	2	12	4%
40-44 yrs	1	0	0	0	0	1	1	3	1%
45-54 yrs	2	0	0	0	0	0	0	2	1%
55-64 yrs	1	0	0	0	0	0	0	1	0%
65+ yrs	0	0	0	0	0	0	0	0	0%
Unknown	1	0	0	0	0	0	0	1	0%
TOTAL	292	11	0	0	0	6	6	315	100%

Table 27. STD, HIV, and AIDS by year of report, Health District 7, 2001–2010

Year of report	Chancroid	Chlamydia	Gonorrhea	Early syphilis	Syphilis, Late/Unk/ Neuro	Congenital syphilis	HIV (not AIDS)	AIDS
2001	0	143	6	0	0	0	1	1
2002	0	172	10	0	0	0	1	1
2003	0	160	3	1	1	0	1	2
2004	0	129	2	1	2	0	1	1
2005	0	110	4	0	0	0	0	0
2006	0	166	4	0	2	0	1	1
2007	0	171	6	1	1	0	2	0
2008	0	232	7	0	0	0	3	4
2009	0	238	7	0	2	0	5	3
2010	0	292	11	0	0	0	6	6

