



Results from the 2008  
Behavioral Risk Factor  
Surveillance System

# Idaho Behavioral Risk Factors

Bureau of Vital Records  
and Health Statistics  
Idaho Department of  
Health and Welfare

October 2009

## Acknowledgments

This project was funded in part by a grant from the Centers for Disease Control and Prevention, number U58/CCU022801, and administered via a cooperative agreement with the Idaho Bureau of Vital Records and Health Statistics. The contents are solely the responsibility of the authors and do not necessarily represent the official views of the Centers for Disease Control and Prevention.

Special thanks go to Jeff Harder of Boise, Idaho, for his excellent artwork and his design of the cover.

The Idaho Department of Health and Welfare would like to thank the citizens of Idaho who have participated in the Behavioral Risk Factor Surveillance System.

For more details on this project or any of the survey results, please contact the Bureau of Vital Records and Health Statistics at (208) 332-7326.

## Suggested Citation

*Idaho Behavioral Risk Factors: Results From the 2008 Behavioral Risk Factor Surveillance System.* Boise: Idaho Department of Health and Welfare, Division of Health, Bureau of Vital Records and Health Statistics, 2009.

# Table of Contents

---

Title Page.....	i
Acknowledgments.....	ii
Table of Contents.....	iii
Map of Idaho Public Health Districts .....	iv
Introduction .....	1
Methodology .....	1
Reporting .....	2
Data Limitations .....	2
Summary Results.....	3
2008 Risk Factors: Idaho and United States Median .....	8
General Health	
General Health Status .....	10
Health Care Coverage .....	12
Dental Insurance .....	14
Chronic Disease Prevalence	
Diabetes .....	16
Asthma .....	18
Health Risk Behaviors	
Physical Activity .....	20
Weight .....	22
Cigarette Smoking.....	26
Alcohol Consumption .....	28
Illicit Drug Use .....	32
Preventive Behaviors	
Seat Belt Use .....	34
Colorectal Cancer Screening .....	36
Breast Cancer Screening .....	38
Cervical Cancer Screening.....	40
Prostate Cancer Screening .....	42
Dental Visits .....	44
Data Tables	
Understanding the Data Tables.....	46
2008 Data Tables by Topic .....	48
Appendix A: Trends of Selected Risk Variables by Public Health District .....	85
Appendix B: 2008 BRFSS Questionnaire (with results) .....	95
Appendix C: References .....	110

# Idaho Public Health Districts


---

Idaho is divided into seven public health districts, with each of the districts containing four to eight counties. The seven health districts are independent agencies governed by a local board appointed by the county commissioners within the district.

## 2008 Estimated Population Adults Aged 18 and Over

Overall:.....	1,088,573
District 1: .....	161,320
District 2: .....	80,560
District 3: .....	170,252
District 4: .....	302,303
District 5: .....	126,852
District 6: .....	116,665
District 7: .....	130,621

*Note:* Population estimates are obtained from the Centers for Disease Control and Prevention and are used in weighting BRFSS data (see page 2). These population estimates may differ from U.S. Census Bureau population figures.



# Idaho Behavioral Risk Factor Surveillance System

---

## Introduction

Health risk behaviors and lack of preventive care are associated with higher rates of morbidity and mortality. Monitoring of modifiable risk factors for diseases and the use of preventive health services is vital to planning, initiating, and developing targeted education and intervention programs at the state and local level. It also provides a means of tracking progress toward achieving public health objectives. The Behavioral Risk Factor Surveillance System (BRFSS) is an ongoing surveillance program developed and partially funded by the Centers for Disease Control and Prevention (CDC). It is designed to estimate the prevalence of risk factors for the major causes of morbidity and mortality in the United States. The survey provides state-specific estimates of the proportion of adults aged 18 and over reporting health risk behaviors. For certain state and national objectives, BRFSS is the only source of data.

Idaho has participated in the BRFSS since its inception in 1984. Since then, the program has grown to encompass all 50 states and several U.S. territories. BRFSS data are valued at the state and national level for measuring progress toward national and state health objectives. BRFSS is unique in providing data that enable state health policymakers to assess their own states' needs and determine progress toward goals. Results from BRFSS have been used to support risk reduction and disease prevention activities by directing program planning, assessing trends, and targeting relevant population groups. This report summarizes major health risk behaviors and use of preventive health practices that affect chronic diseases associated with morbidity and mortality in Idaho. Summarized data in this report are not inclusive of all data that may be provided by BRFSS. For this reason, the 2008 questionnaire appears in its entirety in Appendix B.

## Methodology

The BRFSS is conducted as a random telephone survey of the non-institutionalized adult population. In order to produce health district estimates, Idaho's sample has grown in size from 600 people in 1984 to over 5,000 in 2008. The survey is administered in every month of the calendar year. After annual data collection is complete, individual responses are weighted to be representative of the state's adult population, and analysis is performed on the weighted data.

Additional information regarding BRFSS methodology is available online at <http://www.cdc.gov/brfss>.

*Sampling:* Idaho used disproportionate stratified sampling for its 2008 BRFSS. The sample was stratified by Idaho's seven public health districts. A minimum of 700 Idahoans were interviewed within each health district, for a total sample size

of 5,111. Interviews were conducted by telephone using computer-assisted telephone interviewing (CATI) software to record responses.

*Weighting:* The data were weighted to the respondent's probability of selection as well as age and sex based on 2008 population estimates. For example, some households have more than one telephone line and are more likely to be called. The weighting corrects for this difference in probability of selection as well as differences between the population and the sample.

*Analysis:* Idaho used SUDAAN<sup>®</sup> (Survey Data Analysis) software for statistical testing and calculation of confidence limits. This software takes into account the complex sampling design methodology of BRFSS. Idaho used SAS<sup>®</sup> software for data manipulation and risk factor creation. All "don't know," "not sure," and "refused" responses have been excluded from the analysis.

## **Reporting**

Prevalence estimates based on denominators with fewer than 50 respondents have been suppressed and are indicated in the data tables with an asterisk (\*). The BRFSS has adopted this standard to maintain a high degree of reliability. Differences between estimates are reported throughout the document. Any difference determined to be statistically significantly different (using 95 percent confidence intervals) was designated as such and will be preceded by the words "significantly" or "statistically."

## **Data Limitations**

Errors in estimation are a result of BRFSS data being self-reported and certain behaviors possibly being underreported (Centers for Disease Control and Prevention, 2008). Another source of error is based on sampling as samples will deviate somewhat from the population. Additional possible errors may occur due to the population from which the sample is drawn. In order to be cost effective, the sample is limited to adults aged 18 and older who are non-institutionalized, live in a household with a non-cellular (landline) telephone, and can communicate in English or Spanish. This excludes people in prisons and dormitories, those who exclusively speak a language other than English or Spanish, those with only cellular telephones and others who cannot communicate by telephone. Spanish interviewing for the Idaho BRFSS began in midyear 2004 with 2005 as the first complete year that included Spanish interviewing. The Federal Communications Commission estimates that 95.5 percent of households had landline telephone service in 2006 (Federal Communications Commission, 2009). With ongoing changes in telephone technology, there has been an increase in households that have cellular telephones only. These households are not in the current sampling frame for the BRFSS which may bias results (Centers for Disease Control and Prevention, 2007).

Additional information on telephone coverage in U.S. households is available at: [www.fcc.gov/wcb/iatd/socc.html](http://www.fcc.gov/wcb/iatd/socc.html)

# Summary Results

---

## General Health Status (page 10)

- The percentage of Idaho adults reporting their health as “fair” or “poor” held steady at 14.0 percent in 2008. This figure has not changed significantly over the past ten years.
- Those with household incomes below \$25,000 were three times as likely to say their general health was less than good compared with those with incomes of \$25,000 or higher.
- Married adults were less likely to report “fair” or “poor” general health compared with unmarried adults (which includes those who were divorced, separated, widowed and never married).

## Health Care Coverage (page 12)

- In Idaho, 18.5 percent of adults did not have health care coverage in 2008. This was not a significant change since 2007.
- Health care coverage increased with age. More than one-in-three of those aged 18 to 24 were without coverage. This compares with 7.3 percent of those over the age of 55.
- Hispanics were significantly more likely than non-Hispanics to be without health coverage.

## Dental Insurance (page 14)

- The percentage of Idaho adults who were without dental insurance was unchanged since 2007 at 43.0 percent.
- Those over age 65 were more likely than younger adults to be without dental insurance.
- Hispanics were significantly more likely than non-Hispanics to be without dental insurance.

## Diabetes (page 16)

- Adult diabetes prevalence in 2008 was 7.0 percent. The percentage of Idahoans with diabetes has increased significantly since 1999.
- Those who did not engage in leisure time physical activity reported diabetes twice as often as those who did.

# Summary Results

---

## Asthma (page 18)

- Asthma prevalence (those with current asthma) was 8.9 percent in 2008 and has not changed significantly during the ten year tracking period.
- There were no significant differences in asthma prevalence between males and females, or by level of income, education or employment status.
- Compared with Hispanics, non-Hispanics were significantly more likely to have a current diagnosis of asthma.

## Physical Activity (page 20)

- In Idaho, 21.1 percent of adults did not participate in leisure time physical activity in 2008. This has not changed significantly over the past 10 years.
- Overweight (BMI  $\geq 25$ ) adults were significantly less likely to report physical activity when compared with those who were not overweight.

## Weight (page 22)

- About 6-in-10 (62.2 percent) Idaho adults were overweight (BMI  $\geq 25$ ) in 2008. While this was not a significant change since 2007, the prevalence of overweight adults has increased significantly since 1999.
- About one-in-four (25.2 percent) Idahoans was obese (BMI  $\geq 30$ ) in 2008. The proportion of overweight adults who are also obese has increased significantly over the past ten years.
- College graduates were significantly less likely than those with lower levels of education to be obese.
- Idaho adults categorized as overweight and/or obese were significantly more likely to have diabetes.

# Summary Results

---

## **Cigarette Smoking (page 26)**

- The percentage of adults who were cigarette smokers dropped slightly in 2008 to 16.9 percent. Cigarette smoking in Idaho has decreased significantly since 1999.
- Cigarette smokers were significantly more likely to have “fair” or “poor” general health.
- Smoking prevalence decreased as education level increased. Those with less than a high school education were five times as likely to smoke as college graduates.

## **Alcohol Consumption (page 28)**

- Binge drinking for females is defined as having four or more alcoholic beverages on a single occasion. For males, binge drinking is defined as having five or more alcoholic beverages on a single occasion. In 2008, 13.1 percent of Idaho adults engaged in binge drinking in the past 30 days. This has not changed significantly since 2005.
- Heavy drinking (more than 30 drinks per month for females and more than 60 drinks for males) showed no significant change from recent years with 4.6 percent of adults reporting heavy drinking.
- In every age category, males were significantly more likely to be binge drinkers than females.
- Males and females were not significantly different with regard to heavy drinking.
- Binge drinkers and heavy drinkers were significantly more likely to smoke cigarettes compared with those who were not binge or heavy drinkers.

# Summary Results

---

## **Illicit Drug Use (page 32)**

- The percentage of Idaho adults who used illicit drugs for non-medical reasons in the prior 12 months remained steady at 4.1 percent in 2008.
- Those aged 18 to 34 were significantly more likely to have used illicit drugs compared with older adults.
- In 2008, illicit drug use did not vary significantly by employment status or education.

## **Seat Belt Use (page 34)**

- The percentage of Idaho adults who did not always wear a seat belt when driving or riding in a car was 23.4 percent in 2008.
- Males were significantly more likely to say they did not consistently wear their seat belts compared with females.
- Those over the age of 65 were significantly more likely use their seat belts than those under the age of 25.

## **Colorectal Cancer Screening (page 36)**

- The percentage of those aged 50 and over who had never received a sigmoidoscopy or colonoscopy reached a new low in 2008 at 44.4 percent.
- Idaho adults over the age of 65 were more likely to have received sigmoidoscopy or colonoscopy compared with those aged 50 to 64.

## **Breast Cancer Screening (page 38)**

- In 2008, 36.8 percent of women aged 40 and over had not received a mammogram and clinical breast exam in the prior two years. This has not varied significantly over the ten-year tracking period.
- Women aged 40 and over with health insurance were twice as likely to receive these recommended breast cancer screenings compared with those who did not have health insurance.

# Summary Results

---

## **Cervical Cancer Screening (page 40)**

- In 2008, 23.1 percent of Idaho women with an intact cervix had not received a Pap test in the prior three years. This was not a significant change since 2006 but is significantly higher than the decade low in 2002.
- Women in the youngest (18 to 24 years) and oldest (65 and older) age categories were the most likely to have not received cervical cancer screening.

## **Prostate Cancer Screening (page 42)**

- The percentage of males 40 and over who had not received a Prostate-Specific Antigen (PSA) test in the prior two years did not change significantly between 2001 and 2008. Just under one-half (47.1 percent) of males 40 and older had not received a PSA in 2006.
- Males aged 40 and over without health care coverage were significantly less likely to receive a PSA test compared with those who had health insurance.

## **Dental Visits (page 44)**

- In 2008, 32.0 percent of Idaho adults had not visited a dentist within the previous 12 months.
- The percentage of adults who had annual dental visits did not vary significantly by age or by sex.
- Those who did not have dental insurance were significantly less likely to visit a dentist in the past year compared with those who had dental insurance.

## 2008 Risk Factors: Idaho and United States Median\*

---

Percent of adults who:	Idaho %	Median of U.S. States*
Reported their general health as "fair" or "poor"	14.0	14.4
Did not have health care coverage	18.5	14.5
Did not have dental insurance	43.0	n/a
Had ever been told they had diabetes	7.0	8.3
Were diagnosed and still had asthma	8.9	8.8
Did not participate in leisure time physical activity	21.1	24.6
Were overweight (BMI $\geq$ 25)	62.1	63.2
Were obese (BMI $\geq$ 30)	25.2	26.7
Smoked cigarettes	16.9	18.4
Were binge drinkers ( $\geq$ 4 alcoholic drinks on single occasion for females, $\geq$ 5 alcoholic drinks on single occasion for males)	13.1	15.6
Were heavy drinkers ( >30 alcoholic drinks in past 30 days for females, >60 alcoholic drinks in past 30 days for males)	4.6	5.1

\* Median of all states and District of Columbia (51)

n/a = data not available

## 2008 Risk Factors: Idaho and United States Median\*

---

Percent of adults who:	Idaho %	Median of U.S. States*
Used illicit drugs in the last 12 months	4.1	n/a
Did not always wear a seatbelt when driving or riding	23.4	n/a
Had not visited the dentist in the last 12 months	32.0	28.7
<i>Adults aged 50+ who:</i> Had never had a sigmoidoscopy or colonoscopy	44.4	37.8
<i>Females aged 40+ who:</i> Had not had a mammogram and clinical breast exam in last two years	36.8	n/a
<i>Females aged 18+ who:</i> Had not had a Pap test in the last three years	23.1	17.1
<i>Males aged 40+ who:</i> Had not had a Prostate Specific Antigen test in the last two years	47.1	45.2

\* Median of all states and District of Columbia (51)


n/a = data not available

# General Health

## General Health Status


Respondents were asked to rate their health as “excellent,” “very good,” “good,” “fair,” or “poor.” General health status remained similar to that of recent years with 14.0 percent of Idaho adults rating their health as “fair” or “poor.” The slight increase since 1999 was not statistically significant.

**Percent of Idaho adults who reported their general health status as "fair" or "poor" 1999-2008**



In 2008, “fair” or “poor” general health was highest in Districts 1, 2 and 3, each with 16.5 percent. District 4 had the lowest percentage of “fair” or “poor” health, at 11.5 percent.

**Percent of Idaho adults who reported their general health status as "fair" or "poor," 2008 Health District**




# General Health

## General Health Status

There were no significant differences between males and females in general health status. While females aged 18 to 24 appeared to be more likely to report “fair” or “poor” health compared with males in the same age group, the difference was not statistically significant. Those over the age of 55 were significantly more likely to report “fair” or “poor” health compared with those aged 54 and younger.

**Percent of Idaho adults who reported their general health status as "fair" or "poor," 2008  
Age and Sex**



## Other Highlights:


- Those with annual incomes below \$25,000 were three times as likely to say their general health was less than good compared with those with incomes of \$25,000 or above (27.1 percent compared with 9.0 percent).
- Adults who were employed were significantly less likely to report “fair” or “poor” health (8.0 percent) when compared with other employment categories (19.6 percent of the unemployed and 24.0 percent of “other” employment status, which includes students, homemakers, retirees and those unable to work).
- Those with higher levels of education were less likely to report “fair” or “poor” health status. More than one-quarter (26.7 percent) of those without a high school diploma reported having “fair” or “poor” health compared with 7.5 percent of those with a college degree.
- Married adults were less likely to report less than “good” general health compared with unmarried adults (12.6 percent compared with 17.2 percent). “Unmarried adults” refers to those who were divorced, separated, widowed, or never married as well as cohabitating couples.

# General Health

## Health Care Coverage


In 2008, 18.5 percent of adults in Idaho had no health care coverage. The percentage of uninsured adults has remained steady since 2005.

**Percent of Idaho adults without health care coverage 1999-2008**



Idaho adults living in Health District 6 and District 7 were the least likely to report having no health care coverage in 2008 (14.2 percent and 14.6 percent, respectively). District 3 had the highest percentage of uninsured adults at 25.4 percent.

**Percent of Idaho adults without health care coverage, 2008 Health District**




# General Health

## Health Care Coverage

Overall, males were significantly more likely to be without health care coverage than females (21.0 percent compared with 16.1 percent). However, when factoring in age it becomes apparent that the gender difference is only notable for those under the age of 35. Adults 35 and older of both sexes were significantly more likely to have health coverage.

**Percent of Idaho adults without health care coverage, 2008**  
Age and Sex



### Other Highlights:

- Idahoans with household incomes below \$25,000 were more likely than those with higher incomes to be without insurance coverage. More than one-in-three (36.3%) of those in households with incomes below \$25,000 were uninsured. This compares with just 5.3 percent of those with incomes of \$50,000 or higher.
- Unemployed adults were nearly three times as likely as employed adults to be uninsured (56.9 percent versus 19.2 percent). Still, the majority (63.4 percent) of those without health care coverage were employed for wages.
- Idaho adults with a high school education or below reported being uninsured more than two-and-a-half times as often as those with some college or more (29.7 percent compared with 11.5 percent).
- Hispanics were significantly more likely than non-Hispanics to be without health coverage (44.4 percent compared with 16.4 percent).


# General Health

## Dental Insurance

In 2008, 43.0 percent of Idaho adults did not have dental insurance. There has been no significant change in this figure during the ten year tracking period.

**Percent of Idaho adults without dental insurance, 1999-2008**


Breaks in the trendline indicate data are not available for those years



Compared with the rest of the state, Health District 4 had a significantly lower percentage of adults without dental insurance. None of the other districts varied significantly with regard to dental insurance coverage.

**Percent of Idaho adults without dental insurance, 2008**

Health District




# General Health

## Dental Insurance

Although males and females were equally likely to be without dental insurance coverage, differences were observed between age categories. Those over the age of 65 were significantly more likely than younger adults to be without dental insurance at 64.8 percent. Still, nearly one-half (48.2 percent) of those aged 18 to 24 did not have dental insurance.

**Percent of Idaho adults without dental insurance, 2008**  
**Age and Sex**



### Other Highlights:


- Nine-in-ten (90.0 percent) Idaho adults without health care coverage were also without dental insurance.
- Unemployed adults were significantly more likely to be without dental insurance compared with employed adults (67.6 percent compared with 35.5 percent, respectively). Still, most (50.8 percent) of those without dental insurance were employed.
- Dental insurance coverage varied significantly by income level. About two-thirds (66.3 percent) of Idaho adults with incomes below \$25,000 did not have dental insurance. This compares with about one-in-four (25.3 percent) of those with incomes of \$50,000 or more.
- Those without a high school diploma were more than twice as likely as college graduates to be without dental insurance (67.6 percent compared with 30.6 percent, respectively).

# Chronic Disease Prevalence

## Diabetes


The percentage of Idaho adults with diabetes was 7.0 percent in 2008 which was not a significant change from the prior year. However, the prevalence of diabetes has increased significantly since 1999 (from 4.8 percent to 7.0 percent).

**Percent of Idaho adults who had ever been told they had diabetes 1998-2008**



Diabetes prevalence ranged from a high of 8.9 percent in District 6 to a low of 6.0 percent in District 4. None of the differences between Health Districts were statistically significant.

**Percent of Idaho adults who had ever been told they had diabetes, 2008 Health District**




# Chronic Disease Prevalence

## Diabetes

Males and females did not significantly differ overall with regard to diabetes prevalence (7.4 percent for males and 6.7 percent for females). Diabetes prevalence increased with age with nearly one-in-five (19.2 percent) of those over age 65 being diagnosed with diabetes.

**Percent of Idaho adults who had ever been told they had diabetes, 2008**  
Age and Sex



### Other Highlights:

- Obese (BMI  $\geq 30$ ) adults reported diabetes more than five times as often as those who were not overweight (BMI  $< 25$ ). Among obese adults, 15.4 percent had diabetes compared with 2.9 percent of those who were not overweight or obese.
- Those who did not engage in leisure time physical activity reported diabetes at twice the rate of those who did have leisure time physical activity (11.6 percent compared with 5.8 percent).
- Diabetes prevalence did not differ significantly by education level in 2008. Among those with a high school diploma or less education, 7.6 percent had diabetes. This was not significantly higher than for those with some college or more education (6.7 percent of whom had diabetes).
- Those with lower incomes were significantly more likely to have diabetes than those with mid-level or high incomes. Nearly one-in-ten (9.7 percent) of those with household incomes below \$35,000 had diabetes compared with 5.0 percent of those with incomes of \$35,000 or more.


# Chronic Disease Prevalence

## Asthma

Prior to 2002, the Idaho BRFSS reported asthma prevalence as those who had ever been told they had asthma. Since 2002, Idaho began reporting asthma prevalence as those adults who had ever been told they had asthma and still had it. In 2008, the percentage of adults with current asthma remained steady at 8.9 percent.


**Percent of Idaho adults who were diagnosed and still had asthma 1999-2008**

Breaks in the trendline indicate data are not available for those years



Asthma prevalence was highest in Health District 2 at 11.5 percent. This was significantly higher than for District 7 (6.0 percent). No other district differed significantly on asthma prevalence.

**Percent of Idaho adults who were diagnosed and still had asthma, 2008 Health District**




# Chronic Disease Prevalence

## Asthma

In 2008, males and females did not differ significantly on asthma prevalence (10.2 percent compared with 7.6 percent, overall). However, females aged 45 to 64 were significantly more likely than males of the same age to have asthma (13.2 percent and 7.7 percent, respectively).

**Percent of Idaho adults who were diagnosed and still had asthma, 2008**  
**Age and Sex**



### Other Highlights:

- There were no significant differences in asthma prevalence by level of income, education or employment status.
- Adults with current asthma reported their general health status as “fair” or “poor” more than twice as often as those who did not have asthma (28.5 percent compared with 12.4 percent).
- Non-Hispanics were significantly more likely to have a current diagnosis of asthma compared with Hispanics (9.4 percent compared with 2.0 percent).


# Health Risk Behaviors

## Physical Activity

More than one-in-five (21.1 percent) Idaho adults did not participate in physical activity in 2008. There were no significant changes in the ten year trend of leisure time physical activity.

**Percent of Idaho adults who did not participate in leisure time physical activity, 1999-2008**


Breaks in the trendline indicate data are not available for those years



Idaho adults residing in District 4 were significantly less likely than those in the rest of the state to say they did not participate in physical activity. No other district differed significantly from the state as a whole.

**Percent of Idaho adults who did not participate in leisure time physical activity, 2008**

Health District




# Health Risk Behaviors

## Physical Activity

No significant difference existed between male and female participation in leisure time physical activity in 2008. Adults over 55 were significantly more likely to say they did not participate in physical activity compared with those 54 and younger (28.8 percent compared with 17.8 percent).

**Percent of Idaho adults who did not participate in leisure time physical activity, 2008  
Age and Sex**



### Other Highlights:

- Participation in physical activity varied significantly by income level. One-in-three (33.3 percent) of those with household incomes below \$25,000 reported no physical activity compared with just over one-in-ten (11.5 percent) of those with incomes of \$50,000 or higher.
- Adults with a K to 11<sup>th</sup> grade education reported no leisure time physical activity three times as often as those with a college degree (36.5 percent compared with 11.8 percent).
- Overweight (BMI  $\geq$  25) adults were significantly more likely to report no leisure time physical activity when compared with those who were not overweight (23.3 percent compared with 16.5 percent).

# Health Risk Behaviors

---

## Weight

### Body Mass Index

Body Mass Index (BMI) is an index based on height and weight used to categorize an adult as not overweight or obese, overweight, or obese. BMI is calculated using the following formula:

$$\frac{\text{Weight in kilograms}}{(\text{Height in meters})^2}$$

The following table illustrates common category breakouts based on BMI that are used in this report.

BMI Category	BMI
Not Overweight or Obese*	< 25.0
Overweight**	25.0 +
Obese	30.0 +

\*The category "Not Overweight or Obese" also includes adults who are underweight based on BMI < 18.5). Although underweight is a recognized health risk, in 2008 only 1.4 percent of Idaho's adult population fell into the underweight category.

\*\* The category Overweight *includes* the category Obese.

# Health Risk Behaviors


## Weight

### Overweight and Obesity Trends

In 2008, a majority of Idaho adults were overweight or obese with 62.2 percent having a BMI of 25 or higher. This figure has remained steady in recent years but is up significantly since 1999.

The prevalence of obesity also increased significantly in the past ten years. One-in-four (25.2 percent) of Idaho adults were classified as obese in 2008.

**Percent of Idaho adults who were overweight and/or obese\*  
1999-2008**



\*Results for overweight and obese may differ from previous publications due to revisions in risk factor calculation.


**Note:** For each year, the percentage of "Overweight, Not Obese" plus the percentage of "Obese" may not sum to the exact figure for "Total Overweight" due to rounding.

# Health Risk Behaviors

## Weight


Overweight and obesity prevalence did not vary significantly by health district. District 6 and District 3 had significantly higher percentages of obese adults (32.9 percent and 31.0 percent, respectively) than the rest of the state.

**Percent of Idaho adults who were overweight and/or obese, 2008**  
Health District



Males were significantly more likely to be overweight than females (68.4 percent compared with 55.6 percent). The likelihood of being overweight increased with age until decreasing significantly among those 65 and over.

**Percent of Idaho adults who were overweight, 2008**  
Age and Sex




# Health Risk Behaviors

## Weight

Males (25.2 percent) and females (25.1 percent) were equally likely to be obese. Obesity prevalence increased with age until 65 years and over, where the rate dropped significantly (to 22.1 percent).

**Percent of Idaho adults who were obese, 2008  
Age and Sex**



### Other Highlights:


- The percentage of overweight adults was consistent across socio-economic categories such as income, education and employment status. However, differences were observed when looking at the percentage that were categorized as obese.
  - While the prevalence of overweight adults (BMI  $\geq$  25) did not differ significantly by education level, college graduates were significantly less likely than those with lower levels of education to be obese (20.1 percent compared with 27.4 percent).
  - Those with household incomes of \$75,000 or more were significantly less likely to be obese than those with incomes below \$35,000 (21.1 percent compared with 28.3 percent).
- Those who were overweight were significantly more likely than those who were not to have diabetes. The prevalence of diabetes was highest among those who were obese. The prevalence of diabetes among those who were not overweight was 2.9 percent. This compares with 5.4 percent of those who were overweight/not obese and 15.4 percent of those who were obese.

# Health Risk Behaviors

## Cigarette Smoking


Adults were classified as cigarette smokers if they had smoked at least 100 cigarettes in their lives *and* currently smoked every day or some days. After an increase in 2007, the prevalence of cigarette smoking among Idaho adults fell to 16.9 percent in 2008. Although the single year decrease was not statistically significant, the smoking rate has dropped significantly since 1999.

**Percent of Idaho adults who smoked cigarettes  
1999-2008**



Smoking rates did not vary significantly by health district. About one-in-five (20.5 percent) adults in District 1 were current smokers compared with just over one-in-ten (12.0 percent) in District 6.

**Percent of Idaho adults who smoked cigarettes, 2008  
Health District**




# Health Risk Behaviors

## Cigarette Smoking

In 2008, there was no significant difference between males and females overall in smoking prevalence (18.3 percent for males and 15.4 percent for females). Adults over the age of 55 were significantly less likely to smoke (12.3 percent) compared with those in younger age categories.

Percent of Idaho adults who smoked cigarettes, 2008  
Age and Sex



### Other Highlights:


- Smokers were significantly more likely to have “fair” or “poor” general health compared with non-smokers (20.8 percent compared with 12.6 percent).
- Smoking prevalence decreased as education level increased. Those with less than a high school education were five times as likely to report smoking as college graduates (34.0 percent compared with 6.7 percent, respectively).
- Adults with incomes below \$25,000 were significantly more likely to smoke cigarettes than those with higher incomes. More than one-in-four (27.8 percent) of those with incomes below \$25,000 smoked compared with fewer than one-in-ten (9.5 percent) of those with incomes above \$50,000.
- Unemployed Idahoans were significantly more likely to smoke than those in other employment categories (34.3 percent of the unemployed compared with 17.8 percent of the employed and 12.7 percent of the “other” employment category—students, homemakers, retirees and those unable to work).

# Health Risk Behaviors

## Alcohol Consumption

Binge drinking for females is defined as consuming four or more alcoholic beverages on one occasion in the last 30 days. For males, binge drinking is five or more alcoholic beverages on one occasion in the last 30 days. In 2008, 13.1 percent of Idaho adults reported binge drinking in the prior 30 days. This was not a significant change since 2005.


**Percent of Idaho adults who were binge drinkers  
2005-2008**



**Note:** The CDC adopted a revised definition of binge drinking in 2005. Binge drinking figures reported in previous BRFSS publications may not be comparable.

Binge drinking was significantly lower in District 6 than for the rest of the state. No other district differed significantly.

**Percent of Idaho adults who were binge drinkers, 2008  
Health District**




# Health Risk Behaviors

## Alcohol Consumption

Heavy drinking for females is defined as consuming more than 30 alcoholic beverages in the last 30 days. For males, heavy drinking is more than 60 alcoholic beverages in the last 30 days. The trend of heavy drinking in Idaho has remained flat over the past four years. In 2008, 4.6 percent of Idaho adults were heavy drinkers.


**Percent of Idaho adults who were heavy drinkers  
2005-2008**



**Note:** The CDC adopted a revised definition of heavy drinking in 2005. Heavy drinking figures reported in previous BRFSS publications may not be comparable.

Prevalence of heavy drinking varied from a low of 2.1 percent in District 7 to a high 6.8 percent in District 1.

**Percent of Idaho adults who were heavy drinkers, 2008  
Health District**




# Health Risk Behaviors

## Alcohol Consumption


The percentage of Idaho adults who engaged in binge drinking dropped significantly after the age of 55 (to 5.2 percent). In every age category, males were significantly more likely to binge drink than females.

**Percent of Idaho adults who were binge drinkers, 2008**  
Age and Sex



The percentage of males who engaged in heavy drinking was not significantly higher than that of females (5.3 percent compared with 4.0 percent). Those over the age of 65 were the least likely age category to be heavy drinkers. However, only middle aged adults (aged 45 to 64) were significantly more likely than those over 65 to be heavy drinkers.

**Percent of Idaho adults who were heavy drinkers, 2008**  
Age and Sex



# Health Risk Behaviors

---

## Alcohol Consumption

### Other Highlights:


- Binge and heavy drinking did not vary significantly by socioeconomic factors such as income or education. No significant differences were seen between the employed and unemployed.
- Binge drinkers were more than twice as likely to smoke cigarettes compared with those who were not binge drinkers (38.3 percent of binge drinkers smoked cigarettes compared with 13.6 percent of those who were not binge drinkers).
- Heavy drinkers were also significantly more likely to be smokers than those who were not heavy drinkers (38.6 percent of heavy drinkers smoked compared with 15.6 percent of those who were not heavy drinkers).
- Illicit drug use was reported significantly more often by binge drinkers and heavy drinkers compared with those who did not engage in those behaviors.
  - 14.4 percent of binge drinkers had used illicit drugs in the past year compared with 2.5 percent of those who were not binge drinkers.
  - 17.8 percent of heavy drinkers had used illicit drugs in the past year compared with 3.4 percent of those who were not heavy drinkers.

# Health Risk Behaviors

## Illicit Drug Use

In 2008, 4.1 percent of Idaho adults reported illicit drug use in the prior 12 months. This figure has not changed significantly since 2002. Drug use includes all drugs not prescribed by a doctor or used to get high or for curiosity. This includes marijuana, cocaine (including crack), heroin, methamphetamines and prescription painkillers.


**Percent of Idaho adults who used illicit drugs in the last 12 months, 2002-2008**  
Breaks in the trendline indicate data are not available for those years\*



\*Question wording changes make data prior to 2002 not comparable with 2002 and later.

Illicit drug use was highest in District 4. Both District 1 and District 4 showed significantly higher rates of drug use than District 7. There were no other significant differences by district.

**Percent of Idaho adults who used illicit drugs in the last 12 months, 2008**  
Health District




# Health Risk Behaviors

## Illicit Drug Use

Males and females did not differ significantly with regard to illicit drug use (5.4 percent and 2.8 percent, respectively). Adults aged 18 to 34 were significantly more likely to report drug use (7.8 percent) than older adults (2.1 percent of those 35 and older).

**Percent of Idaho adults who used illicit drugs  
in the last 12 months, 2008  
Age and Sex**



## Other Highlights:

- Those with a high school or less education reported using illicit drugs twice as often as those with some college or more (6.2 percent compared with 2.7 percent).
- Illicit drug use did not vary significantly by employment status or income level.
- More than one half (54.5 percent) of illicit drug users were current smokers.
- Illicit drug users reported binge drinking nearly four times as often as non-users (46.1 percent compared with 11.7 percent). Drug users reported heavy drinking five times as often as non-users (20.2 percent compared with 4.0 percent).


# Preventive Behaviors

## Seat Belt Use

The percentage of Idaho adults who did not consistently use a seat belt has decreased significantly from levels seen in 2003 and prior years. In 2008, just under one-quarter (23.4 percent) of adults did not always wear a seatbelt.


**Percent of Idaho adults who did not always wear a seat belt, 1999-2008**

Breaks in the trendline indicate data are not available for those years



Adults in District 4 were significantly less likely than those in the rest of the state to drive or ride without a seat belt. Districts 6 and 7 had the highest percentage of adults who did not consistently wear a seatbelt (38.5 percent and 31.3 percent, respectively).

**Percent of Idaho adults who did not always wear a seat belt, 2008**  
**Health District**




# Preventive Behaviors

## Seat Belt Use

Males were significantly more likely to not always wear their seat belt when driving or riding in a car compared with females (29.4 percent compared with 17.6 percent). Although there was no clear relationship between age and seat belt use in 2008, those over age 65 were the most likely to use a seat belt while young adults (aged 18 to 24) were the least likely.

**Percent of Idaho adults who did not always wear a seat belt, 2008  
Age and Sex**



### Other Highlights:

- Seat belt usage did not differ significantly by income level or between the employed and unemployed.
- Seat belt usage did increase with education level. Those with a high school diploma or less education were significantly more likely to report that they did not always use a seat belt compared with college graduates (28.6 percent compared with 16.7 percent).
- Seat belt usage was not significantly different for those with children in the household compared with those without children (24.5 percent and 22.5 percent, respectively).


# Preventive Behaviors

## Colorectal Cancer Screening

The American Cancer Society recommends colorectal cancer screening for adults of average risk aged 50 and older. The percentage of Idaho adults in this age group who had never received a sigmoidoscopy or colonoscopy has dropped steadily in the past ten years reaching a new low of 44.4 percent in 2008. This is a significant decrease since 1999.

**Percent of Idaho adults aged 50 and older who had never had a sigmoidoscopy or colonoscopy, 1999-2008**


Breaks in the trendline indicate data are not available for those years



District 6 and District 7 had the highest percentages of adults over 50 who had never received a sigmoidoscopy or colonoscopy (51.2 percent and 50.0 percent respectively). Compared with the state as a whole, those in District 4 were significantly less likely to have never had colorectal cancer screening.

**Percent of Idaho adults aged 50 and older who had never had a sigmoidoscopy or colonoscopy, 2008**


Health District



# Preventive Behaviors

## Colorectal Cancer Screening

Males and females did not differ significantly with regard to colorectal cancer screening. Around one-half of males (45.4 percent) and females (43.4 percent) had never had a sigmoidoscopy or colonoscopy. There were significant differences between age categories, however. Those aged 50 to 64 were significantly more likely to not have been screened than those aged 65 and older (52.8 percent compared with 31.4 percent).



### Other Highlights:


- College graduates were significantly less likely than those with less education to say they had not had colorectal cancer screening (37.3 percent of college graduates compared with 47.6 percent of those with some college or less).
- Those with incomes below \$25,000 were significantly more likely to have never had colorectal cancer screening compared with those with incomes of \$25,000 or higher (51.1 percent compared with 43.3 percent).
- Idaho adults aged 50 and over who did not have health insurance coverage reported never having a sigmoidoscopy or colonoscopy nearly twice as often as those who did have coverage (77.9 percent compared with 41.1 percent).

# Preventive Behaviors

## Breast Cancer Screening


The National Cancer Institute recommends that women aged 40 and above receive screening for breast cancer with mammography every one to two years. The percentage of Idaho women aged 40 and older not receiving this breast cancer screening has not changed significantly over the past decade. In 2008, 36.8 percent of women aged 40 and over had not received breast cancer screening in the prior two years.

**Percent of Idaho females aged 40 and older who had not had a mammogram and clinical breast exam in the last two years, 1999-2008**



Breast cancer screening varied widely by health district. Those in District 3 were the most likely to have not had the recommended screening while those in District 4 were the least likely.

**Percent of Idaho females aged 40 and older who had not had a mammogram and clinical breast exam in the last two years, 2008**




# Preventive Behaviors

---

## Breast Cancer Screening

Those aged 40 to 49 were significantly more likely than those 50 and over to have not had a mammogram and clinical breast exam in the prior two years.

**Percent of Idaho females aged 40 and older who had not had a mammogram and clinical breast exam in the last two years, 2008**  
**Age Groups**



### Other Highlights:


- About two-thirds (66.3 percent) of 40-plus women without health care coverage reported that they did not have breast cancer screening compared with about one-third (33.2 percent) of same aged women with health insurance.
- Those who had someone they considered a personal health care provider were significantly more likely to have had the recommended breast cancer screening compared with those who did not have a health care provider. Nearly two-in-three (62.4 percent) of 40-plus women who did not have a health care provider had not had breast cancer screening compared with 32.3 percent of those who had a provider.
- Higher incomes and higher education levels were associated with breast cancer screening.
  - o More than one-half (53.2 percent ) of those who had less than a high school education had not had the recommended breast cancer screening compared with 29.8 percent of college graduates.
  - o Those with incomes below \$25,000 were significantly more likely to have missed this screening compared with those with incomes above \$50,000 (51.0 percent compared with 32.5 percent).

# Preventive Behaviors

## Cervical Cancer Screening


The American Cancer Society recommends that women have a Pap test at least once every three years. The percentage of those with an intact cervix who had not received the test in the previous three years was 23.1 percent in 2008. While this was not a significant change since 2006, it is significantly higher than the decade low in 2002 (16.2 percent).

**Percent of Idaho females who had not had a Pap test in the last three years, 1999-2008**



Women residing in Health District 7 were significantly more likely than those in the rest of the state to have not had cervical cancer screening with about one-in-three (33.7 percent) reporting no Pap test in the previous three years. Those in District 1 were the least likely to have not had a Pap test in three years (16.2 percent).

**Percent of Idaho females who had not had a Pap test in the last three years, 2008**  
**Health District**




# Preventive Behaviors

## Cervical Cancer Screening

There were significant differences by age in the percentage receiving screening for cervical cancer through Pap tests. Women in the youngest (18 to 24 years) and oldest (65 years and over) age categories were significantly more likely to have not had cervical cancer screening compared with other age categories.

**Percent of Idaho females who had not had a Pap test within the last three years, 2008**  
**Age Groups**



### Other Highlights:

- Those with incomes below \$35,000 reported not having a Pap test twice as often as those with incomes of \$35,000 or higher (31.2 percent compared with 15.5 percent).
- College graduates were significantly less likely to have not received a Pap test in the previous three years (11.8 percent) when compared with those at all other education levels (less than high school: 31.4 percent; high school graduates: 26.9 percent; some college: 28.6 percent).
- Those who had someone they considered a personal health care doctor or provider were significantly more likely to have had a Pap test compared with those who did not. More than one-in-three (39.3 percent) of those who did not have a health care provider had not a Pap test in the previous three years compared with 17.1 percent of those who did have a provider.
- Women without healthcare coverage were significantly more likely to have not had a Pap test than those with coverage (35.4 percent compared with 20.1 percent).


# Preventive Behaviors

## Prostate Cancer Screening

While there is currently no standard recommendation for prostate cancer screening, the National Cancer Institute reports that age is the most common risk factor for prostate cancer. In 2008, about one-half (47.1 percent) of Idaho males aged 40 and older had not received a Prostate-Specific Antigen (PSA) test in the previous two years. This was not a significant change since 2001.


**Percent of Idaho males aged 40 and older who had not had a PSA test in the last two years, 2001-2008**

Breaks in the trendline indicate data are not available for those years



Prostate cancer screening through PSA tests among males 40 and over did not differ significantly by Health District.

**Percent of Idaho males aged 40 and older who had not had a PSA test in the last two years, 2008**  
Health District




# Preventive Behaviors

---

## Prostate Cancer Screening

Prostate cancer screening rates differed significantly by age. Around one-in-four (23.2 percent) men aged 70 and older had not received a PSA exam in the prior two years. This compares with more than three-in-four (74.5 percent) among men aged 40 to 49 who had not at a PSA test.

**Percent of Idaho males aged 40 and older who had not had a PSA test in the last two years, 2008**  
Age Groups



### Other Highlights:

- Men aged 40 and over with household incomes below \$15,000 were significantly more likely to have not received a PSA test in the prior two years (64.1 percent) compared with those with incomes of \$35,000 or more (45.9 percent).
- Nearly eight-in-ten (77.8 percent) of 40-plus men with no health insurance had not had a PSA test within the past two years. This compares with less than half (43.3 percent) of 40-plus men with health insurance.
- Those who did not have a personal doctor or health care provider were less likely to have had a PSA test. About three-quarters (75.7 percent) of those without a provider had not had a PSA test in the past two years. Among those who did have a personal provider, 38.9 percent had not had this screening.


# Preventive Behaviors

## Dental Visits

The percentage of Idaho adults who had not visited a dentist in the prior year remained consistent over the past ten years. In 2008, about one-in-three (32.0 percent) adults had not visited the dentist in the prior 12 months.


**Percent of Idaho adults who had not visited the dentist in the last 12 months, 1999-2008**

Breaks in the trendline indicate data are not available for those years



Adults residing in Health Districts 7 and 4 were significantly less likely than those in the rest of the state to have not had a dental visit in the prior 12 months (28.8 percent and 26.8 percent, respectively).

**Percent of Idaho adults who had not visited the dentist in the last 12 months, 2008**  
Health District




# Preventive Behaviors

## Dental Visits

There were no statistically significant differences by age or sex relating to dental visits in 2008.

**Percent of Idaho adults who had not visited the dentist  
in the last 12 months, 2008  
by Age and Sex**



### Other Highlights:

- The percentage of adults who had not visited a dentist in the previous year decreased as income level increased. More than one-half (50.6 percent) of those with incomes below \$25,000 had not had a dental visit compared with 18.0 percent of those with incomes above \$50,000.
- Those with less than a high school education reported not having a dental visit in the past year more than three times as often as college graduates (54.8 percent compared with 17.9 percent).
- Those who did not have dental insurance were significantly more likely to report not visiting a dentist in the past year compared with those who did have dental insurance (47.0 percent compared with 20.7 percent).

# Understanding the Data Tables

---

The data tables on the following pages provide both state and health district data for various demographic subcategories. An understanding of the tables is important to interpreting the data correctly.

- A. This title refers to the risk factor.
- B. This title refers to the population surveyed. For most questions, this will be the adult (aged 18 and over) Idaho population. However, for some calculated variables and some risk factors, the population surveyed may be a subset of adults aged 18 and over. For instance, the Breast Cancer Screening risk factor table will include only the female population aged 40 and over, because only females were asked those questions and breast cancer screening is recommended for that age group. Colorectal Cancer Screening will include only adults over the age of 50, because only they were asked questions about colorectal cancer screening. It is important to understand the population the data represent in order to accurately interpret the data in the tables.
- C. The labels across the top of the table (i.e., Statewide and District) refer to the geographic region of residence of the population in (B). For more information on geographic break-outs, please see page iv of this report for a map of Idaho's seven public health districts.
- D. The labels on the left side of the table refer to the particular group the numbers represent. For example, the row labeled "Female" will contain data for women for the geographic region indicated at the top of the table (C). In the following example table, the circled data point should be interpreted as:

“In 2008, 15.3 percent of females in Health District 1 were at risk for this risk factor.”

- E. The symbol "n" refers to the number of people in the sample within the indicated demographic group and geographic region who responded to the question. It does NOT represent the number of people who have the risk factor in the population. An asterisk indicates that fewer than 50 people in the group responded (see Methodology and Reporting section, pages 1-2 for explanation on sampling and suppressing data for sample sizes less than 50). In the following example table, the boxed data point should be interpreted as:

“In 2008, there were 728 people in Health District 1 sampled for this risk factor.”

# Understanding the Data Tables

---

F. The shaded columns contain the prevalence (in percent) of the risk factor. This percentage is a weighted percentage and is not calculated using the sample frequencies from the column containing n. Therefore, it is not possible to calculate the prevalence by simply dividing n by the total number of people sampled for the year (see Methodology section, pages 1-2 for explanation on weighting procedures).

G. The two numbers in the column labeled "95% CI" are the lower and upper limits of the confidence interval. Confidence intervals are a way to measure sampling error and define the range of values where the "true" percentage would be found. This interval can be interpreted to mean there is a 95% certainty the true prevalence of the risk factor in the population falls within the confidence interval. In the following example table, the confidence interval data should be interpreted as:

"For this risk factor, the prevalence in Idaho was 14.0 percent in 2008. We can say, with 95 percent certainty, the actual population prevalence falls within the range of 12.9 percent to 15.2 percent."

<b>A Risk Factor</b>																
<b>B Population</b>																
<b>D</b>	<b>C</b>			<b>E</b>	Public Health District											
	Statewide				District 1			District 2			District 3					
	%	95% CI			n	%	95% CI		n	%	95% CI		n	%	95% CI	
<b>TOTAL</b>	<b>F</b> 14.0	<b>G</b> 12.9	15.2	5,099	16.5	13.5	19.9	728	16.5	13.7	19.8	751	16.5	13.5	20.1	749
<b>SEX</b>																
Male	13.8	12.1	15.6	1,986	17.6	12.9	23.6	282	13.8	10.0	18.7	319	15.8	11.9	20.8	267
Female	14.2	12.8	15.8	3,113	15.3	2.1	19.3	446	19.3	15.5	23.9	432	17.2	12.9	22.7	482

# General Health Status

Idaho adults who reported their general health status as "fair" or "poor," 2008

	Statewide			Public Health District												
				District 1			District 2			District 3						
	%	95% CI		n	%	95% CI		n	%	95% CI		n	%	95% CI		n
<b>TOTAL</b>	<b>14.0</b>	12.9	15.2	5,099	<b>16.5</b>	13.5	19.9	728	<b>16.5</b>	13.7	19.8	751	<b>16.5</b>	13.5	20.1	749
<b>SEX</b>																
Male	<b>13.8</b>	12.1	15.6	1,986	<b>17.6</b>	12.9	23.6	282	<b>13.8</b>	10.0	18.7	319	<b>15.8</b>	11.9	20.8	267
Female	<b>14.2</b>	12.8	15.8	3,113	<b>15.3</b>	12.1	19.3	446	<b>19.3</b>	15.5	23.9	432	<b>17.2</b>	12.9	22.7	482
<b>AGE</b>																
18-24	<b>6.2</b>	3.2	11.8	164	*	*	*	*	*	*	*	*	*	*	*	*
25-34	<b>7.2</b>	5.3	9.9	629	<b>16.5</b>	8.4	30.0	67	<b>12.1</b>	6.8	20.8	85	<b>8.4</b>	4.4	15.2	108
35-44	<b>10.6</b>	8.4	13.5	771	<b>12.0</b>	6.5	21.1	104	<b>18.9</b>	12.0	28.3	110	<b>10.6</b>	5.7	19.0	103
45-54	<b>14.5</b>	12.1	17.4	1,038	<b>16.8</b>	11.6	23.7	169	<b>14.8</b>	9.6	22.0	154	<b>20.6</b>	14.0	29.2	131
55-64	<b>21.5</b>	18.7	24.5	1,063	<b>20.3</b>	14.6	27.6	160	<b>20.3</b>	14.3	28.0	164	<b>28.5</b>	20.8	37.6	151
65+	<b>25.5</b>	22.9	28.3	1,392	<b>23.3</b>	17.3	30.6	199	<b>25.1</b>	19.0	32.5	214	<b>26.3</b>	20.0	33.7	224
18-34	<b>6.9</b>	5.1	9.3	793	<b>11.8</b>	6.1	21.9	89	<b>10.0</b>	5.7	17.0	106	<b>9.1</b>	4.4	17.9	130
35-54	<b>12.6</b>	10.9	14.6	1,809	<b>14.6</b>	10.5	19.8	273	<b>16.6</b>	12.2	22.2	264	<b>15.4</b>	11.0	21.1	234
55+	<b>23.6</b>	21.7	25.6	2,455	<b>21.9</b>	17.6	26.9	359	<b>23.1</b>	18.6	28.3	378	<b>27.2</b>	22.2	32.9	375
<b>SEX and AGE</b>																
Male																
18-34	<b>6.3</b>	3.9	10.1	290	*	*	*	*	*	*	*	*	<b>6.6</b>	2.8	14.7	52
35-54	<b>11.9</b>	9.4	14.9	728	<b>14.7</b>	8.7	23.7	104	<b>14.0</b>	8.2	22.9	127	<b>13.1</b>	7.5	21.8	88
55+	<b>25.1</b>	22.1	28.4	958	<b>23.0</b>	16.6	30.9	147	<b>19.6</b>	13.2	28.1	147	<b>31.6</b>	23.1	41.5	123
Female																
18-34	<b>7.5</b>	5.1	10.8	503	<b>8.2</b>	3.3	18.8	59	<b>11.9</b>	5.8	22.8	61	<b>11.7</b>	4.1	28.8	78
35-54	<b>13.4</b>	11.2	16.0	1,081	<b>14.5</b>	9.8	20.9	169	<b>19.3</b>	13.3	27.2	137	<b>17.6</b>	11.6	25.8	146
55+	<b>22.2</b>	19.9	24.7	1,497	<b>20.9</b>	15.6	27.6	212	<b>26.2</b>	20.4	33.0	231	<b>23.4</b>	18.1	29.7	252
<b>INCOME</b>																
Less than \$15,000	<b>35.7</b>	29.9	41.9	478	<b>38.7</b>	26.6	52.3	76	<b>35.0</b>	23.6	48.4	84	<b>37.7</b>	25.0	52.2	79
\$15,000 - \$24,999	<b>22.9</b>	19.3	27.0	826	<b>27.5</b>	18.4	38.8	119	<b>22.8</b>	15.3	32.5	134	<b>26.8</b>	18.9	36.5	149
\$25,000 - \$34,999	<b>14.9</b>	11.7	18.7	662	<b>17.1</b>	9.4	29.2	78	<b>21.6</b>	14.0	31.9	105	<b>18.0</b>	9.0	32.8	109
\$35,000 - \$49,999	<b>9.8</b>	7.8	12.1	853	<b>12.4</b>	7.5	19.9	116	<b>7.4</b>	4.2	12.8	128	<b>11.1</b>	6.4	18.6	130
\$50,000-\$74,999	<b>7.4</b>	5.4	10.0	793	<b>4.5</b>	2.0	9.7	114	<b>7.8</b>	3.6	16.0	110	<b>11.0</b>	5.4	20.8	106
\$75,000+	<b>5.9</b>	4.2	8.2	911	<b>10.0</b>	4.7	19.9	135	<b>8.6</b>	4.0	17.4	111	<b>5.0</b>	1.9	12.8	86
<b>EMPLOYMENT</b>																
Employed	<b>8.0</b>	6.9	9.4	2,721	<b>9.3</b>	6.3	13.4	378	<b>9.6</b>	6.7	13.5	387	<b>9.1</b>	6.4	12.6	382
Unemployed	<b>19.6</b>	13.2	28.2	174	*	*	*	*	*	*	*	*	*	*	*	*
Other**	<b>24.0</b>	21.8	26.3	2,196	<b>27.9</b>	22.5	34.1	320	<b>25.8</b>	20.6	31.8	333	<b>29.7</b>	23.1	37.3	336
<b>EDUCATION</b>																
K-11th Grade	<b>26.7</b>	21.6	32.4	407	<b>26.6</b>	13.9	44.7	54	<b>31.5</b>	18.5	48.1	54	<b>22.7</b>	14.8	33.2	93
12th Grade or GED	<b>16.8</b>	14.5	19.3	1,515	<b>18.7</b>	13.7	25.0	231	<b>17.3</b>	12.9	23.0	251	<b>21.5</b>	15.0	30.0	255
Some College	<b>13.7</b>	11.8	15.8	1,621	<b>17.3</b>	12.4	23.6	245	<b>20.6</b>	15.0	27.5	226	<b>13.8</b>	9.5	19.6	228
College Graduate+	<b>7.5</b>	6.1	9.3	1,546	<b>9.2</b>	5.4	15.1	195	<b>6.3</b>	3.8	10.1	219	<b>8.3</b>	4.8	14.2	171

\*Figure not reliable by BRFSS standards (n<50)

\*\*Other includes students, homemakers, retirees, and persons unable to work

# General Health Status

Idaho adults who reported their general health status as "fair" or "poor," 2008

	Public Health District															
	District 4				District 5				District 6				District 7			
	%	95% CI		n	%	95% CI		n	%	95% CI		n	%	95% CI		n
<b>TOTAL</b>	<b>11.5</b>	9.3	14.3	751	<b>13.1</b>	10.7	15.9	724	<b>14.2</b>	11.5	17.3	704	<b>12.6</b>	10.0	15.8	692
<b>SEX</b>																
Male	<b>12.6</b>	9.0	17.2	294	<b>11.2</b>	8.1	15.3	282	<b>14.3</b>	10.4	19.4	256	<b>11.2</b>	7.6	16.2	286
Female	<b>10.5</b>	7.9	13.8	457	<b>14.9</b>	11.6	19.0	442	<b>14.0</b>	10.7	18.3	448	<b>14.0</b>	10.6	18.3	406
<b>AGE</b>																
18-24	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
25-34	<b>2.1</b>	0.6	6.5	87	<b>6.8</b>	3.3	13.6	78	<b>3.3</b>	1.1	9.5	93	<b>9.3</b>	3.6	22.1	111
35-44	<b>10.3</b>	6.0	17.1	149	<b>4.8</b>	2.0	11.0	106	<b>13.2</b>	7.1	23.3	88	<b>9.0</b>	5.0	15.5	111
45-54	<b>14.2</b>	8.9	21.8	150	<b>8.1</b>	4.6	13.8	153	<b>17.0</b>	10.6	26.2	141	<b>9.3</b>	5.5	15.3	140
55-64	<b>17.7</b>	12.2	25.1	164	<b>21.6</b>	15.1	30.0	140	<b>23.9</b>	17.4	31.8	163	<b>21.8</b>	14.8	30.9	121
65+	<b>21.9</b>	16.0	29.3	171	<b>27.8</b>	21.8	34.6	218	<b>27.3</b>	20.8	35.0	190	<b>31.1</b>	24.2	39.0	176
18-34	<b>4.3</b>	1.8	10.1	110	<b>8.0</b>	4.2	14.5	100	<b>2.9</b>	1.0	7.8	117	<b>5.8</b>	2.4	13.5	141
35-54	<b>12.2</b>	8.6	17.0	299	<b>6.6</b>	4.1	10.3	259	<b>15.2</b>	10.5	21.7	229	<b>9.1</b>	6.2	13.3	251
55+	<b>19.8</b>	15.5	24.8	335	<b>25.1</b>	20.5	30.3	358	<b>25.7</b>	20.9	31.1	353	<b>26.6</b>	21.4	32.4	297
<b>SEX and AGE</b>																
Male																
18-34	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
35-54	<b>12.8</b>	7.7	20.7	118	<b>5.6</b>	2.6	11.6	95	<b>15.7</b>	8.8	26.4	88	<b>5.9</b>	2.9	11.6	108
55+	<b>21.8</b>	15.3	30.1	137	<b>26.7</b>	19.8	35.0	149	<b>28.3</b>	20.6	37.4	127	<b>26.8</b>	19.3	35.9	128
Female																
18-34	<b>2.8</b>	0.9	8.8	72	<b>14.0</b>	7.5	24.8	64	<b>4.2</b>	1.3	13.1	77	<b>6.4</b>	2.4	16.0	92
35-54	<b>11.5</b>	7.2	17.8	181	<b>7.6</b>	4.2	13.3	164	<b>14.8</b>	9.0	23.2	141	<b>12.4</b>	7.8	19.1	143
55+	<b>18.0</b>	12.9	24.6	198	<b>23.6</b>	18.0	30.4	209	<b>23.4</b>	17.9	29.9	226	<b>26.3</b>	19.8	34.2	169
<b>INCOME</b>																
Less than \$15,000	<b>40.9</b>	25.8	58.0	55	<b>39.3</b>	26.3	54.1	66	<b>31.6</b>	18.2	48.9	68	<b>25.4</b>	13.0	43.6	50
\$15,000 - \$24,999	<b>17.4</b>	10.2	28.2	81	<b>27.0</b>	18.4	37.7	121	<b>19.6</b>	11.6	31.1	114	<b>20.4</b>	11.7	33.2	108
\$25,000 - \$34,999	<b>12.4</b>	7.1	20.6	88	<b>13.9</b>	7.8	23.5	101	<b>15.0</b>	8.7	24.8	101	<b>9.7</b>	5.2	17.3	80
\$35,000 - \$49,999	<b>8.4</b>	4.8	14.4	133	<b>5.9</b>	3.0	11.3	113	<b>12.1</b>	7.1	19.9	121	<b>10.6</b>	5.8	18.6	112
\$50,000-\$74,999	<b>6.8</b>	2.8	15.4	116	<b>5.0</b>	2.2	10.9	109	<b>9.2</b>	5.0	16.3	111	<b>8.7</b>	5.1	14.5	127
\$75,000+	<b>6.0</b>	3.2	10.8	201	<b>3.4</b>	1.3	9.0	119	<b>5.2</b>	2.3	11.4	118	<b>3.0</b>	1.1	7.5	141
<b>EMPLOYMENT</b>																
Employed	<b>7.3</b>	5.0	10.6	417	<b>6.8</b>	4.6	10.0	409	<b>8.0</b>	5.3	12.0	358	<b>7.4</b>	4.7	11.4	390
Unemployed	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Other**	<b>19.3</b>	15.0	24.6	300	<b>25.2</b>	20.1	31.0	297	<b>23.3</b>	18.1	29.5	328	<b>19.7</b>	15.0	25.4	282
<b>EDUCATION</b>																
K-11th Grade	*	*	*	*	<b>20.4</b>	12.5	31.6	71	<b>26.6</b>	14.1	44.3	51	*	*	*	*
12th Grade or GED	<b>11.8</b>	7.7	17.9	173	<b>14.2</b>	9.7	20.3	207	<b>17.3</b>	12.5	23.4	219	<b>19.5</b>	13.5	27.3	179
Some College	<b>12.8</b>	8.7	18.4	220	<b>14.6</b>	10.5	19.8	246	<b>12.9</b>	8.8	18.5	230	<b>7.6</b>	4.9	11.6	226
College Graduate+	<b>7.0</b>	4.5	10.8	317	<b>6.3</b>	3.7	10.6	199	<b>7.5</b>	4.4	12.5	202	<b>8.5</b>	5.6	12.7	243

\*Figure not reliable by BRFSS standards (n<50)

\*\*Other includes students, homemakers, retirees, and persons unable to work

# Health Care Coverage

## Idaho adults without health care coverage, 2008

	Statewide			Public Health District												
				District 1			District 2			District 3						
	%	95% CI		n	%	95% CI		n	%	95% CI		n	%	95% CI		n
<b>TOTAL</b>	<b>18.5</b>	16.9	20.4	5,099	<b>20.5</b>	16.7	25.0	727	<b>16.2</b>	12.8	20.2	750	<b>25.4</b>	20.9	30.5	750
<b>SEX</b>																
Male	<b>21.0</b>	18.2	24.1	1,986	<b>22.6</b>	16.4	30.3	283	<b>16.0</b>	11.1	22.6	318	<b>31.1</b>	24.0	39.3	268
Female	<b>16.1</b>	14.4	18.1	3,113	<b>18.5</b>	14.4	23.5	444	<b>16.3</b>	12.2	21.5	432	<b>19.8</b>	14.8	25.9	482
<b>AGE</b>																
18-24	<b>37.5</b>	29.0	46.8	160	*	*	*	*	*	*	*	*	*	*	*	*
25-34	<b>28.8</b>	24.4	33.6	628	<b>30.4</b>	19.7	43.9	67	<b>23.1</b>	14.5	34.6	85	<b>36.8</b>	27.4	47.4	108
35-44	<b>15.8</b>	13.0	19.0	772	<b>21.6</b>	14.2	31.4	104	<b>21.5</b>	13.6	32.2	111	<b>24.0</b>	16.2	34.0	103
45-54	<b>15.9</b>	13.4	18.8	1,038	<b>19.3</b>	13.6	26.5	169	<b>17.4</b>	11.6	25.2	154	<b>18.9</b>	12.7	27.3	131
55-64	<b>13.4</b>	11.2	16.0	1,067	<b>17.2</b>	11.6	24.7	161	<b>12.6</b>	7.9	19.3	165	<b>15.7</b>	9.8	24.2	152
65+	<b>1.8</b>	1.1	2.9	1,393	<b>0.4</b>	0.1	1.7	199	<b>3.9</b>	1.1	12.3	213	<b>2.7</b>	1.1	6.6	224
18-34	<b>31.9</b>	27.7	36.4	788	<b>36.4</b>	25.7	48.8	88	<b>21.7</b>	14.2	31.6	104	<b>43.6</b>	33.2	54.6	130
35-54	<b>15.8</b>	13.9	17.9	1,810	<b>20.3</b>	15.6	26.1	273	<b>19.2</b>	14.2	25.5	265	<b>21.6</b>	16.3	28.0	234
55+	<b>7.3</b>	6.1	8.6	2,460	<b>8.4</b>	5.6	12.3	360	<b>7.7</b>	4.8	12.1	378	<b>8.5</b>	5.6	12.8	376
<b>SEX and AGE</b>																
Male																
18-34	<b>37.7</b>	30.9	44.9	288	*	*	*	*	*	*	*	*	<b>55.6</b>	39.8	70.3	52
35-54	<b>16.0</b>	13.2	19.4	728	<b>21.6</b>	14.4	31.0	104	<b>17.5</b>	10.8	26.9	127	<b>23.5</b>	15.5	34.1	88
55+	<b>7.6</b>	5.9	9.9	960	<b>6.8</b>	3.7	12.1	148	<b>7.8</b>	3.4	16.5	147	<b>9.7</b>	4.9	18.4	124
Female																
18-34	<b>25.7</b>	21.2	30.8	500	<b>29.5</b>	18.5	43.5	58	<b>20.9</b>	11.8	34.3	60	<b>31.0</b>	19.1	46.1	78
35-54	<b>15.6</b>	13.3	18.3	1,082	<b>19.2</b>	13.5	26.5	169	<b>21.1</b>	14.2	30.1	138	<b>19.7</b>	13.6	27.7	146
55+	<b>6.9</b>	5.5	8.7	1,500	<b>9.8</b>	5.8	16.2	212	<b>7.6</b>	4.5	12.5	231	<b>7.4</b>	4.7	11.6	252
<b>INCOME</b>																
Less than \$15,000	<b>33.8</b>	27.3	40.9	482	<b>25.7</b>	15.9	38.7	76	<b>21.5</b>	12.0	35.5	85	<b>47.3</b>	32.5	62.6	80
\$15,000 - \$24,999	<b>37.5</b>	32.1	43.2	828	<b>39.1</b>	27.9	51.5	120	<b>38.7</b>	28.0	50.6	133	<b>43.7</b>	32.8	55.2	149
\$25,000 - \$34,999	<b>25.3</b>	20.0	31.4	661	<b>34.4</b>	21.4	50.3	78	<b>12.8</b>	6.3	24.0	104	<b>28.1</b>	17.4	42.1	109
\$35,000 - \$49,999	<b>16.0</b>	12.8	19.8	852	<b>22.6</b>	13.7	35.1	115	<b>14.2</b>	7.5	25.3	128	<b>17.0</b>	10.0	27.6	130
\$50,000-\$74,999	<b>6.3</b>	4.3	9.1	793	<b>9.6</b>	4.8	18.2	114	<b>5.8</b>	2.3	13.7	110	<b>4.4</b>	1.4	12.6	106
\$75,000+	<b>4.6</b>	2.8	7.5	911	<b>2.1</b>	0.7	5.8	135	<b>0.8</b>	0.1	5.7	111	<b>10.2</b>	3.2	27.8	86
<b>EMPLOYMENT</b>																
Employed	<b>19.2</b>	17.0	21.6	2,717	<b>21.5</b>	16.3	27.8	377	<b>16.7</b>	12.4	22.2	386	<b>26.7</b>	21.2	33.0	382
Unemployed	<b>56.9</b>	46.3	66.8	176	*	*	*	*	*	*	*	*	*	*	*	*
Other**	<b>12.1</b>	10.0	14.6	2,198	<b>12.9</b>	8.3	19.5	320	<b>10.6</b>	6.8	16.2	333	<b>13.6</b>	8.1	22.0	336
<b>EDUCATION</b>																
K-11th Grade	<b>46.1</b>	39.1	53.3	406	<b>50.2</b>	32.2	68.2	54	<b>29.6</b>	16.5	47.2	54	<b>54.8</b>	40.8	67.9	93
12th Grade or GED	<b>24.0</b>	20.3	28.0	1,513	<b>24.8</b>	17.9	33.3	231	<b>21.5</b>	14.9	30.0	250	<b>25.6</b>	17.9	35.2	256
Some College	<b>15.9</b>	13.6	18.6	1,622	<b>18.9</b>	13.8	25.4	244	<b>14.6</b>	9.7	21.4	225	<b>21.7</b>	14.4	31.4	228
College Graduate+	<b>6.9</b>	5.2	9.1	1,547	<b>6.9</b>	3.9	11.9	195	<b>5.2</b>	2.9	9.3	220	<b>7.7</b>	3.9	14.6	171

\*Figure not reliable by BRFSS standards (n<50)

\*\*Other includes students, homemakers, retirees, and persons unable to work

# Health Care Coverage

## Idaho adults without health care coverage, 2008

	Public Health District															
	District 4				District 5				District 6				District 7			
	%	95% CI		n	%	95% CI		n	%	95% CI		n	%	95% CI		n
<b>TOTAL</b>	<b>16.5</b>	12.8	21.1	750	<b>21.2</b>	17.2	26.0	727	<b>14.2</b>	10.6	18.6	703	<b>14.6</b>	11.0	19.0	692
<b>SEX</b>																
Male	<b>19.5</b>	13.5	27.4	294	<b>21.6</b>	15.1	29.9	283	<b>15.8</b>	10.3	23.6	256	<b>16.0</b>	10.3	24.2	284
Female	<b>13.4</b>	9.8	18.1	456	<b>20.9</b>	16.4	26.2	444	<b>12.5</b>	8.7	17.8	447	<b>13.2</b>	9.6	17.9	408
<b>AGE</b>																
18-24	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
25-34	<b>23.9</b>	15.1	35.8	87	<b>43.6</b>	31.5	56.5	78	<b>18.3</b>	10.1	31.1	93	<b>25.9</b>	16.8	37.7	110
35-44	<b>10.4</b>	5.9	17.5	149	<b>14.0</b>	8.3	22.7	107	<b>10.3</b>	5.4	18.9	88	<b>15.6</b>	9.7	23.9	110
45-54	<b>14.0</b>	8.8	21.5	150	<b>15.9</b>	10.3	23.7	153	<b>17.9</b>	11.5	26.8	141	<b>9.6</b>	5.5	16.1	140
55-64	<b>10.7</b>	6.6	16.7	164	<b>14.3</b>	9.3	21.4	141	<b>12.1</b>	7.3	19.4	163	<b>11.9</b>	6.8	20.2	121
65+	<b>1.6</b>	0.4	5.8	171	<b>3.1</b>	1.3	7.2	219	<b>0.7</b>	0.2	2.8	189	<b>0.7</b>	0.1	5.1	178
18-34	<b>30.2</b>	20.8	41.5	109	<b>43.4</b>	32.7	54.8	100	<b>20.8</b>	12.7	32.2	117	<b>22.8</b>	14.7	33.5	140
35-54	<b>12.1</b>	8.5	17.0	299	<b>15.0</b>	10.7	20.5	260	<b>14.3</b>	9.9	20.3	229	<b>12.4</b>	8.7	17.4	250
55+	<b>6.2</b>	4.0	9.7	335	<b>8.0</b>	5.5	11.6	360	<b>6.1</b>	3.7	9.8	352	<b>6.2</b>	3.6	10.5	299
<b>SEX and AGE</b>																
Male																
18-34	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
35-54	<b>13.6</b>	8.3	21.6	118	<b>12.4</b>	7.0	21.2	95	<b>11.9</b>	6.1	21.9	88	<b>12.2</b>	6.8	20.7	108
55+	<b>5.2</b>	2.3	11.4	137	<b>10.3</b>	6.1	16.9	150	<b>7.4</b>	3.8	14.1	127	<b>8.7</b>	4.3	17.0	127
Female																
18-34	<b>22.5</b>	13.5	35.0	71	<b>43.4</b>	31.4	56.3	64	<b>15.0</b>	7.0	29.0	77	<b>20.7</b>	12.6	32.0	92
35-54	<b>10.5</b>	6.3	16.7	181	<b>17.6</b>	11.9	25.3	165	<b>16.7</b>	11.0	24.7	141	<b>12.6</b>	8.2	19.0	142
55+	<b>7.2</b>	4.3	11.8	198	<b>6.0</b>	3.5	10.2	210	<b>4.9</b>	2.4	9.7	225	<b>3.9</b>	1.7	8.6	172
<b>INCOME</b>																
Less than \$15,000	<b>49.5</b>	32.2	66.9	55	<b>46.4</b>	31.5	62.0	67	<b>27.4</b>	12.5	49.9	68	<b>10.0</b>	4.2	21.7	51
\$15,000 - \$24,999	<b>40.7</b>	25.9	57.4	81	<b>34.4</b>	21.1	50.7	123	<b>27.8</b>	17.8	40.6	114	<b>29.5</b>	18.3	43.9	108
\$25,000 - \$34,999	<b>20.2</b>	9.5	37.8	88	<b>29.5</b>	19.2	42.5	101	<b>22.4</b>	11.6	38.9	101	<b>30.1</b>	15.3	50.7	80
\$35,000 - \$49,999	<b>10.9</b>	5.6	19.9	133	<b>21.9</b>	13.5	33.6	113	<b>8.3</b>	3.8	17.2	121	<b>19.8</b>	11.1	32.6	112
\$50,000-\$74,999	<b>5.5</b>	1.9	15.2	116	<b>8.3</b>	3.2	19.9	109	<b>6.1</b>	2.6	13.5	111	<b>4.8</b>	2.0	10.8	127
\$75,000+	<b>5.1</b>	2.1	11.9	201	<b>5.6</b>	2.5	12.0	119	<b>5.5</b>	1.7	16.3	118	<b>1.1</b>	0.2	7.5	141
<b>EMPLOYMENT</b>																
Employed	<b>14.9</b>	10.2	21.2	416	<b>22.9</b>	17.6	29.4	410	<b>17.9</b>	12.7	24.5	358	<b>16.3</b>	11.8	22.2	388
Unemployed	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Other**	<b>13.1</b>	8.6	19.5	300	<b>16.2</b>	10.8	23.8	298	<b>7.5</b>	4.3	12.8	327	<b>9.2</b>	4.6	17.7	284
<b>EDUCATION</b>																
K-11th Grade	*	*	*	*	<b>48.6</b>	32.9	64.6	71	<b>30.1</b>	15.3	50.7	51	*	*	*	*
12th Grade or GED	<b>29.0</b>	19.1	41.4	172	<b>23.5</b>	16.3	32.5	208	<b>15.9</b>	9.0	26.5	218	<b>19.2</b>	12.8	27.7	178
Some College	<b>13.1</b>	8.6	19.4	220	<b>20.5</b>	14.5	28.3	247	<b>10.8</b>	6.6	17.4	230	<b>11.7</b>	7.3	18.3	228
College Graduate+	<b>7.2</b>	3.9	12.8	317	<b>6.3</b>	3.2	12.2	199	<b>9.9</b>	5.5	17.3	202	<b>4.3</b>	2.2	8.3	243

\*Figure not reliable by BRFSS standards (n<50)

\*\*Other includes students, homemakers, retirees, and persons unable to work

# Dental Insurance

Idaho adults without dental insurance, 2008

	Statewide			Public Health District												
				District 1			District 2			District 3						
	%	95% CI		n	%	95% CI		n	%	95% CI		n	%	95% CI		n
<b>TOTAL</b>	<b>43.0</b>	41.0	44.9	4,814	<b>48.6</b>	43.9	53.3	685	<b>46.1</b>	41.5	50.7	708	<b>48.5</b>	43.5	53.5	702
<b>SEX</b>																
Male	<b>43.9</b>	40.7	47.1	1,871	<b>49.3</b>	41.9	56.8	271	<b>48.0</b>	40.9	55.2	300	<b>51.6</b>	43.9	59.3	248
Female	<b>42.1</b>	39.7	44.4	2,943	<b>47.9</b>	42.3	53.5	414	<b>44.1</b>	38.4	50.0	408	<b>45.4</b>	39.2	51.8	454
<b>AGE</b>																
18-24	<b>48.2</b>	38.9	57.6	151	*	*	*	*	*	*	*	*	*	*	*	*
25-34	<b>39.9</b>	35.0	45.0	600	<b>44.9</b>	31.8	58.7	62	<b>45.3</b>	34.1	57.1	80	<b>50.0</b>	39.6	60.3	103
35-44	<b>31.2</b>	27.4	35.3	724	<b>38.4</b>	28.2	49.8	98	<b>34.8</b>	25.3	45.6	103	<b>38.9</b>	28.7	50.2	94
45-54	<b>34.7</b>	31.3	38.4	990	<b>43.1</b>	35.0	51.5	161	<b>36.8</b>	28.8	45.6	146	<b>30.6</b>	22.6	39.9	125
55-64	<b>44.9</b>	41.4	48.6	1,022	<b>46.2</b>	37.9	54.8	149	<b>45.9</b>	37.5	54.4	159	<b>50.3</b>	41.0	59.5	147
65+	<b>64.8</b>	61.7	67.9	1,294	<b>64.8</b>	56.7	72.2	188	<b>70.2</b>	62.8	76.6	202	<b>65.1</b>	57.3	72.1	205
18-34	<b>42.8</b>	38.2	47.5	751	<b>49.2</b>	37.2	61.3	83	<b>41.9</b>	31.7	52.8	97	<b>54.1</b>	43.0	64.7	124
35-54	<b>33.0</b>	30.4	35.7	1,714	<b>40.9</b>	34.4	47.8	259	<b>35.9</b>	29.6	42.7	249	<b>34.8</b>	28.2	42.1	219
55+	<b>55.3</b>	52.9	57.7	2,316	<b>56.1</b>	50.3	61.8	337	<b>59.5</b>	53.8	65.0	361	<b>58.3</b>	52.2	64.1	352
<b>SEX and AGE</b>																
Male																
18-34	<b>47.5</b>	40.3	54.7	276	*	*	*	*	*	*	*	*	*	*	*	*
35-54	<b>33.1</b>	29.1	37.4	679	<b>47.8</b>	37.2	58.5	98	<b>36.5</b>	27.5	46.7	116	<b>37.0</b>	26.6	48.9	82
55+	<b>53.3</b>	49.5	57.0	907	<b>50.5</b>	41.9	59.2	142	<b>60.7</b>	51.6	69.0	143	<b>56.1</b>	45.9	65.8	114
Female																
18-34	<b>37.8</b>	32.5	43.5	475	<b>48.8</b>	34.7	63.2	53	<b>37.2</b>	24.9	51.4	56	<b>45.6</b>	31.7	60.3	75
35-54	<b>32.9</b>	29.7	36.3	1,035	<b>34.2</b>	26.9	42.5	161	<b>35.3</b>	27.0	44.6	133	<b>32.7</b>	24.8	41.6	137
55+	<b>57.1</b>	54.0	60.2	1,409	<b>61.4</b>	53.6	68.7	195	<b>58.3</b>	51.2	65.2	218	<b>60.1</b>	53.2	66.7	238
<b>INCOME</b>																
Less than \$15,000	<b>64.2</b>	56.9	70.9	446	<b>65.2</b>	50.0	77.9	71	<b>48.8</b>	34.4	63.5	79	<b>74.1</b>	59.8	84.6	75
\$15,000 - \$24,999	<b>67.2</b>	61.4	72.6	782	<b>66.8</b>	54.6	77.1	118	<b>75.7</b>	64.3	84.3	123	<b>71.2</b>	58.8	81.0	140
\$25,000 - \$34,999	<b>51.7</b>	45.6	57.8	620	<b>55.3</b>	39.8	69.9	70	<b>53.7</b>	42.2	64.8	99	<b>50.7</b>	37.3	64.0	101
\$35,000 - \$49,999	<b>37.8</b>	33.8	42.0	819	<b>45.7</b>	34.5	57.2	108	<b>41.5</b>	31.3	52.4	126	<b>40.5</b>	31.0	50.8	124
\$50,000-\$74,999	<b>27.1</b>	23.5	31.1	757	<b>37.5</b>	27.6	48.6	111	<b>28.0</b>	19.0	39.3	105	<b>27.6</b>	18.4	39.2	98
\$75,000+	<b>24.0</b>	20.5	27.8	877	<b>30.7</b>	22.1	40.9	127	<b>25.1</b>	17.2	35.1	110	<b>22.0</b>	12.2	36.4	85
<b>EMPLOYMENT</b>																
Employed	<b>35.5</b>	32.9	38.1	2,585	<b>41.0</b>	34.6	47.6	355	<b>36.3</b>	30.6	42.5	367	<b>40.6</b>	34.5	47.1	359
Unemployed	<b>67.6</b>	56.6	76.9	161	*	*	*	*	*	*	*	*	*	*	*	*
Other**	<b>53.4</b>	50.3	56.4	2,062	<b>57.3</b>	50.5	63.8	302	<b>57.8</b>	50.4	65.0	312	<b>55.8</b>	47.6	63.8	313
<b>EDUCATION</b>																
K-11th Grade	<b>67.6</b>	60.4	74.1	374	<b>80.6</b>	65.2	90.2	52	*	*	*	*	<b>71.7</b>	56.6	83.1	84
12th Grade or GED	<b>47.8</b>	43.9	51.8	1,405	<b>49.6</b>	41.1	58.2	213	<b>50.1</b>	42.0	58.1	235	<b>53.1</b>	43.8	62.1	239
Some College	<b>42.7</b>	39.4	46.1	1,544	<b>45.9</b>	38.3	53.7	231	<b>44.0</b>	36.3	52.1	213	<b>46.6</b>	37.9	55.6	215
College Graduate+	<b>30.6</b>	27.7	33.7	1,483	<b>38.5</b>	30.9	46.7	187	<b>38.5</b>	30.8	47.0	210	<b>27.8</b>	20.8	36.0	164

\*Figure not reliable by BRFSS standards (n<50)

\*\*Other includes students, homemakers, retirees, and persons unable to work

# Dental Insurance

## Idaho adults without dental insurance, 2008

	Public Health District															
	District 4			District 5			District 6			District 7						
	%	95% CI		n	%	95% CI		n	%	95% CI		n	%	95% CI		n
<b>TOTAL</b>	<b>31.6</b>	27.4	36.2	710	<b>49.5</b>	44.7	54.2	680	<b>43.7</b>	38.4	49.1	664	<b>46.6</b>	41.2	52.1	665
<b>SEX</b>																
Male	<b>32.2</b>	25.5	39.8	273	<b>49.2</b>	41.6	56.8	268	<b>43.6</b>	35.8	51.7	238	<b>47.2</b>	38.8	55.7	273
Female	<b>31.0</b>	26.2	36.2	437	<b>49.7</b>	44.2	55.3	412	<b>43.8</b>	36.9	50.9	426	<b>46.1</b>	39.3	53.1	392
<b>AGE</b>																
18-24	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
25-34	<b>25.1</b>	15.9	37.2	87	<b>59.8</b>	46.9	71.4	74	<b>29.5</b>	19.5	41.9	89	<b>47.1</b>	35.7	58.9	105
35-44	<b>22.7</b>	15.9	31.4	141	<b>25.8</b>	17.6	36.1	99	<b>35.8</b>	25.4	47.7	81	<b>35.2</b>	26.5	45.0	108
45-54	<b>31.4</b>	23.6	40.5	139	<b>42.8</b>	34.2	51.9	148	<b>37.3</b>	28.7	46.9	137	<b>24.2</b>	17.4	32.6	134
55-64	<b>40.7</b>	32.5	49.5	159	<b>48.4</b>	39.5	57.5	135	<b>40.5</b>	32.4	49.1	156	<b>46.2</b>	36.7	56.0	117
65+	<b>46.0</b>	37.8	54.5	155	<b>75.6</b>	68.6	81.4	197	<b>73.9</b>	65.8	80.6	176	<b>77.1</b>	69.7	83.1	171
18-34	<b>28.1</b>	19.3	39.0	109	<b>52.8</b>	41.3	63.9	95	<b>38.1</b>	26.3	51.5	110	<b>51.1</b>	39.0	63.0	133
35-54	<b>26.8</b>	21.5	33.0	280	<b>34.9</b>	28.6	41.7	247	<b>36.6</b>	29.7	44.1	218	<b>29.5</b>	23.8	35.9	242
55+	<b>43.2</b>	37.4	49.3	314	<b>63.2</b>	57.3	68.7	332	<b>58.1</b>	52.1	63.8	332	<b>62.0</b>	55.6	68.1	288
<b>SEX and AGE</b>																
Male																
18-34	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
35-54	<b>23.8</b>	16.2	33.6	106	<b>34.0</b>	24.6	44.9	90	<b>35.8</b>	25.3	47.8	83	<b>28.1</b>	20.0	38.0	104
55+	<b>40.5</b>	31.7	49.9	128	<b>61.6</b>	52.7	69.9	142	<b>57.9</b>	48.0	67.1	116	<b>62.4</b>	52.3	71.5	122
Female																
18-34	<b>19.2</b>	10.9	31.7	71	<b>49.6</b>	36.8	62.4	61	<b>36.7</b>	20.7	56.2	72	<b>48.1</b>	33.4	63.0	87
35-54	<b>29.9</b>	23.0	37.9	174	<b>35.8</b>	28.1	44.3	157	<b>37.4</b>	29.0	46.7	135	<b>30.9</b>	23.4	39.4	138
55+	<b>45.7</b>	38.0	53.6	186	<b>64.6</b>	56.8	71.8	190	<b>58.3</b>	51.1	65.1	216	<b>61.7</b>	53.5	69.3	166
<b>INCOME</b>																
Less than \$15,000	<b>61.2</b>	42.9	76.8	50	<b>83.0</b>	69.5	91.3	57	<b>55.8</b>	33.6	75.8	63	<b>62.2</b>	38.2	81.3	51
\$15,000 - \$24,999	<b>60.7</b>	43.1	75.9	73	<b>71.3</b>	56.7	82.5	114	<b>61.8</b>	49.2	72.9	110	<b>68.7</b>	52.2	81.5	104
\$25,000 - \$34,999	<b>36.0</b>	23.4	50.9	88	<b>63.6</b>	51.8	73.9	93	<b>65.5</b>	51.0	77.7	95	<b>60.7</b>	44.0	75.2	74
\$35,000 - \$49,999	<b>22.6</b>	15.5	31.6	129	<b>41.9</b>	31.4	53.1	110	<b>38.2</b>	28.2	49.2	115	<b>51.5</b>	40.3	62.5	107
\$50,000-\$74,999	<b>16.3</b>	9.6	26.2	106	<b>34.4</b>	25.0	45.2	108	<b>26.3</b>	18.2	36.4	104	<b>30.0</b>	22.4	39.0	125
\$75,000+	<b>21.3</b>	15.2	29.0	196	<b>29.2</b>	20.8	39.3	112	<b>22.9</b>	15.0	33.3	111	<b>22.4</b>	15.4	31.3	136
<b>EMPLOYMENT</b>																
Employed	<b>24.7</b>	19.5	30.8	399	<b>44.2</b>	38.1	50.5	391	<b>34.4</b>	28.0	41.5	339	<b>40.3</b>	33.6	47.4	375
Unemployed	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Other**	<b>41.4</b>	34.7	48.4	280	<b>60.1</b>	52.4	67.3	273	<b>58.4</b>	49.9	66.5	309	<b>55.4</b>	45.6	64.8	273
<b>EDUCATION</b>																
K-11th Grade	*	*	*	*	<b>75.3</b>	61.4	85.4	66	*	*	*	*	*	*	*	*
12th Grade or GED	<b>37.9</b>	28.1	48.9	165	<b>48.0</b>	39.0	57.1	191	<b>52.2</b>	42.7	61.6	200	<b>55.3</b>	44.1	66.1	162
Some College	<b>34.1</b>	27.1	41.9	209	<b>51.0</b>	43.3	58.7	232	<b>39.9</b>	30.5	50.2	219	<b>44.4</b>	34.9	54.3	225
College Graduate+	<b>21.7</b>	16.5	27.9	299	<b>36.2</b>	28.9	44.2	189	<b>37.2</b>	29.6	45.4	196	<b>37.4</b>	30.0	45.3	238

\*Figure not reliable by BRFSS standards (n<50)

\*\*Other includes students, homemakers, retirees, and persons unable to work

# Diabetes

## Idaho adults who had ever been told they had diabetes, 2008

	Statewide			Public Health District												
				District 1			District 2			District 3						
	%	95% CI		n	%	95% CI		n	%	95% CI		n	%	95% CI		n
<b>TOTAL</b>	<b>7.0</b>	6.3	7.8	5,108	<b>6.4</b>	4.8	8.4	729	<b>7.9</b>	6.1	10.1	754	<b>7.1</b>	5.5	9.1	749
<b>SEX</b>																
Male	7.4	6.3	8.6	1,988	7.4	4.9	11.0	283	7.5	5.0	11.1	320	6.6	4.3	9.9	267
Female	6.7	5.8	7.6	3,120	5.3	3.6	7.8	446	8.3	6.1	11.2	434	7.6	5.6	10.4	482
<b>AGE</b>																
18-24	0.1	0.0	0.9	164	*	*	*	*	*	*	*	*	*	*	*	*
25-34	1.0	0.5	1.9	629	0.0	0.0	0.0	67	0.9	0.1	6.4	85	0.0	0.0	0.0	108
35-44	2.8	1.8	4.4	773	3.1	0.7	12.3	104	3.6	0.9	12.9	111	2.2	0.7	6.9	103
45-54	7.2	5.4	9.5	1,037	6.0	3.3	10.6	169	8.2	4.6	14.0	154	6.9	3.7	12.7	131
55-64	13.5	11.3	16.0	1,068	12.2	7.7	18.7	161	13.4	8.8	19.7	165	18.7	12.0	27.9	152
65+	19.2	16.8	21.8	1,395	14.3	9.5	21.0	199	19.5	14.0	26.6	215	19.6	14.4	26.1	223
18-34	0.7	0.4	1.3	793	0.0	0.0	0.0	89	0.7	0.1	4.8	106	0.0	0.0	0.0	130
35-54	5.0	3.9	6.4	1,810	4.7	2.6	8.4	273	6.1	3.5	10.3	265	4.5	2.6	7.7	234
55+	16.5	14.8	18.3	2,463	13.3	9.8	17.7	360	16.9	13.0	21.6	380	19.2	14.8	24.4	375
<b>SEX and AGE</b>																
Male																
18-34	0.6	0.2	1.7	290	*	*	*	*	*	*	*	*	0.0	0.0	0.0	52
35-54	5.7	4.0	8.2	727	4.6	1.6	12.1	104	4.4	1.5	12.2	127	4.4	1.9	10.0	88
55+	17.6	15.1	20.5	961	16.8	11.3	24.4	148	18.9	12.7	27.2	148	18.4	11.9	27.4	123
Female																
18-34	0.8	0.4	1.7	503	0.0	0.0	0.0	59	1.5	0.2	9.9	61	0.0	0.0	0.0	78
35-54	4.3	3.2	5.7	1,083	4.8	2.4	9.2	169	7.8	4.2	13.9	138	4.5	2.2	9.3	146
55+	15.5	13.4	17.8	1,502	10.0	6.3	15.5	212	14.9	10.7	20.4	232	19.9	14.6	26.4	252
<b>INCOME</b>																
Less than \$15,000	8.9	6.8	11.7	484	13.3	7.5	22.5	76	9.5	5.3	16.5	86	9.7	4.9	18.4	80
\$15,000 - \$24,999	10.6	8.6	13.1	829	5.6	2.9	10.6	120	12.7	7.5	20.7	134	11.4	6.8	18.5	148
\$25,000 - \$34,999	9.1	6.9	11.9	662	7.2	2.9	16.5	78	11.7	7.0	19.0	105	10.4	6.1	17.1	109
\$35,000 - \$49,999	5.9	4.5	7.8	852	4.6	2.2	9.5	116	6.8	3.4	13.1	128	5.5	2.7	10.8	130
\$50,000-\$74,999	4.1	2.9	5.8	794	6.0	2.8	12.2	114	1.3	0.3	5.3	110	3.4	1.3	9.0	106
\$75,000+	4.8	3.5	6.7	911	5.1	2.1	11.8	135	3.9	1.7	8.7	111	2.6	0.8	7.8	86
<b>EMPLOYMENT</b>																
Employed	4.3	3.5	5.2	2,722	3.7	2.2	6.2	378	4.2	2.6	6.6	388	4.4	2.8	7.0	382
Unemployed	7.9	4.7	12.9	176	*	*	*	*	*	*	*	*	*	*	*	*
Other**	11.8	10.4	13.4	2,202	10.4	7.4	14.6	321	13.2	9.8	17.5	335	11.7	8.6	15.8	335
<b>EDUCATION</b>																
K-11th Grade	6.5	4.7	9.0	406	5.9	2.5	13.2	54	7.8	2.8	20.4	54	9.7	5.2	17.3	93
12th Grade or GED	8.0	6.6	9.6	1,519	7.2	4.7	11.0	232	9.2	6.1	13.7	253	7.7	5.0	11.7	255
Some College	6.8	5.6	8.1	1,624	5.5	3.1	9.4	245	8.5	5.5	12.8	226	5.3	3.3	8.5	228
College Graduate+	6.5	5.3	8.1	1,548	6.7	3.6	12.1	195	5.8	3.6	9.4	220	7.0	4.0	12.0	171

\*Figure not reliable by BRFSS standards (n<50)

\*\*Other includes students, homemakers, retirees, and persons unable to work

# Diabetes

## Idaho adults who had ever been told they had diabetes, 2008

	Public Health District															
	District 4			District 5			District 6			District 7						
	%	95% CI		n	%	95% CI		n	%	95% CI		n	%	95% CI		n
<b>TOTAL</b>	<b>6.0</b>	4.6	8.0	751	<b>7.7</b>	5.9	9.8	726	<b>8.9</b>	6.8	11.5	704	<b>7.1</b>	5.5	9.1	695
<b>SEX</b>																
Male	<b>6.8</b>	4.5	10.1	294	<b>8.9</b>	6.3	12.5	283	<b>8.1</b>	5.3	12.2	256	<b>7.4</b>	5.0	10.7	285
Female	<b>5.3</b>	3.7	7.6	457	<b>6.4</b>	4.5	9.1	443	<b>9.6</b>	6.9	13.2	448	<b>6.8</b>	4.9	9.4	410
<b>AGE</b>																
18-24	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
25-34	<b>0.0</b>	0.0	0.0	87	<b>0.9</b>	0.1	6.0	78	<b>5.7</b>	2.1	14.5	93	<b>2.6</b>	0.8	8.1	111
35-44	<b>1.2</b>	0.2	5.9	149	<b>1.9</b>	0.5	7.4	107	<b>8.1</b>	3.5	17.6	88	<b>3.6</b>	1.4	8.8	111
45-54	<b>8.3</b>	4.3	15.7	150	<b>6.1</b>	2.8	12.8	153	<b>9.5</b>	5.2	16.8	141	<b>4.6</b>	2.3	9.2	139
55-64	<b>10.0</b>	6.1	16.2	164	<b>17.3</b>	11.7	25.0	141	<b>14.3</b>	9.4	21.1	163	<b>12.4</b>	7.5	19.8	122
65+	<b>21.2</b>	15.3	28.6	171	<b>19.4</b>	14.2	25.9	218	<b>16.8</b>	11.8	23.5	190	<b>24.0</b>	17.9	31.5	179
18-34	<b>0.0</b>	0.0	0.0	110	<b>0.7</b>	0.1	4.5	100	<b>3.0</b>	1.1	8.0	117	<b>1.6</b>	0.6	4.3	141
35-54	<b>4.6</b>	2.5	8.5	299	<b>4.1</b>	2.1	8.0	260	<b>8.8</b>	5.4	14.2	229	<b>4.1</b>	2.4	7.2	250
55+	<b>15.5</b>	11.7	20.1	335	<b>18.5</b>	14.5	23.3	359	<b>15.6</b>	11.9	20.2	353	<b>18.4</b>	14.2	23.5	301
<b>SEX and AGE</b>																
Male																
18-34	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
35-54	<b>7.7</b>	3.8	14.8	118	<b>5.7</b>	2.3	13.3	95	<b>5.9</b>	2.3	14.3	88	<b>4.1</b>	1.7	9.7	107
55+	<b>14.2</b>	9.2	21.4	137	<b>22.4</b>	16.1	30.2	150	<b>18.0</b>	12.0	26.1	127	<b>18.4</b>	12.4	26.5	128
Female																
18-34	<b>0.0</b>	0.0	0.0	72	<b>1.4</b>	0.2	9.1	64	<b>3.6</b>	1.2	10.3	77	<b>0.6</b>	0.1	4.3	92
35-54	<b>1.3</b>	0.4	4.7	181	<b>2.4</b>	0.9	6.2	165	<b>11.8</b>	6.7	20.0	141	<b>4.2</b>	2.1	8.2	143
55+	<b>16.6</b>	11.6	23.2	198	<b>15.0</b>	10.3	21.3	209	<b>13.5</b>	9.5	18.8	226	<b>18.3</b>	13.0	25.2	173
<b>INCOME</b>																
Less than \$15,000	<b>8.6</b>	3.9	18.0	55	<b>6.0</b>	2.7	12.5	67	<b>12.0</b>	6.2	22.0	68	<b>3.3</b>	1.1	9.6	52
\$15,000 - \$24,999	<b>8.1</b>	4.2	15.1	81	<b>16.2</b>	10.2	24.7	123	<b>14.0</b>	7.9	23.8	114	<b>11.7</b>	6.9	19.2	109
\$25,000 - \$34,999	<b>9.7</b>	4.7	18.7	88	<b>10.2</b>	6.0	16.9	101	<b>7.9</b>	4.0	15.2	101	<b>6.1</b>	2.7	13.0	80
\$35,000 - \$49,999	<b>5.5</b>	2.7	10.9	133	<b>5.5</b>	2.5	11.7	113	<b>8.8</b>	4.6	16.2	121	<b>6.5</b>	3.3	12.4	111
\$50,000-\$74,999	<b>4.4</b>	2.0	9.3	116	<b>4.2</b>	1.8	9.4	109	<b>3.0</b>	1.3	6.9	111	<b>4.7</b>	2.2	9.9	128
\$75,000+	<b>4.6</b>	2.4	8.8	201	<b>4.6</b>	1.8	11.2	119	<b>7.8</b>	3.7	15.8	118	<b>5.2</b>	2.7	10.1	141
<b>EMPLOYMENT</b>																
Employed	<b>4.0</b>	2.5	6.4	417	<b>3.9</b>	2.4	6.3	410	<b>5.9</b>	3.7	9.1	358	<b>4.8</b>	3.2	7.3	389
Unemployed	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Other**	<b>10.9</b>	7.9	14.9	300	<b>14.5</b>	10.9	19.1	297	<b>13.9</b>	10.0	19.0	328	<b>10.4</b>	7.4	14.4	286
<b>EDUCATION</b>																
K-11th Grade	*	*	*	*	<b>5.1</b>	2.3	10.6	71	<b>6.7</b>	2.8	15.4	51	*	*	*	*
12th Grade or GED	<b>6.7</b>	3.7	11.7	173	<b>8.5</b>	5.5	12.9	208	<b>7.9</b>	4.9	12.3	219	<b>11.4</b>	7.6	16.6	179
Some College	<b>6.5</b>	4.1	10.4	220	<b>8.6</b>	5.6	13.0	246	<b>11.2</b>	7.4	16.6	230	<b>4.2</b>	2.4	7.4	229
College Graduate+	<b>5.8</b>	3.7	8.9	317	<b>6.4</b>	3.6	11.2	199	<b>8.2</b>	4.6	14.2	202	<b>7.4</b>	4.8	11.3	244

\*Figure not reliable by BRFSS standards (n<50)

\*\*Other includes students, homemakers, retirees, and persons unable to work

# Asthma

## Idaho adults who were diagnosed and still had asthma, 2008

	Statewide			Public Health District												
				District 1			District 2			District 3						
	%	95% CI		n	%	95% CI		n	%	95% CI		n	%	95% CI		n
<b>TOTAL</b>	<b>8.9</b>	7.8	10.1	5,079	<b>9.5</b>	7.2	12.5	724	<b>11.5</b>	9.0	14.6	752	<b>10.3</b>	7.7	13.6	743
<b>SEX</b>																
Male	<b>7.6</b>	6.0	9.5	1,981	<b>6.1</b>	3.2	11.3	281	<b>8.1</b>	5.1	12.7	319	<b>8.9</b>	5.8	13.6	266
Female	<b>10.2</b>	8.8	11.6	3,098	<b>12.9</b>	9.6	16.9	443	<b>14.9</b>	11.2	19.5	433	<b>11.6</b>	7.8	16.9	477
<b>AGE</b>																
18-24	<b>7.2</b>	3.5	14.3	162	*	*	*	*	*	*	*	*	*	*	*	*
25-34	<b>8.2</b>	5.7	11.6	626	<b>6.2</b>	2.4	15.2	67	<b>11.9</b>	6.3	21.2	85	<b>7.7</b>	3.6	15.8	106
35-44	<b>8.8</b>	6.6	11.6	769	<b>7.9</b>	4.0	14.9	104	<b>9.4</b>	5.1	16.7	111	<b>9.4</b>	4.7	18.0	102
45-54	<b>10.0</b>	8.0	12.4	1,033	<b>10.8</b>	6.8	16.6	168	<b>15.4</b>	10.0	22.9	153	<b>16.5</b>	10.3	25.2	131
55-64	<b>11.0</b>	9.0	13.4	1,059	<b>12.3</b>	7.8	18.9	160	<b>12.5</b>	7.9	19.3	164	<b>13.3</b>	8.2	20.8	150
65+	<b>7.8</b>	6.4	9.5	1,388	<b>9.4</b>	5.8	14.9	196	<b>9.2</b>	5.7	14.6	215	<b>8.8</b>	5.3	14.5	222
18-34	<b>7.8</b>	5.6	10.9	788	<b>8.3</b>	3.5	18.6	89	<b>11.2</b>	6.3	19.2	106	<b>7.3</b>	3.1	16.4	128
35-54	<b>9.4</b>	7.9	11.2	1,802	<b>9.4</b>	6.4	13.6	272	<b>12.6</b>	8.9	17.7	264	<b>12.8</b>	8.6	18.5	233
55+	<b>9.3</b>	8.1	10.7	2,447	<b>10.8</b>	7.8	14.8	356	<b>10.7</b>	7.7	14.7	379	<b>10.8</b>	7.6	15.2	372
<b>SEX and AGE</b>																
Male																
18-34	<b>7.2</b>	4.0	12.6	289	*	*	*	*	*	*	*	*	<b>3.9</b>	1.0	14.6	52
35-54	<b>8.0</b>	5.9	10.8	727	<b>6.0</b>	2.8	12.4	104	<b>8.2</b>	4.2	15.4	127	<b>13.0</b>	7.1	22.7	87
55+	<b>7.5</b>	5.8	9.7	955	<b>6.6</b>	3.4	12.2	146	<b>7.8</b>	4.1	14.3	147	<b>10.6</b>	5.8	18.6	123
Female																
18-34	<b>8.5</b>	5.8	12.2	499	<b>10.9</b>	4.7	23.2	59	<b>14.5</b>	7.3	26.8	61	<b>11.0</b>	3.8	27.8	76
35-54	<b>10.8</b>	8.8	13.2	1,075	<b>12.8</b>	8.3	19.2	168	<b>17.1</b>	11.3	25.0	137	<b>12.5</b>	7.6	20.0	146
55+	<b>11.0</b>	9.3	12.9	1,492	<b>14.7</b>	10.1	20.9	210	<b>13.4</b>	9.2	19.0	232	<b>11.0</b>	7.4	16.3	249
<b>INCOME</b>																
Less than \$15,000	<b>11.8</b>	8.7	16.0	479	<b>13.7</b>	6.6	26.0	75	<b>29.2</b>	17.8	44.1	86	<b>8.3</b>	3.8	17.2	79
\$15,000 - \$24,999	<b>9.6</b>	7.4	12.2	823	<b>10.6</b>	6.2	17.5	120	<b>9.6</b>	5.7	15.7	134	<b>13.9</b>	8.3	22.5	146
\$25,000 - \$34,999	<b>9.5</b>	6.0	14.7	658	<b>13.0</b>	6.2	25.4	78	<b>9.9</b>	5.2	18.0	103	<b>14.7</b>	6.2	30.7	108
\$35,000 - \$49,999	<b>8.5</b>	6.5	11.1	850	<b>11.1</b>	6.2	18.9	116	<b>5.2</b>	2.5	10.4	128	<b>12.8</b>	7.2	21.6	128
\$50,000-\$74,999	<b>8.0</b>	5.4	11.8	793	<b>3.1</b>	1.2	7.8	114	<b>11.2</b>	5.3	22.3	110	<b>12.2</b>	6.4	22.2	106
\$75,000+	<b>8.0</b>	6.0	10.6	907	<b>6.3</b>	3.3	11.7	134	<b>9.1</b>	4.7	17.0	111	<b>4.3</b>	1.7	10.3	86
<b>EMPLOYMENT</b>																
Employed	<b>8.2</b>	6.7	9.8	2,710	<b>7.6</b>	4.8	11.9	376	<b>10.3</b>	7.4	14.3	387	<b>9.7</b>	6.8	13.7	379
Unemployed	<b>9.7</b>	5.8	15.9	172	*	*	*	*	*	*	*	*	*	*	*	*
Other**	<b>10.0</b>	8.4	11.9	2,189	<b>12.4</b>	8.9	17.2	318	<b>12.3</b>	8.4	17.8	334	<b>11.9</b>	7.0	19.5	334
<b>EDUCATION</b>																
K-11th Grade	<b>8.7</b>	5.8	12.9	402	<b>21.5</b>	8.8	43.5	54	<b>8.9</b>	3.1	23.1	54	<b>8.1</b>	4.1	15.3	91
12th Grade or GED	<b>9.4</b>	7.1	12.3	1,509	<b>9.2</b>	5.9	13.9	230	<b>10.6</b>	7.0	15.8	252	<b>11.6</b>	6.3	20.5	254
Some College	<b>10.0</b>	8.3	12.0	1,619	<b>9.0</b>	6.0	13.3	245	<b>16.3</b>	11.1	23.3	226	<b>11.1</b>	7.1	17.0	226
College Graduate+	<b>7.2</b>	5.7	9.2	1,539	<b>6.6</b>	3.5	11.9	192	<b>8.7</b>	4.9	15.0	219	<b>8.9</b>	5.0	15.2	171

\*Figure not reliable by BRFSS standards (n<50)

\*\*Other includes students, homemakers, retirees, and persons unable to work

# Asthma

## Idaho adults who were diagnosed and still had asthma, 2008

	Public Health District															
	District 4				District 5				District 6				District 7			
	%	95% CI		n	%	95% CI		n	%	95% CI		n	%	95% CI		n
<b>TOTAL</b>	<b>9.9</b>	7.3	13.2	750	<b>7.1</b>	5.3	9.4	723	<b>6.5</b>	4.6	9.2	699	<b>6.0</b>	4.3	8.3	688
<b>SEX</b>																
Male	<b>10.9</b>	6.8	17.1	294	<b>6.2</b>	3.8	9.8	282	<b>2.7</b>	1.3	5.4	256	<b>4.9</b>	2.7	8.7	283
Female	<b>8.8</b>	6.2	12.2	456	<b>8.1</b>	5.6	11.4	441	<b>10.3</b>	6.9	15.1	443	<b>7.1</b>	4.8	10.2	405
<b>AGE</b>																
18-24	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
25-34	<b>11.9</b>	6.0	22.2	87	<b>4.4</b>	1.6	12.0	78	<b>5.2</b>	2.1	12.5	93	<b>3.6</b>	1.3	9.6	110
35-44	<b>11.4</b>	6.7	18.7	149	<b>6.5</b>	2.9	13.9	106	<b>3.9</b>	1.1	13.7	86	<b>7.7</b>	3.8	15.2	111
45-54	<b>9.4</b>	5.4	15.9	149	<b>6.1</b>	3.3	11.0	152	<b>7.1</b>	3.9	12.8	141	<b>6.0</b>	3.1	11.3	139
55-64	<b>9.9</b>	5.9	16.2	164	<b>9.3</b>	5.4	15.4	139	<b>8.8</b>	5.3	14.2	162	<b>11.7</b>	7.0	19.1	120
65+	<b>4.7</b>	2.5	8.7	171	<b>10.5</b>	6.8	15.8	219	<b>7.5</b>	4.7	11.9	188	<b>6.1</b>	3.1	11.6	177
18-34	<b>10.9</b>	5.6	20.3	110	<b>5.2</b>	2.3	11.6	100	<b>6.1</b>	2.6	13.9	117	<b>3.1</b>	1.1	8.5	138
35-54	<b>10.4</b>	7.1	15.1	298	<b>6.3</b>	3.8	10.2	258	<b>5.6</b>	3.1	10.0	227	<b>6.8</b>	4.2	11.0	250
55+	<b>7.4</b>	4.9	11.0	335	<b>10.0</b>	7.1	13.7	358	<b>8.1</b>	5.7	11.4	350	<b>8.9</b>	5.9	13.1	297
<b>SEX and AGE</b>																
Male																
18-34	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
35-54	<b>10.9</b>	6.2	18.6	118	<b>5.1</b>	2.1	12.2	95	<b>0.9</b>	0.1	6.4	88	<b>5.4</b>	2.4	11.9	108
55+	<b>5.9</b>	2.7	12.5	137	<b>11.5</b>	7.0	18.2	149	<b>4.3</b>	2.0	9.0	127	<b>6.8</b>	3.4	13.0	126
Female																
18-34	<b>6.9</b>	2.8	15.7	72	<b>8.2</b>	3.4	18.6	64	<b>9.1</b>	3.1	23.6	77	<b>3.2</b>	1.1	8.8	90
35-54	<b>9.9</b>	6.0	16.0	180	<b>7.6</b>	4.3	12.9	163	<b>10.4</b>	5.7	18.4	139	<b>8.3</b>	4.5	14.7	142
55+	<b>8.6</b>	5.4	13.5	198	<b>8.6</b>	5.5	13.3	209	<b>11.6</b>	8.0	16.7	223	<b>10.7</b>	6.4	17.3	171
<b>INCOME</b>																
Less than \$15,000	<b>11.7</b>	4.1	28.7	54	<b>11.8</b>	6.1	21.7	67	<b>11.2</b>	5.4	21.9	68	<b>3.7</b>	1.3	10.1	50
\$15,000 - \$24,999	<b>7.5</b>	3.4	15.7	81	<b>11.4</b>	6.6	19.0	122	<b>6.0</b>	2.1	15.9	114	<b>6.4</b>	3.0	13.1	106
\$25,000 - \$34,999	<b>9.7</b>	2.7	29.4	88	<b>2.5</b>	0.4	12.5	100	<b>7.9</b>	3.1	19.0	101	<b>5.8</b>	2.5	13.0	80
\$35,000 - \$49,999	<b>10.5</b>	6.0	17.8	133	<b>7.3</b>	3.2	15.5	113	<b>2.5</b>	0.9	6.8	120	<b>4.0</b>	1.5	10.2	112
\$50,000-\$74,999	<b>12.3</b>	5.5	25.4	116	<b>2.5</b>	1.0	6.1	108	<b>6.2</b>	2.8	12.9	111	<b>4.4</b>	1.9	9.8	128
\$75,000+	<b>10.5</b>	6.5	16.5	201	<b>7.8</b>	3.9	15.0	118	<b>7.4</b>	3.0	17.5	116	<b>5.6</b>	2.9	10.8	141
<b>EMPLOYMENT</b>																
Employed	<b>10.2</b>	6.8	15.2	416	<b>5.4</b>	3.5	8.3	407	<b>4.9</b>	2.6	9.2	357	<b>6.1</b>	3.9	9.4	388
Unemployed	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Other**	<b>9.5</b>	6.2	14.4	300	<b>9.8</b>	6.5	14.4	298	<b>8.4</b>	5.6	12.5	324	<b>5.6</b>	3.4	9.0	281
<b>EDUCATION</b>																
K-11th Grade	*	*	*	*	<b>6.9</b>	2.9	15.7	70	<b>5.0</b>	2.1	11.2	51	*	*	*	*
12th Grade or GED	<b>13.7</b>	7.3	24.2	172	<b>4.3</b>	2.4	7.6	206	<b>3.5</b>	2.0	6.0	218	<b>7.1</b>	3.4	14.3	177
Some College	<b>11.9</b>	7.8	17.7	220	<b>9.2</b>	5.8	14.4	247	<b>9.1</b>	5.1	15.7	229	<b>4.6</b>	2.6	8.0	226
College Graduate+	<b>6.8</b>	3.9	11.4	317	<b>7.5</b>	4.3	12.9	198	<b>7.5</b>	3.7	14.5	199	<b>6.8</b>	4.1	11.0	243

\*Figure not reliable by BRFSS standards (n<50)

\*\*Other includes students, homemakers, retirees, and persons unable to work

# Physical Activity

Idaho adults who did not participate in leisure time physical activity, 2008

	Statewide			Public Health District												
				District 1			District 2			District 3						
	%	95% CI		n	%	95% CI		n	%	95% CI		n	%	95% CI		n
<b>TOTAL</b>	<b>21.1</b>	19.7	22.7	5,104	<b>24.5</b>	20.9	28.6	729	<b>21.2</b>	17.9	25.0	752	<b>25.6</b>	21.9	29.7	748
<b>SEX</b>																
Male	<b>20.0</b>	17.8	22.5	1,990	<b>24.0</b>	18.5	30.5	283	<b>18.6</b>	14.0	24.4	320	<b>26.2</b>	20.5	32.8	268
Female	<b>22.3</b>	20.5	24.1	3,114	<b>25.1</b>	20.6	30.2	446	<b>23.9</b>	19.4	29.0	432	<b>25.0</b>	20.5	30.2	480
<b>AGE</b>																
18-24	<b>11.4</b>	7.3	17.4	164	*	*	*	*	*	*	*	*	*	*	*	*
25-34	<b>17.0</b>	13.3	21.4	629	<b>26.5</b>	16.1	40.5	67	<b>10.7</b>	5.6	19.4	85	<b>16.0</b>	10.1	24.5	108
35-44	<b>18.0</b>	15.2	21.3	773	<b>21.0</b>	13.6	30.8	104	<b>21.2</b>	13.7	31.2	111	<b>29.8</b>	20.8	40.6	103
45-54	<b>22.5</b>	19.7	25.7	1,037	<b>26.3</b>	19.6	34.4	169	<b>25.0</b>	18.3	33.1	154	<b>29.4</b>	21.4	38.9	131
55-64	<b>25.0</b>	22.1	28.2	1,068	<b>22.8</b>	16.7	30.4	161	<b>25.1</b>	18.5	33.1	165	<b>35.3</b>	27.0	44.7	152
65+	<b>32.1</b>	29.2	35.1	1,391	<b>31.3</b>	24.6	39.0	199	<b>30.9</b>	24.1	38.5	213	<b>32.4</b>	25.7	39.9	222
18-34	<b>15.0</b>	12.1	18.4	793	<b>21.9</b>	13.8	33.0	89	<b>12.2</b>	6.7	21.3	106	<b>14.9</b>	9.2	23.1	130
35-54	<b>20.3</b>	18.3	22.6	1,810	<b>23.8</b>	18.6	29.9	273	<b>23.3</b>	18.1	29.4	265	<b>29.6</b>	23.3	36.7	234
55+	<b>28.8</b>	26.7	30.9	2,459	<b>27.3</b>	22.5	32.6	360	<b>28.3</b>	23.4	33.8	378	<b>33.7</b>	28.3	39.5	374
<b>SEX and AGE</b>																
Male																
18-34	<b>16.8</b>	12.1	22.8	290	*	*	*	*	*	*	*	*	<b>10.8</b>	4.3	24.4	52
35-54	<b>19.4</b>	16.4	22.7	728	<b>27.8</b>	19.5	38.0	104	<b>20.6</b>	14.0	29.3	127	<b>33.9</b>	24.0	45.5	88
55+	<b>24.8</b>	21.8	28.1	962	<b>23.6</b>	17.0	31.7	148	<b>26.3</b>	19.0	35.1	148	<b>36.2</b>	27.4	46.0	124
Female																
18-34	<b>13.1</b>	10.3	16.7	503	<b>24.1</b>	13.8	38.8	59	<b>14.9</b>	7.5	27.3	61	<b>19.2</b>	11.1	31.2	78
35-54	<b>21.3</b>	18.6	24.3	1,082	<b>19.9</b>	14.2	27.2	169	<b>25.9</b>	18.5	35.1	138	<b>25.3</b>	18.3	33.8	146
55+	<b>32.3</b>	29.5	35.2	1,497	<b>30.7</b>	24.3	38.0	212	<b>30.3</b>	24.1	37.2	230	<b>31.5</b>	25.4	38.3	250
<b>INCOME</b>																
Less than \$15,000	<b>39.6</b>	33.2	46.4	484	<b>37.6</b>	25.1	52.2	76	<b>27.6</b>	17.8	40.3	86	<b>46.2</b>	31.9	61.2	80
\$15,000 - \$24,999	<b>30.3</b>	25.9	35.1	829	<b>41.0</b>	30.1	52.8	120	<b>28.8</b>	20.2	39.4	133	<b>31.4</b>	22.8	41.6	149
\$25,000 - \$34,999	<b>24.7</b>	20.2	29.8	661	<b>30.7</b>	18.6	46.2	78	<b>29.3</b>	20.2	40.3	105	<b>32.4</b>	21.4	45.8	108
\$35,000 - \$49,999	<b>19.7</b>	16.8	23.0	852	<b>22.1</b>	15.0	31.4	116	<b>16.8</b>	11.1	24.7	128	<b>25.1</b>	17.8	34.2	130
\$50,000-\$74,999	<b>13.7</b>	11.0	16.8	793	<b>8.3</b>	4.4	15.2	114	<b>10.6</b>	6.0	18.1	110	<b>18.6</b>	11.4	28.9	105
\$75,000+	<b>9.8</b>	7.6	12.5	911	<b>13.9</b>	8.8	21.2	135	<b>10.0</b>	5.8	16.9	111	<b>8.8</b>	4.2	17.6	86
<b>EMPLOYMENT</b>																
Employed	<b>19.3</b>	17.4	21.3	2,722	<b>19.0</b>	14.8	24.1	378	<b>16.4</b>	12.7	20.9	388	<b>25.3</b>	20.4	31.0	382
Unemployed	<b>18.3</b>	12.5	26.1	176	*	*	*	*	*	*	*	*	*	*	*	*
Other**	<b>25.0</b>	22.7	27.4	2,198	<b>32.3</b>	26.3	38.9	321	<b>29.4</b>	23.3	36.3	333	<b>26.8</b>	21.2	33.3	334
<b>EDUCATION</b>																
K-11th Grade	<b>36.5</b>	30.3	43.2	407	<b>39.6</b>	23.8	57.9	54	<b>34.7</b>	20.1	53.0	54	<b>36.8</b>	25.3	50.1	93
12th Grade or GED	<b>27.3</b>	24.3	30.5	1,516	<b>29.1</b>	22.7	36.4	232	<b>24.8</b>	18.9	31.8	251	<b>31.7</b>	24.6	39.6	255
Some College	<b>19.3</b>	16.9	21.9	1,623	<b>22.2</b>	16.4	29.3	245	<b>21.8</b>	16.3	28.4	226	<b>18.6</b>	13.2	25.7	228
College Graduate+	<b>11.8</b>	10.0	13.8	1,547	<b>14.8</b>	10.0	21.3	195	<b>11.7</b>	7.8	17.1	220	<b>17.9</b>	12.2	25.4	170

\*Figure not reliable by BRFSS standards (n<50)

\*\*Other includes students, homemakers, retirees, and persons unable to work

# Physical Activity

Idaho adults who did not participate in leisure time physical activity, 2008

	Public Health District															
	District 4			District 5			District 6			District 7						
	%	95% CI		n	%	95% CI		n	%	95% CI		n	%	95% CI		n
<b>TOTAL</b>	<b>15.5</b>	12.6	18.9	751	<b>23.3</b>	19.5	27.6	725	<b>22.3</b>	18.5	26.6	704	<b>21.1</b>	17.4	25.4	695
<b>SEX</b>																
Male	<b>13.0</b>	8.7	18.8	294	<b>18.4</b>	12.7	26.0	283	<b>25.4</b>	19.2	32.7	256	<b>21.7</b>	15.8	29.1	286
Female	<b>18.1</b>	14.6	22.2	457	<b>28.2</b>	23.7	33.2	442	<b>19.4</b>	15.4	24.0	448	<b>20.5</b>	16.4	25.5	409
<b>AGE</b>																
18-24	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
25-34	<b>12.3</b>	5.8	24.3	87	<b>26.9</b>	16.4	40.9	78	<b>11.9</b>	5.9	22.3	93	<b>18.3</b>	10.5	29.9	111
35-44	<b>10.1</b>	6.2	16.0	149	<b>16.7</b>	10.7	25.1	107	<b>24.0</b>	15.9	34.5	88	<b>14.5</b>	8.9	22.7	111
45-54	<b>16.2</b>	10.7	23.8	150	<b>18.8</b>	12.7	26.8	152	<b>28.5</b>	20.9	37.6	141	<b>21.5</b>	15.0	29.9	140
55-64	<b>20.5</b>	14.5	28.2	164	<b>23.5</b>	16.8	32.0	141	<b>24.1</b>	17.7	32.0	163	<b>28.4</b>	20.7	37.6	122
65+	<b>33.3</b>	25.8	41.7	171	<b>30.3</b>	24.2	37.2	218	<b>31.3</b>	24.5	39.0	190	<b>34.6</b>	27.5	42.5	178
18-34	<b>8.9</b>	4.1	18.1	110	<b>25.4</b>	16.4	37.1	100	<b>13.5</b>	7.1	24.0	117	<b>16.0</b>	9.3	26.1	141
35-54	<b>13.0</b>	9.5	17.6	299	<b>17.8</b>	13.3	23.4	259	<b>26.4</b>	20.6	33.1	229	<b>18.2</b>	13.6	23.8	251
55+	<b>26.7</b>	21.7	32.4	335	<b>27.3</b>	22.6	32.6	359	<b>27.9</b>	23.0	33.3	353	<b>31.6</b>	26.2	37.6	300
<b>SEX and AGE</b>																
Male																
18-34	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
35-54	<b>7.5</b>	4.0	13.5	118	<b>12.1</b>	6.7	21.1	95	<b>29.9</b>	20.8	40.9	88	<b>18.9</b>	12.3	28.0	108
55+	<b>18.2</b>	12.0	26.6	137	<b>18.6</b>	12.8	26.1	150	<b>27.3</b>	19.9	36.3	127	<b>29.7</b>	21.9	38.9	128
Female																
18-34	<b>1.3</b>	0.3	4.9	72	<b>25.7</b>	16.2	38.1	64	<b>7.6</b>	3.0	18.0	77	<b>13.4</b>	7.2	23.5	92
35-54	<b>19.0</b>	13.3	26.3	181	<b>23.6</b>	17.3	31.3	164	<b>22.9</b>	16.4	31.0	141	<b>17.5</b>	11.8	25.0	143
55+	<b>34.3</b>	27.3	42.1	198	<b>35.1</b>	28.4	42.5	209	<b>28.4</b>	22.5	35.2	226	<b>33.3</b>	26.2	41.3	172
<b>INCOME</b>																
Less than \$15,000	<b>37.3</b>	21.3	56.7	55	<b>57.8</b>	42.6	71.7	67	<b>49.6</b>	29.9	69.4	68	<b>25.7</b>	13.4	43.6	52
\$15,000 - \$24,999	<b>20.8</b>	12.4	32.9	81	<b>39.8</b>	26.8	54.4	123	<b>22.3</b>	14.5	32.8	114	<b>30.0</b>	17.9	45.8	109
\$25,000 - \$34,999	<b>15.7</b>	8.0	28.4	88	<b>25.5</b>	16.5	37.4	101	<b>24.9</b>	14.5	39.3	101	<b>23.7</b>	12.8	39.8	80
\$35,000 - \$49,999	<b>18.8</b>	12.6	27.2	133	<b>14.2</b>	8.8	22.1	112	<b>17.8</b>	10.9	27.8	121	<b>20.6</b>	13.5	30.3	112
\$50,000-\$74,999	<b>11.4</b>	6.2	20.1	116	<b>14.2</b>	8.3	23.3	109	<b>17.0</b>	10.7	26.0	111	<b>17.9</b>	11.9	26.0	128
\$75,000+	<b>8.2</b>	4.4	14.7	201	<b>4.6</b>	1.7	11.5	119	<b>14.1</b>	8.5	22.6	118	<b>11.7</b>	7.2	18.4	141
<b>EMPLOYMENT</b>																
Employed	<b>14.2</b>	10.4	18.9	417	<b>22.0</b>	17.0	28.0	409	<b>23.7</b>	18.4	29.9	358	<b>19.6</b>	15.1	25.0	390
Unemployed	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Other**	<b>20.6</b>	16.1	26.0	300	<b>27.2</b>	21.9	33.3	297	<b>20.7</b>	16.0	26.3	328	<b>21.3</b>	15.4	28.6	285
<b>EDUCATION</b>																
K-11th Grade	*	*	*	*	<b>52.6</b>	36.9	67.8	71	<b>28.6</b>	15.5	46.6	51	*	*	*	*
12th Grade or GED	<b>23.1</b>	15.9	32.2	173	<b>21.8</b>	16.4	28.4	207	<b>35.8</b>	27.4	45.3	219	<b>26.5</b>	19.1	35.4	179
Some College	<b>17.1</b>	11.9	24.0	220	<b>24.3</b>	18.6	31.2	246	<b>17.2</b>	12.6	23.1	230	<b>16.9</b>	11.3	24.6	228
College Graduate+	<b>7.9</b>	5.4	11.5	317	<b>9.2</b>	5.9	14.1	199	<b>10.7</b>	6.7	16.8	202	<b>17.8</b>	12.4	24.8	244

\*Figure not reliable by BRFSS standards (n<50)

\*\*Other includes students, homemakers, retirees, and persons unable to work

# Overweight

## Idaho adults who were overweight (BMI ≥ 25), 2008

	Statewide			Public Health District												
				District 1			District 2			District 3						
	%	95% CI		n	%	95% CI		n	%	95% CI		n	%	95% CI		n
<b>TOTAL</b>	<b>62.2</b>	60.2	64.2	4,851	<b>63.3</b>	58.6	67.8	685	<b>63.5</b>	58.8	67.9	717	<b>67.2</b>	62.3	71.7	712
<b>SEX</b>																
Male	<b>68.4</b>	65.2	71.5	1,962	<b>67.8</b>	60.1	74.6	279	<b>68.6</b>	61.1	75.3	315	<b>72.7</b>	64.6	79.5	263
Female	<b>55.6</b>	53.1	58.1	2,889	<b>58.7</b>	53.1	64.2	406	<b>57.8</b>	51.9	63.5	402	<b>61.5</b>	55.3	67.3	449
<b>AGE</b>																
18-24	<b>42.5</b>	33.7	51.8	157	*	*	*	*	*	*	*	*	*	*	*	*
25-34	<b>55.0</b>	49.8	60.1	596	<b>64.5</b>	50.4	76.6	62	<b>58.2</b>	46.5	69.0	80	<b>64.0</b>	53.5	73.3	100
35-44	<b>65.6</b>	61.5	69.4	732	<b>58.9</b>	47.8	69.1	96	<b>70.1</b>	60.1	78.5	104	<b>72.6</b>	62.7	80.7	99
45-54	<b>70.1</b>	66.6	73.4	976	<b>64.2</b>	56.0	71.7	162	<b>74.9</b>	66.3	81.9	145	<b>74.7</b>	65.6	82.1	120
55-64	<b>73.2</b>	70.0	76.2	1,017	<b>73.0</b>	65.2	79.6	156	<b>75.5</b>	67.7	81.9	156	<b>79.6</b>	71.5	85.9	147
65+	<b>64.2</b>	61.2	67.1	1,349	<b>66.1</b>	58.3	73.1	187	<b>65.0</b>	57.5	71.9	211	<b>60.8</b>	53.2	67.9	216
18-34	<b>50.5</b>	45.8	55.1	753	<b>57.2</b>	44.6	68.9	82	<b>48.3</b>	37.9	58.8	100	<b>59.0</b>	47.8	69.4	122
35-54	<b>67.8</b>	65.2	70.4	1,708	<b>61.8</b>	55.1	68.1	258	<b>72.7</b>	66.3	78.3	249	<b>73.6</b>	66.9	79.3	219
55+	<b>68.4</b>	66.2	70.5	2,366	<b>69.4</b>	64.0	74.3	343	<b>69.5</b>	64.1	74.4	367	<b>69.2</b>	63.7	74.2	363
<b>SEX and AGE</b>																
Male																
18-34	<b>52.1</b>	44.9	59.3	280	*	*	*	*	*	*	*	*	*	*	*	*
35-54	<b>78.0</b>	74.2	81.4	717	<b>71.0</b>	60.6	79.6	104	<b>82.7</b>	73.9	89.0	124	<b>82.7</b>	73.0	89.5	87
55+	<b>75.1</b>	71.9	78.1	956	<b>74.8</b>	66.7	81.5	147	<b>78.0</b>	69.8	84.5	147	<b>75.7</b>	66.8	82.9	123
Female																
18-34	<b>48.6</b>	42.9	54.4	473	<b>60.2</b>	45.5	73.4	54	<b>50.0</b>	36.6	63.4	56	<b>57.9</b>	43.5	71.1	73
35-54	<b>56.6</b>	53.0	60.1	991	<b>51.9</b>	43.4	60.2	154	<b>61.6</b>	52.1	70.4	125	<b>63.7</b>	54.5	71.9	132
55+	<b>62.0</b>	59.0	64.9	1,410	<b>64.1</b>	56.6	70.9	196	<b>61.1</b>	53.9	67.8	220	<b>63.1</b>	56.2	69.6	240
<b>INCOME</b>																
Less than \$15,000	<b>60.7</b>	53.1	67.8	448	<b>63.9</b>	48.7	76.8	68	<b>62.1</b>	46.9	75.3	83	<b>67.0</b>	51.7	79.3	77
\$15,000 - \$24,999	<b>59.8</b>	54.0	65.4	790	<b>67.4</b>	54.9	77.7	115	<b>78.5</b>	68.2	86.1	131	<b>60.5</b>	48.5	71.3	138
\$25,000 - \$34,999	<b>65.2</b>	58.9	70.9	630	<b>63.0</b>	47.6	76.1	76	<b>65.6</b>	53.9	75.7	99	<b>74.5</b>	61.9	84.0	104
\$35,000 - \$49,999	<b>65.9</b>	61.6	69.9	814	<b>63.1</b>	51.0	73.8	108	<b>65.1</b>	53.4	75.3	121	<b>75.7</b>	66.8	82.8	126
\$50,000-\$74,999	<b>63.0</b>	58.4	67.4	766	<b>63.2</b>	52.5	72.7	109	<b>59.1</b>	47.1	70.2	109	<b>60.4</b>	48.4	71.3	101
\$75,000+	<b>62.1</b>	57.8	66.2	887	<b>62.7</b>	52.3	72.0	131	<b>60.3</b>	49.3	70.4	107	<b>67.3</b>	53.5	78.6	84
<b>EMPLOYMENT</b>																
Employed	<b>64.1</b>	61.4	66.7	2,577	<b>62.2</b>	55.5	68.5	354	<b>64.2</b>	57.9	70.0	365	<b>72.1</b>	66.1	77.5	361
Unemployed	<b>63.5</b>	52.2	73.5	172	*	*	*	*	*	*	*	*	*	*	*	*
Other**	<b>58.7</b>	55.6	61.7	2,095	<b>65.8</b>	59.3	71.7	303	<b>61.2</b>	53.6	68.3	322	<b>59.5</b>	51.5	67.0	320
<b>EDUCATION</b>																
K-11th Grade	<b>63.5</b>	55.9	70.4	378	<b>70.2</b>	48.8	85.4	51	<b>50.8</b>	33.0	68.3	52	<b>68.8</b>	53.1	81.1	88
12th Grade or GED	<b>60.7</b>	56.6	64.6	1,438	<b>62.1</b>	53.2	70.2	217	<b>64.4</b>	56.0	72.0	235	<b>67.8</b>	58.6	75.7	245
Some College	<b>65.9</b>	62.4	69.2	1,532	<b>67.4</b>	59.8	74.2	232	<b>73.0</b>	65.4	79.4	215	<b>66.1</b>	57.2	74.0	215
College Graduate+	<b>59.5</b>	56.1	62.8	1,494	<b>57.6</b>	49.3	65.6	184	<b>56.7</b>	48.3	64.6	214	<b>66.7</b>	58.0	74.3	162

\*Figure not reliable by BRFSS standards (n<50)

\*\*Other includes students, homemakers, retirees, and persons unable to work

# Overweight

## Idaho adults who were overweight (BMI ≥ 25), 2008

	Public Health District															
	District 4			District 5			District 6			District 7						
	%	95% CI		n	%	95% CI		n	%	95% CI		n	%	95% CI		n
<b>TOTAL</b>	<b>59.4</b>	54.7	64.0	720	<b>62.8</b>	58.0	67.2	679	<b>65.1</b>	59.6	70.2	674	<b>56.8</b>	51.1	62.3	664
<b>SEX</b>																
Male	<b>65.5</b>	57.7	72.5	293	<b>71.8</b>	64.2	78.4	273	<b>74.2</b>	66.1	80.9	254	<b>62.4</b>	53.1	70.9	285
Female	<b>52.8</b>	47.2	58.4	427	<b>53.2</b>	47.6	58.8	406	<b>56.0</b>	48.6	63.0	420	<b>51.1</b>	44.1	58.2	379
<b>AGE</b>																
18-24	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
25-34	<b>46.0</b>	34.8	57.7	85	<b>52.2</b>	39.2	64.9	72	<b>65.8</b>	54.7	75.5	90	<b>48.8</b>	37.3	60.5	107
35-44	<b>59.6</b>	50.6	67.9	143	<b>69.6</b>	59.3	78.2	99	<b>76.0</b>	66.1	83.7	87	<b>65.2</b>	55.3	73.9	104
45-54	<b>71.2</b>	62.7	78.5	144	<b>69.5</b>	60.4	77.2	140	<b>73.4</b>	64.9	80.5	132	<b>64.8</b>	55.6	73.1	133
55-64	<b>69.5</b>	61.3	76.6	156	<b>77.1</b>	68.9	83.7	133	<b>74.0</b>	65.9	80.8	154	<b>67.4</b>	57.8	75.8	115
65+	<b>66.5</b>	58.3	73.7	166	<b>59.4</b>	52.1	66.3	210	<b>61.2</b>	53.1	68.6	185	<b>69.5</b>	61.8	76.2	174
18-34	<b>46.4</b>	36.0	57.1	107	<b>50.2</b>	38.7	61.8	92	<b>54.2</b>	41.8	66.0	114	<b>41.1</b>	30.3	52.9	136
35-54	<b>65.1</b>	58.9	70.8	287	<b>69.5</b>	62.8	75.5	239	<b>74.7</b>	68.3	80.1	219	<b>65.0</b>	58.3	71.1	237
55+	<b>68.0</b>	62.3	73.2	322	<b>67.1</b>	61.6	72.2	343	<b>67.2</b>	61.4	72.4	339	<b>68.5</b>	62.5	74.0	289
<b>SEX and AGE</b>																
Male																
18-34	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
35-54	<b>76.0</b>	66.9	83.3	118	<b>77.8</b>	67.6	85.5	90	<b>88.7</b>	80.4	93.7	86	<b>74.8</b>	65.0	82.6	108
55+	<b>73.8</b>	65.2	80.9	136	<b>80.0</b>	72.4	86.0	148	<b>68.9</b>	59.7	76.9	127	<b>75.9</b>	67.0	83.0	128
Female																
18-34	<b>45.8</b>	33.8	58.2	69	<b>43.1</b>	30.6	56.5	59	<b>43.6</b>	28.4	60.1	74	<b>41.4</b>	27.4	56.9	88
35-54	<b>52.3</b>	44.2	60.2	169	<b>60.5</b>	51.7	68.6	149	<b>60.4</b>	51.2	69.0	133	<b>54.2</b>	45.0	63.2	129
55+	<b>62.6</b>	54.8	69.7	186	<b>55.0</b>	47.3	62.4	195	<b>65.5</b>	58.3	72.0	212	<b>61.3</b>	53.0	69.0	161
<b>INCOME</b>																
Less than \$15,000	<b>65.2</b>	44.1	81.7	50	<b>63.1</b>	47.9	76.1	60	<b>55.5</b>	34.2	74.9	63	*	*	*	*
\$15,000 - \$24,999	<b>56.6</b>	40.6	71.3	80	<b>64.4</b>	50.0	76.6	114	<b>62.7</b>	49.3	74.4	107	<b>36.9</b>	24.8	50.8	105
\$25,000 - \$34,999	<b>59.9</b>	44.4	73.8	84	<b>66.9</b>	54.4	77.4	93	<b>63.1</b>	45.1	78.2	97	<b>65.6</b>	47.2	80.3	77
\$35,000 - \$49,999	<b>65.0</b>	54.9	74.0	128	<b>57.3</b>	45.3	68.4	106	<b>65.5</b>	53.7	75.6	119	<b>68.0</b>	56.8	77.5	106
\$50,000-\$74,999	<b>59.2</b>	47.7	69.7	113	<b>64.6</b>	52.8	74.9	105	<b>73.3</b>	63.2	81.4	106	<b>66.6</b>	57.0	74.9	123
\$75,000+	<b>59.5</b>	51.1	67.3	195	<b>60.1</b>	49.6	69.8	118	<b>68.0</b>	55.7	78.2	115	<b>62.5</b>	51.9	72.1	137
<b>EMPLOYMENT</b>																
Employed	<b>59.1</b>	52.7	65.2	400	<b>65.8</b>	59.7	71.5	385	<b>71.7</b>	65.3	77.3	340	<b>59.7</b>	52.6	66.4	372
Unemployed	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Other**	<b>60.2</b>	53.2	66.8	286	<b>56.8</b>	49.6	63.8	276	<b>52.9</b>	44.2	61.5	316	<b>51.3</b>	41.5	60.9	272
<b>EDUCATION</b>																
K-11th Grade	*	*	*	*	<b>68.4</b>	52.7	80.8	61	*	*	*	*	*	*	*	*
12th Grade or GED	<b>57.2</b>	46.2	67.5	168	<b>55.0</b>	45.6	64.1	194	<b>63.6</b>	54.0	72.1	211	<b>55.2</b>	43.5	66.3	168
Some College	<b>69.3</b>	61.1	76.5	204	<b>65.9</b>	58.2	72.9	232	<b>65.0</b>	54.0	74.5	218	<b>55.3</b>	44.9	65.1	216
College Graduate+	<b>54.4</b>	47.6	61.1	308	<b>64.9</b>	56.9	72.2	190	<b>66.0</b>	57.8	73.4	196	<b>62.0</b>	54.0	69.4	240

\*Figure not reliable by BRFSS standards (n<50)

\*\*Other includes students, homemakers, retirees, and persons unable to work

# Obesity

## Idaho adults who were obese (BMI ≥ 30), 2008

	Statewide			Public Health District												
				District 1			District 2			District 3						
	%	95% CI		n	%	95% CI		n	%	95% CI		n	%	95% CI		n
<b>TOTAL</b>	<b>25.2</b>	23.6	26.8	4,851	<b>24.3</b>	20.7	28.3	685	<b>28.1</b>	24.3	32.2	717	<b>31.0</b>	26.5	35.9	712
<b>SEX</b>																
Male	<b>25.2</b>	22.9	27.8	1,962	<b>22.9</b>	17.7	29.1	279	<b>29.7</b>	24.1	36.1	315	<b>30.4</b>	24.0	37.8	263
Female	<b>25.1</b>	23.0	27.2	2,889	<b>25.8</b>	21.2	31.0	406	<b>26.3</b>	21.6	31.5	402	<b>31.5</b>	25.5	38.2	449
<b>AGE</b>																
18-24	<b>14.6</b>	9.4	22.0	157	*	*	*	*	*	*	*	*	*	*	*	*
25-34	<b>22.5</b>	18.6	26.9	596	<b>25.1</b>	15.0	38.9	62	<b>22.9</b>	14.6	33.9	80	<b>23.7</b>	15.8	34.0	100
35-44	<b>26.0</b>	22.5	29.9	732	<b>17.7</b>	10.9	27.5	96	<b>28.3</b>	19.9	38.5	104	<b>28.1</b>	19.2	39.1	99
45-54	<b>31.2</b>	27.8	34.8	976	<b>22.5</b>	16.4	30.0	162	<b>40.1</b>	31.9	48.9	145	<b>43.7</b>	34.4	53.4	120
55-64	<b>32.9</b>	29.6	36.3	1,017	<b>36.3</b>	28.7	44.7	156	<b>38.4</b>	30.3	47.1	156	<b>37.7</b>	29.2	47.0	147
65+	<b>22.1</b>	19.7	24.8	1,349	<b>27.7</b>	21.0	35.6	187	<b>22.3</b>	16.9	28.9	211	<b>20.1</b>	14.9	26.5	216
18-34	<b>19.6</b>	16.4	23.3	753	<b>20.1</b>	12.4	31.0	82	<b>20.6</b>	13.4	30.3	100	<b>29.2</b>	19.8	40.7	122
35-54	<b>28.6</b>	26.2	31.2	1,708	<b>20.3</b>	15.5	26.1	258	<b>34.7</b>	28.6	41.3	249	<b>35.3</b>	28.6	42.7	219
55+	<b>27.2</b>	25.1	29.3	2,366	<b>31.8</b>	26.6	37.5	343	<b>29.2</b>	24.3	34.5	367	<b>27.9</b>	23.0	33.5	363
<b>SEX and AGE</b>																
Male																
18-34	<b>18.3</b>	13.8	23.9	280	*	*	*	*	*	*	*	*	*	*	*	*
35-54	<b>29.9</b>	26.1	33.9	717	<b>18.5</b>	12.0	27.4	104	<b>36.5</b>	28.0	46.0	124	<b>37.8</b>	27.5	49.3	87
55+	<b>27.3</b>	24.2	30.6	956	<b>33.0</b>	25.3	41.6	147	<b>30.3</b>	22.9	38.8	147	<b>21.8</b>	14.9	30.7	123
Female																
18-34	<b>21.0</b>	16.7	26.2	473	<b>24.2</b>	14.1	38.3	54	<b>17.7</b>	9.6	30.1	56	<b>29.0</b>	16.2	46.2	73
35-54	<b>27.3</b>	24.2	30.6	991	<b>22.2</b>	15.8	30.2	154	<b>32.6</b>	24.4	42.1	125	<b>32.7</b>	24.7	41.8	132
55+	<b>27.1</b>	24.4	29.9	1,410	<b>30.7</b>	24.0	38.4	196	<b>28.1</b>	22.1	34.9	220	<b>33.6</b>	27.2	40.7	240
<b>INCOME</b>																
Less than \$15,000	<b>31.6</b>	25.5	38.3	448	<b>31.4</b>	20.3	45.3	68	<b>34.2</b>	22.0	48.9	83	<b>32.1</b>	18.5	49.6	77
\$15,000 - \$24,999	<b>23.3</b>	19.7	27.4	790	<b>28.0</b>	18.7	39.7	115	<b>38.5</b>	28.5	49.6	131	<b>25.4</b>	17.6	35.2	138
\$25,000 - \$34,999	<b>32.4</b>	27.1	38.1	630	<b>32.6</b>	20.4	47.7	76	<b>32.8</b>	23.6	43.7	99	<b>46.0</b>	33.1	59.6	104
\$35,000 - \$49,999	<b>25.4</b>	21.9	29.3	814	<b>20.8</b>	13.7	30.1	108	<b>17.2</b>	11.1	25.7	121	<b>27.6</b>	19.3	37.9	126
\$50,000-\$74,999	<b>24.5</b>	20.9	28.4	766	<b>25.3</b>	17.1	35.7	109	<b>27.2</b>	18.3	38.6	109	<b>29.4</b>	19.6	41.7	101
\$75,000+	<b>21.1</b>	17.9	24.7	887	<b>16.3</b>	10.7	24.1	131	<b>24.2</b>	16.7	33.6	107	<b>27.8</b>	18.1	40.0	84
<b>EMPLOYMENT</b>																
Employed	<b>26.1</b>	24.0	28.3	2,577	<b>22.2</b>	17.6	27.6	354	<b>29.5</b>	24.4	35.3	365	<b>32.3</b>	26.8	38.5	361
Unemployed	<b>33.2</b>	24.0	44.0	172	*	*	*	*	*	*	*	*	*	*	*	*
Other**	<b>22.4</b>	20.2	24.8	2,095	<b>26.0</b>	20.7	32.1	303	<b>25.9</b>	20.6	32.0	322	<b>24.6</b>	18.3	32.3	320
<b>EDUCATION</b>																
K-11th Grade	<b>27.7</b>	22.1	34.1	378	<b>24.4</b>	13.7	39.6	51	<b>32.4</b>	19.1	49.2	52	<b>29.9</b>	19.4	43.1	88
12th Grade or GED	<b>25.6</b>	22.6	28.9	1,438	<b>27.0</b>	20.7	34.5	217	<b>26.2</b>	20.3	33.1	235	<b>32.2</b>	23.7	41.9	245
Some College	<b>29.0</b>	26.0	32.2	1,532	<b>26.2</b>	19.8	33.8	232	<b>34.4</b>	27.1	42.4	215	<b>32.2</b>	24.3	41.1	215
College Graduate+	<b>20.1</b>	17.8	22.7	1,494	<b>18.4</b>	13.0	25.4	184	<b>22.9</b>	16.8	30.5	214	<b>28.5</b>	21.1	37.2	162

\*Figure not reliable by BRFSS standards (n<50)

\*\*Other includes students, homemakers, retirees, and persons unable to work

# Obesity

## Idaho adults who were obese (BMI ≥ 30), 2008

	Public Health District															
	District 4			District 5			District 6			District 7						
	%	95% CI		n	%	95% CI		n	%	95% CI		n	%	95% CI		n
<b>TOTAL</b>	<b>21.1</b>	17.8	24.9	720	<b>22.8</b>	19.4	26.7	679	<b>32.9</b>	28.2	38.1	674	<b>21.4</b>	17.9	25.4	664
<b>SEX</b>																
Male	<b>21.3</b>	16.4	27.2	293	<b>22.0</b>	16.9	28.1	273	<b>34.6</b>	27.5	42.4	254	<b>22.6</b>	17.3	29.1	285
Female	<b>20.9</b>	16.8	25.7	427	<b>23.6</b>	19.3	28.6	406	<b>31.3</b>	25.1	38.2	420	<b>20.2</b>	16.0	25.3	379
<b>AGE</b>																
18-24	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
25-34	<b>19.2</b>	11.7	29.9	85	<b>16.6</b>	9.4	27.6	72	<b>37.3</b>	26.6	49.4	90	<b>20.3</b>	13.0	30.2	107
35-44	<b>23.1</b>	16.6	31.3	143	<b>32.3</b>	23.1	43.0	99	<b>33.1</b>	23.2	44.7	87	<b>28.2</b>	20.0	38.1	104
45-54	<b>27.5</b>	20.2	36.3	144	<b>26.1</b>	18.7	35.1	140	<b>44.5</b>	35.4	54.1	132	<b>25.5</b>	18.2	34.5	133
55-64	<b>26.2</b>	19.3	34.6	156	<b>32.8</b>	24.8	41.9	133	<b>37.8</b>	29.9	46.5	154	<b>28.6</b>	20.5	38.5	115
65+	<b>19.4</b>	13.8	26.7	166	<b>18.8</b>	13.8	25.0	210	<b>22.6</b>	16.7	29.9	185	<b>25.4</b>	19.1	33.1	174
18-34	<b>14.9</b>	9.2	23.3	107	<b>13.2</b>	7.6	21.9	92	<b>30.1</b>	20.2	42.2	114	<b>12.8</b>	7.7	20.5	136
35-54	<b>25.2</b>	20.1	31.1	287	<b>29.1</b>	23.1	35.9	239	<b>38.9</b>	31.9	46.4	219	<b>26.8</b>	21.1	33.3	237
55+	<b>22.9</b>	18.2	28.3	322	<b>24.9</b>	20.3	30.2	343	<b>29.7</b>	24.7	35.3	339	<b>27.0</b>	21.7	33.0	289
<b>SEX and AGE</b>																
Male																
18-34	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
35-54	<b>28.3</b>	20.6	37.4	118	<b>26.8</b>	18.3	37.5	90	<b>41.1</b>	30.5	52.5	86	<b>28.1</b>	20.2	37.7	108
55+	<b>23.6</b>	16.9	31.9	136	<b>27.4</b>	20.4	35.7	148	<b>31.8</b>	23.8	41.0	127	<b>27.0</b>	19.2	36.4	128
Female																
18-34	<b>19.3</b>	11.4	30.7	69	<b>15.6</b>	8.5	27.1	59	<b>29.4</b>	16.6	46.5	74	<b>11.3</b>	6.0	20.0	88
35-54	<b>21.6</b>	15.7	29.0	169	<b>31.6</b>	23.9	40.4	149	<b>36.7</b>	28.0	46.4	133	<b>25.3</b>	17.9	34.4	129
55+	<b>22.2</b>	16.1	29.7	186	<b>22.5</b>	16.8	29.4	195	<b>27.7</b>	21.9	34.4	212	<b>27.0</b>	20.2	35.1	161
<b>INCOME</b>																
Less than \$15,000	<b>34.6</b>	20.5	52.0	50	<b>37.0</b>	23.8	52.5	60	<b>39.0</b>	20.0	62.0	63	*	*	*	*
\$15,000 - \$24,999	<b>14.2</b>	7.7	24.8	80	<b>22.3</b>	14.1	33.5	114	<b>31.0</b>	21.2	42.9	107	<b>16.5</b>	10.1	25.8	105
\$25,000 - \$34,999	<b>28.9</b>	17.9	43.2	84	<b>26.1</b>	17.6	37.0	93	<b>30.5</b>	19.3	44.6	97	<b>26.3</b>	14.1	43.8	77
\$35,000 - \$49,999	<b>25.2</b>	17.5	34.9	128	<b>21.0</b>	13.6	31.0	106	<b>41.4</b>	30.7	52.8	119	<b>22.6</b>	14.9	32.8	106
\$50,000-\$74,999	<b>21.5</b>	14.1	31.3	113	<b>19.5</b>	12.6	29.0	105	<b>33.4</b>	24.3	44.0	106	<b>20.1</b>	13.6	28.8	123
\$75,000+	<b>17.2</b>	11.9	24.4	195	<b>22.4</b>	14.7	32.7	118	<b>30.6</b>	19.8	44.0	115	<b>22.8</b>	15.9	31.5	137
<b>EMPLOYMENT</b>																
Employed	<b>22.3</b>	17.9	27.4	400	<b>23.7</b>	19.2	28.9	385	<b>35.7</b>	29.4	42.6	340	<b>24.2</b>	19.4	29.8	372
Unemployed	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Other**	<b>20.3</b>	15.5	26.1	286	<b>19.8</b>	15.1	25.5	276	<b>25.6</b>	20.0	32.1	316	<b>16.1</b>	11.7	21.6	272
<b>EDUCATION</b>																
K-11th Grade	*	*	*	*	<b>16.8</b>	8.4	30.7	61	*	*	*	*	*	*	*	*
12th Grade or GED	<b>23.0</b>	16.2	31.6	168	<b>20.8</b>	14.9	28.1	194	<b>30.7</b>	23.5	39.0	211	<b>18.3</b>	12.5	26.0	168
Some College	<b>31.4</b>	24.0	39.8	204	<b>25.6</b>	19.7	32.5	232	<b>34.5</b>	26.6	43.4	218	<b>20.1</b>	14.6	27.1	216
College Graduate+	<b>12.7</b>	9.2	17.3	308	<b>23.8</b>	17.5	31.5	190	<b>27.9</b>	21.0	36.1	196	<b>26.0</b>	19.7	33.6	240

\*Figure not reliable by BRFSS standards (n<50)

\*\*Other includes students, homemakers, retirees, and persons unable to work

# Cigarette Smoking

Idaho adults who smoked cigarettes, 2008

	Statewide			Public Health District												
				District 1			District 2			District 3						
	%	95% CI		n	%	95% CI		n	%	95% CI		n	%	95% CI		n
<b>TOTAL</b>	<b>16.9</b>	15.4	18.5	5,095	<b>20.5</b>	16.9	24.7	729	<b>18.3</b>	15.0	22.0	748	<b>17.7</b>	14.3	21.7	748
<b>SEX</b>																
Male	<b>18.3</b>	15.9	21.0	1,979	<b>19.9</b>	14.3	26.9	283	<b>16.4</b>	12.1	22.0	317	<b>21.6</b>	16.0	28.4	267
Female	<b>15.4</b>	13.8	17.2	3,116	<b>21.2</b>	16.9	26.3	446	<b>20.1</b>	15.7	25.5	431	<b>13.9</b>	10.5	18.3	481
<b>AGE</b>																
18-24	<b>17.2</b>	11.4	25.1	164	*	*	*	*	*	*	*	*	*	*	*	*
25-34	<b>22.3</b>	18.2	26.9	628	<b>26.0</b>	15.9	39.4	67	<b>26.2</b>	17.5	37.3	85	<b>21.3</b>	14.0	31.0	108
35-44	<b>17.8</b>	14.7	21.3	773	<b>22.5</b>	15.0	32.5	104	<b>22.1</b>	14.9	31.4	111	<b>19.2</b>	12.0	29.3	103
45-54	<b>17.3</b>	14.8	20.1	1,034	<b>24.7</b>	18.4	32.3	169	<b>17.8</b>	12.1	25.5	153	<b>19.4</b>	13.1	27.8	130
55-64	<b>16.0</b>	13.6	18.7	1,067	<b>17.5</b>	12.0	24.8	161	<b>13.8</b>	9.0	20.5	164	<b>21.2</b>	14.5	30.0	152
65+	<b>8.9</b>	7.3	10.8	1,388	<b>7.2</b>	4.3	11.9	199	<b>13.6</b>	9.2	19.6	211	<b>10.9</b>	6.9	16.8	223
18-34	<b>20.5</b>	17.0	24.4	792	<b>27.4</b>	18.0	39.4	89	<b>21.5</b>	14.5	30.8	106	<b>18.4</b>	11.7	27.7	130
35-54	<b>17.5</b>	15.5	19.7	1,807	<b>23.7</b>	18.7	29.6	273	<b>19.7</b>	15.0	25.5	264	<b>19.3</b>	14.3	25.7	233
55+	<b>12.3</b>	10.8	13.8	2,455	<b>12.1</b>	8.9	16.2	360	<b>13.7</b>	10.3	18.0	375	<b>15.5</b>	11.6	20.4	375
<b>SEX and AGE</b>																
Male																
18-34	<b>22.2</b>	16.7	28.9	290	*	*	*	*	*	*	*	*	<b>22.8</b>	12.3	38.5	52
35-54	<b>18.4</b>	15.3	22.0	724	<b>22.3</b>	14.8	32.1	104	<b>19.2</b>	12.8	28.0	126	<b>20.4</b>	12.7	31.2	87
55+	<b>13.8</b>	11.5	16.5	956	<b>12.0</b>	7.5	18.7	148	<b>13.8</b>	9.0	20.6	146	<b>22.2</b>	15.2	31.2	124
Female																
18-34	<b>18.6</b>	14.8	23.1	502	<b>28.6</b>	17.5	43.2	59	<b>27.5</b>	17.1	41.0	61	<b>13.7</b>	7.3	24.4	78
35-54	<b>16.6</b>	14.3	19.2	1,083	<b>25.1</b>	18.9	32.6	169	<b>20.2</b>	14.0	28.4	138	<b>18.3</b>	12.5	25.9	146
55+	<b>10.9</b>	9.2	12.8	1,499	<b>12.2</b>	8.1	17.8	212	<b>13.5</b>	9.2	19.4	229	<b>9.5</b>	6.1	14.7	251
<b>INCOME</b>																
Less than \$15,000	<b>27.6</b>	22.2	33.7	481	<b>31.8</b>	19.8	46.9	76	<b>34.0</b>	22.1	48.4	85	<b>26.6</b>	13.5	45.8	79
\$15,000 - \$24,999	<b>27.9</b>	23.3	33.1	828	<b>32.1</b>	22.1	44.2	120	<b>23.7</b>	15.6	34.2	133	<b>31.2</b>	21.5	42.9	149
\$25,000 - \$34,999	<b>22.1</b>	17.0	28.2	659	<b>19.3</b>	9.4	35.7	78	<b>17.9</b>	10.2	29.7	104	<b>18.1</b>	11.2	27.8	109
\$35,000 - \$49,999	<b>15.8</b>	13.0	19.2	851	<b>19.1</b>	11.8	29.4	116	<b>17.4</b>	10.6	27.3	127	<b>15.8</b>	9.9	24.2	130
\$50,000-\$74,999	<b>11.2</b>	8.6	14.5	793	<b>19.1</b>	12.1	28.9	114	<b>8.4</b>	4.4	15.4	110	<b>11.0</b>	5.8	20.1	106
\$75,000+	<b>8.2</b>	6.1	10.9	910	<b>7.5</b>	3.7	14.6	135	<b>10.4</b>	5.3	19.1	111	<b>8.5</b>	4.0	17.1	85
<b>EMPLOYMENT</b>																
Employed	<b>17.8</b>	15.8	20.0	2,720	<b>21.8</b>	17.0	27.7	378	<b>17.4</b>	13.1	22.6	387	<b>17.6</b>	13.5	22.6	381
Unemployed	<b>34.3</b>	24.7	45.4	174	*	*	*	*	*	*	*	*	*	*	*	*
Other**	<b>12.7</b>	10.9	14.7	2,193	<b>16.5</b>	11.7	22.6	321	<b>16.9</b>	12.6	22.2	330	<b>12.9</b>	9.3	17.6	335
<b>EDUCATION</b>																
K-11th Grade	<b>34.0</b>	27.5	41.2	405	<b>47.1</b>	29.2	65.8	54	<b>18.5</b>	9.6	32.7	53	<b>19.4</b>	11.7	30.5	93
12th Grade or GED	<b>23.2</b>	20.1	26.6	1,514	<b>27.8</b>	21.1	35.8	232	<b>26.6</b>	20.2	34.2	251	<b>23.4</b>	16.6	31.9	255
Some College	<b>15.4</b>	13.2	17.7	1,621	<b>18.6</b>	13.5	24.9	245	<b>19.3</b>	13.6	26.6	225	<b>17.5</b>	11.8	25.1	228
College Graduate+	<b>6.7</b>	5.0	8.9	1,544	<b>4.3</b>	2.3	7.8	195	<b>7.1</b>	3.9	12.7	218	<b>8.6</b>	4.9	14.6	170

\*Figure not reliable by BRFSS standards (n<50)

\*\*Other includes students, homemakers, retirees, and persons unable to work.

Note: Cigarette smoking is defined as smoking at least 100 cigarettes in entire life and currently smoking every day or some days.

# Cigarette Smoking

Idaho adults who smoked cigarettes, 2008

	Public Health District															
	District 4				District 5				District 6				District 7			
	%	95% CI		n	%	95% CI		n	%	95% CI		n	%	95% CI		n
<b>TOTAL</b>	<b>18.3</b>	14.8	22.3	749	<b>16.1</b>	12.6	20.3	725	<b>12.0</b>	9.3	15.4	701	<b>12.2</b>	9.1	16.1	695
<b>SEX</b>																
Male	<b>21.5</b>	15.9	28.5	292	<b>18.7</b>	12.9	26.4	281	<b>10.5</b>	6.8	15.9	253	<b>12.4</b>	7.6	19.5	286
Female	<b>15.0</b>	11.4	19.3	457	<b>13.5</b>	10.3	17.6	444	<b>13.5</b>	9.9	18.1	448	<b>12.1</b>	8.7	16.5	409
<b>AGE</b>																
18-24	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
25-34	<b>24.2</b>	15.4	35.9	87	<b>25.2</b>	14.9	39.4	78	<b>14.1</b>	6.9	26.5	93	<b>13.2</b>	7.7	21.8	110
35-44	<b>18.0</b>	12.0	26.1	149	<b>16.2</b>	9.6	26.2	107	<b>12.6</b>	7.2	21.1	88	<b>12.5</b>	7.0	21.3	111
45-54	<b>14.4</b>	9.5	21.3	149	<b>17.2</b>	11.6	24.6	152	<b>16.3</b>	10.6	24.2	141	<b>12.1</b>	7.8	18.5	140
55-64	<b>15.4</b>	10.6	21.9	164	<b>12.1</b>	7.6	18.8	141	<b>19.0</b>	13.2	26.5	163	<b>11.1</b>	6.5	18.2	122
65+	<b>10.9</b>	7.0	16.8	170	<b>9.0</b>	5.7	13.8	218	<b>4.5</b>	2.4	8.2	188	<b>4.9</b>	2.6	9.1	179
18-34	<b>24.5</b>	16.4	34.9	110	<b>21.9</b>	13.4	33.7	100	<b>10.2</b>	5.2	18.9	117	<b>15.2</b>	8.8	25.0	140
35-54	<b>16.3</b>	12.2	21.5	298	<b>16.7</b>	12.2	22.5	259	<b>14.6</b>	10.4	20.1	229	<b>12.3</b>	8.6	17.3	251
55+	<b>13.2</b>	9.9	17.4	334	<b>10.4</b>	7.5	14.1	359	<b>11.4</b>	8.2	15.5	351	<b>7.9</b>	5.2	11.8	301
<b>SEX and AGE</b>																
Male																
18-34	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
35-54	<b>21.3</b>	14.6	30.1	117	<b>16.6</b>	9.9	26.4	94	<b>10.9</b>	6.0	19.1	88	<b>10.8</b>	5.9	19.0	108
55+	<b>14.6</b>	9.5	21.8	136	<b>12.2</b>	7.7	18.8	149	<b>10.5</b>	6.1	17.4	125	<b>8.6</b>	4.7	15.3	128
Female																
18-34	<b>21.9</b>	13.4	33.6	72	<b>15.4</b>	8.6	26.1	64	<b>10.3</b>	4.3	22.4	77	<b>13.9</b>	7.4	24.5	91
35-54	<b>10.9</b>	7.1	16.4	181	<b>16.8</b>	11.5	24.1	165	<b>18.1</b>	12.1	26.3	141	<b>13.8</b>	9.0	20.8	143
55+	<b>12.0</b>	8.2	17.3	198	<b>8.7</b>	5.5	13.5	210	<b>12.1</b>	8.2	17.6	226	<b>7.3</b>	4.2	12.5	173
<b>INCOME</b>																
Less than \$15,000	<b>36.9</b>	22.5	54.0	55	<b>24.7</b>	14.6	38.5	67	<b>15.4</b>	7.7	28.4	67	<b>20.0</b>	10.6	34.7	52
\$15,000 - \$24,999	<b>33.0</b>	20.6	48.3	81	<b>28.8</b>	16.3	45.6	123	<b>16.6</b>	10.8	24.8	114	<b>18.9</b>	10.7	31.2	108
\$25,000 - \$34,999	<b>30.9</b>	17.8	48.1	86	<b>16.0</b>	9.2	26.4	101	<b>15.8</b>	8.0	28.9	101	<b>28.2</b>	14.0	48.7	80
\$35,000 - \$49,999	<b>16.5</b>	10.6	24.7	133	<b>18.4</b>	10.8	29.6	112	<b>13.4</b>	6.8	24.7	121	<b>7.6</b>	3.9	14.2	112
\$50,000-\$74,999	<b>14.9</b>	8.7	24.5	116	<b>8.6</b>	3.9	18.0	109	<b>6.4</b>	3.2	12.4	110	<b>2.4</b>	0.8	6.6	128
\$75,000+	<b>8.5</b>	4.6	15.1	201	<b>7.7</b>	3.5	16.1	119	<b>9.7</b>	4.9	18.3	118	<b>5.7</b>	2.7	11.7	141
<b>EMPLOYMENT</b>																
Employed	<b>18.8</b>	14.3	24.4	417	<b>18.7</b>	13.9	24.7	410	<b>13.0</b>	9.2	18.2	357	<b>14.2</b>	10.4	19.2	390
Unemployed	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Other**	<b>12.7</b>	8.9	17.9	299	<b>11.2</b>	7.7	16.1	296	<b>9.6</b>	6.7	13.5	326	<b>9.0</b>	4.4	17.3	286
<b>EDUCATION</b>																
K-11th Grade	*	*	*	*	<b>30.2</b>	15.7	50.1	70	<b>24.2</b>	11.8	43.3	51	*	*	*	*
12th Grade or GED	<b>26.6</b>	18.4	36.7	173	<b>21.4</b>	14.9	29.7	207	<b>12.3</b>	8.1	18.1	218	<b>17.8</b>	11.6	26.4	178
Some College	<b>15.5</b>	10.9	21.5	218	<b>13.7</b>	9.4	19.5	247	<b>13.6</b>	8.8	20.3	229	<b>10.0</b>	6.0	16.3	229
College Graduate+	<b>8.8</b>	5.2	14.5	317	<b>6.2</b>	3.5	10.7	199	<b>4.9</b>	2.3	10.0	201	<b>3.1</b>	1.6	6.1	244

\*Figure not reliable by BRFSS standards (n<50)

\*\*Other includes students, homemakers, retirees, and persons unable to work.

Note: Cigarette smoking is defined as smoking at least 100 cigarettes in entire life and currently smoking every day or some days.

# Binge Drinking

## Idaho adults who were binge drinkers, 2008

	Statewide			Public Health District												
				District 1			District 2			District 3						
	%	95% CI		n	%	95% CI		n	%	95% CI		n	%	95% CI		n
<b>TOTAL</b>	<b>13.1</b>	11.8	14.6	5,028	<b>15.8</b>	12.2	20.1	713	<b>17.5</b>	14.1	21.5	741	<b>11.2</b>	8.4	14.7	735
<b>SEX</b>																
Male	<b>17.0</b>	14.7	19.6	1,949	<b>21.7</b>	15.6	29.5	275	<b>21.3</b>	16.0	27.6	313	<b>14.8</b>	10.2	21.0	263
Female	<b>9.3</b>	7.9	10.9	3,079	<b>10.0</b>	7.0	14.1	438	<b>13.6</b>	9.6	19.0	428	<b>7.6</b>	5.1	11.3	472
<b>AGE</b>																
18-24	<b>18.0</b>	12.1	25.9	160	*	*	*	*	*	*	*	*	*	*	*	*
25-34	<b>18.8</b>	15.2	23.1	622	<b>28.7</b>	17.6	43.2	67	<b>21.4</b>	13.2	32.7	84	<b>16.3</b>	9.9	25.7	105
35-44	<b>15.6</b>	12.6	19.1	767	<b>18.8</b>	11.8	28.4	104	<b>22.2</b>	15.1	31.6	109	<b>14.3</b>	8.2	23.9	102
45-54	<b>14.1</b>	11.8	16.8	1,020	<b>17.2</b>	11.9	24.3	165	<b>19.9</b>	14.0	27.6	151	<b>15.5</b>	9.9	23.4	130
55-64	<b>8.5</b>	6.7	10.7	1,052	<b>7.2</b>	4.0	12.4	158	<b>12.0</b>	7.6	18.5	163	<b>9.4</b>	5.3	16.2	150
65+	<b>2.2</b>	1.5	3.3	1,370	<b>1.1</b>	0.4	3.3	192	<b>3.4</b>	1.8	6.4	214	<b>0.4</b>	0.1	2.9	218
18-34	<b>18.5</b>	15.3	22.3	782	<b>27.7</b>	17.9	40.3	88	<b>24.7</b>	16.7	35.0	102	<b>13.2</b>	7.7	21.6	127
35-54	<b>14.8</b>	12.9	17.0	1,787	<b>17.9</b>	13.4	23.6	269	<b>21.0</b>	16.2	26.7	260	<b>14.9</b>	10.4	20.8	232
55+	<b>5.2</b>	4.2	6.4	2,422	<b>4.0</b>	2.4	6.7	350	<b>7.1</b>	4.8	10.3	377	<b>4.4</b>	2.5	7.7	368
<b>SEX and AGE</b>																
Male																
18-34	<b>21.8</b>	16.6	28.2	285	*	*	*	*	*	*	*	*	<b>17.7</b>	8.8	32.2	51
35-54	<b>19.6</b>	16.3	23.5	714	<b>23.3</b>	15.7	33.1	102	<b>25.9</b>	18.5	34.9	124	<b>17.9</b>	10.8	28.2	88
55+	<b>8.0</b>	6.2	10.2	941	<b>5.9</b>	3.1	10.9	142	<b>12.2</b>	8.0	18.3	146	<b>7.5</b>	3.8	14.0	121
Female																
18-34	<b>15.0</b>	11.6	19.3	497	<b>16.9</b>	8.9	29.9	58	<b>24.0</b>	13.9	38.2	59	<b>8.5</b>	3.8	18.1	76
35-54	<b>10.0</b>	8.2	12.1	1,073	<b>12.8</b>	8.4	18.9	167	<b>16.1</b>	10.7	23.5	136	<b>11.9</b>	7.4	18.7	144
55+	<b>2.7</b>	1.9	3.8	1,481	<b>2.3</b>	0.9	5.7	208	<b>2.3</b>	1.0	5.5	231	<b>1.8</b>	0.7	4.3	247
<b>INCOME</b>																
Less than \$15,000	<b>9.7</b>	6.4	14.4	476	<b>8.0</b>	3.5	17.5	75	<b>17.6</b>	7.7	35.3	83	<b>9.6</b>	3.7	22.4	80
\$15,000 - \$24,999	<b>15.4</b>	11.4	20.5	816	<b>9.3</b>	4.4	18.7	118	<b>13.6</b>	7.8	22.6	131	<b>12.4</b>	5.8	24.5	144
\$25,000 - \$34,999	<b>11.3</b>	8.1	15.7	649	<b>18.4</b>	8.4	35.4	76	<b>11.8</b>	6.8	19.6	105	<b>8.8</b>	3.9	18.5	105
\$35,000 - \$49,999	<b>13.0</b>	10.2	16.4	846	<b>16.3</b>	8.4	29.2	116	<b>24.1</b>	15.3	35.9	126	<b>11.2</b>	6.4	18.9	130
\$50,000-\$74,999	<b>14.0</b>	10.8	17.8	783	<b>22.5</b>	13.9	34.3	112	<b>15.8</b>	8.8	26.9	108	<b>11.9</b>	6.0	22.1	104
\$75,000+	<b>15.5</b>	12.5	19.2	904	<b>18.5</b>	11.3	28.9	133	<b>19.4</b>	12.7	28.4	110	<b>17.3</b>	9.8	28.6	85
<b>EMPLOYMENT</b>																
Employed	<b>16.8</b>	14.9	19.0	2,680	<b>22.1</b>	16.7	28.7	369	<b>24.8</b>	19.6	31.0	378	<b>13.4</b>	9.8	18.0	376
Unemployed	<b>19.1</b>	11.2	30.5	174	*	*	*	*	*	*	*	*	*	*	*	*
Other**	<b>5.6</b>	4.3	7.3	2,167	<b>5.5</b>	3.0	10.1	314	<b>6.3</b>	3.8	10.4	332	<b>4.4</b>	2.6	7.4	327
<b>EDUCATION</b>																
K-11th Grade	<b>16.8</b>	11.5	24.0	395	<b>20.0</b>	6.7	46.7	50	<b>12.3</b>	4.8	28.2	53	<b>6.9</b>	2.6	17.4	92
12th Grade or GED	<b>15.9</b>	13.1	19.0	1,492	<b>20.5</b>	14.1	28.9	226	<b>22.3</b>	16.0	30.2	249	<b>12.9</b>	7.5	21.3	252
Some College	<b>11.7</b>	9.6	14.2	1,602	<b>13.8</b>	8.6	21.4	242	<b>18.0</b>	12.4	25.5	223	<b>12.7</b>	8.4	18.7	222
College Graduate+	<b>10.9</b>	8.9	13.3	1,530	<b>10.9</b>	6.6	17.7	192	<b>13.0</b>	7.8	20.8	215	<b>9.7</b>	5.4	16.8	169

\*Figure not reliable by BRFSS standards (n<50)

\*\*Other includes students, homemakers, retirees, and persons unable to work.

Note: Definition of binge drinking is males who consumed 5+ drinks and females who consumed 4+ drinks on an occasion in the last 30 days. This definition changed beginning in 2006.

# Binge Drinking

## Idaho adults who were binge drinkers , 2008

	Public Health District															
	District 4			District 5			District 6			District 7						
	%	95% CI		n	%	95% CI		n	%	95% CI		n	%	95% CI		n
<b>TOTAL</b>	<b>15.0</b>	11.8	18.8	738	<b>11.9</b>	8.6	16.2	715	<b>8.6</b>	6.2	11.9	698	<b>10.8</b>	7.7	15.0	688
<b>SEX</b>																
Male	<b>19.1</b>	13.9	25.6	286	<b>16.7</b>	11.0	24.6	278	<b>10.6</b>	6.7	16.4	252	<b>12.6</b>	7.7	20.1	282
Female	<b>10.8</b>	7.6	15.1	452	<b>7.1</b>	4.8	10.4	437	<b>6.6</b>	4.1	10.6	446	<b>9.0</b>	5.8	13.8	406
<b>AGE</b>																
18-24	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
25-34	<b>17.3</b>	10.7	26.6	87	<b>18.6</b>	9.5	33.1	78	<b>13.4</b>	7.1	24.0	92	<b>17.6</b>	9.6	30.0	109
35-44	<b>17.2</b>	10.9	26.1	147	<b>13.5</b>	8.2	21.5	106	<b>11.4</b>	5.6	21.8	88	<b>10.6</b>	5.8	18.8	111
45-54	<b>14.2</b>	9.0	21.6	144	<b>14.4</b>	9.0	22.1	150	<b>8.4</b>	4.6	14.8	140	<b>9.6</b>	5.5	16.2	140
55-64	<b>10.1</b>	6.0	16.5	162	<b>3.7</b>	1.6	8.3	139	<b>8.9</b>	5.0	15.4	160	<b>7.8</b>	3.6	16.1	120
65+	<b>4.1</b>	1.9	8.4	168	<b>3.0</b>	1.4	6.5	213	<b>2.0</b>	0.7	5.7	189	<b>1.0</b>	0.3	3.0	176
18-34	<b>20.4</b>	13.4	29.7	110	<b>18.6</b>	10.4	31.1	100	<b>10.4</b>	5.5	18.7	116	<b>16.2</b>	9.4	26.6	139
35-54	<b>15.8</b>	11.4	21.4	291	<b>14.0</b>	10.0	19.2	256	<b>9.8</b>	6.1	15.4	228	<b>10.1</b>	6.7	15.0	251
55+	<b>7.1</b>	4.6	10.9	330	<b>3.3</b>	1.8	5.8	352	<b>5.2</b>	3.1	8.6	349	<b>4.3</b>	2.1	8.5	296
<b>SEX and AGE</b>																
Male																
18-34	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
35-54	<b>24.5</b>	16.9	34.2	113	<b>15.6</b>	9.5	24.4	92	<b>11.7</b>	5.9	21.8	87	<b>11.9</b>	6.8	20.0	108
55+	<b>10.7</b>	6.2	17.9	134	<b>6.0</b>	3.2	11.0	148	<b>6.4</b>	3.3	12.2	124	<b>6.5</b>	2.8	14.6	126
Female																
18-34	<b>21.9</b>	13.4	33.6	72	<b>7.9</b>	3.5	16.6	64	<b>7.7</b>	2.8	19.2	76	<b>14.6</b>	7.6	26.3	92
35-54	<b>6.6</b>	3.8	11.1	178	<b>12.4</b>	7.9	19.0	164	<b>8.0</b>	4.3	14.3	141	<b>8.2</b>	4.4	14.8	143
55+	<b>4.0</b>	2.0	7.7	196	<b>0.8</b>	0.2	3.5	204	<b>4.2</b>	1.9	9.1	225	<b>2.3</b>	0.7	6.9	170
<b>INCOME</b>																
Less than \$15,000	<b>10.7</b>	3.4	28.7	53	<b>4.1</b>	1.1	14.2	65	<b>13.4</b>	4.5	34.1	68	<b>5.9</b>	1.8	17.7	52
\$15,000 - \$24,999	<b>21.5</b>	11.1	37.6	80	<b>23.5</b>	11.0	43.3	122	<b>9.0</b>	4.4	17.5	114	<b>15.4</b>	7.2	30.1	107
\$25,000 - \$34,999	<b>8.2</b>	3.4	18.1	88	<b>6.2</b>	2.8	13.1	100	<b>7.7</b>	3.6	15.9	99	<b>27.8</b>	12.6	50.5	76
\$35,000 - \$49,999	<b>13.0</b>	7.6	21.2	132	<b>11.8</b>	6.1	21.6	110	<b>9.3</b>	3.8	20.7	120	<b>8.5</b>	4.3	16.0	112
\$50,000-\$74,999	<b>17.9</b>	10.6	28.6	113	<b>12.8</b>	6.7	23.1	109	<b>7.0</b>	3.1	15.2	109	<b>3.6</b>	1.4	8.6	128
\$75,000+	<b>17.8</b>	11.9	25.8	198	<b>9.8</b>	5.2	17.6	119	<b>10.6</b>	5.1	20.4	118	<b>10.7</b>	6.3	17.6	141
<b>EMPLOYMENT</b>																
Employed	<b>17.1</b>	12.9	22.3	409	<b>17.1</b>	12.3	23.3	406	<b>11.9</b>	8.2	17.0	356	<b>13.9</b>	9.9	19.3	386
Unemployed	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Other**	<b>8.1</b>	5.0	13.0	296	<b>2.1</b>	1.0	4.7	291	<b>3.6</b>	1.9	6.6	324	<b>6.3</b>	2.3	16.0	283
<b>EDUCATION</b>																
K-11th Grade	*	*	*	*	<b>13.9</b>	3.6	40.7	69	*	*	*	*	*	*	*	*
12th Grade or GED	<b>16.4</b>	9.9	25.8	172	<b>14.6</b>	8.4	24.2	203	<b>9.0</b>	5.2	15.2	217	<b>14.8</b>	9.2	23.1	173
Some College	<b>13.6</b>	8.5	21.1	216	<b>9.7</b>	5.9	15.4	242	<b>6.1</b>	2.9	12.6	228	<b>7.9</b>	4.2	14.3	229
College Graduate+	<b>12.9</b>	9.0	18.3	310	<b>10.6</b>	6.6	16.7	199	<b>9.9</b>	5.2	18.1	202	<b>5.8</b>	3.4	9.7	243

\*Figure not reliable by BRFSS standards (n<50)

\*\*Other includes students, homemakers, retirees, and persons unable to work.

Note: Definition of binge drinking is males who consumed 5+ drinks and females who consumed 4+ drinks on an occasion in the last 30 days. This definition changed beginning in 2006.

# Heavy Drinking

## Idaho adults who were heavy drinkers, 2008

	Statewide			Public Health District												
				District 1			District 2			District 3						
	%	95% CI		n	%	95% CI		n	%	95% CI		n	%	95% CI		n
<b>TOTAL</b>	<b>4.6</b>	3.9	5.6	5,010	<b>6.8</b>	4.9	9.4	710	<b>5.5</b>	3.8	7.9	739	<b>3.8</b>	2.5	5.8	736
<b>SEX</b>																
Male	<b>5.3</b>	4.1	6.9	1,942	<b>6.5</b>	3.7	11.1	276	<b>4.8</b>	2.6	8.8	312	<b>4.9</b>	2.8	8.7	264
Female	<b>4.0</b>	3.2	5.0	3,068	<b>7.1</b>	4.9	10.3	434	<b>6.2</b>	4.0	9.7	427	<b>2.7</b>	1.5	4.9	472
<b>AGE</b>																
18-24	<b>4.6</b>	2.0	10.1	160	*	*	*	*	*	*	*	*	*	*	*	*
25-34	<b>4.1</b>	2.4	6.9	621	<b>8.2</b>	3.2	19.6	67	<b>4.7</b>	1.9	11.4	84	<b>3.4</b>	1.1	10.0	107
35-44	<b>5.2</b>	3.6	7.6	764	<b>10.3</b>	5.7	18.0	103	<b>4.9</b>	2.0	11.4	110	<b>5.3</b>	2.1	12.7	102
45-54	<b>6.5</b>	5.0	8.6	1,017	<b>11.2</b>	6.8	18.1	164	<b>6.8</b>	3.8	12.1	153	<b>8.4</b>	4.3	15.9	129
55-64	<b>4.0</b>	2.8	5.7	1,046	<b>3.9</b>	1.8	8.2	155	<b>3.1</b>	1.3	7.6	163	<b>3.6</b>	1.4	8.9	150
65+	<b>3.1</b>	2.3	4.2	1,365	<b>3.3</b>	1.5	7.2	194	<b>5.3</b>	2.9	9.4	209	<b>1.6</b>	0.6	4.4	218
18-34	<b>4.3</b>	2.7	6.7	781	<b>5.6</b>	2.2	13.9	88	<b>6.3</b>	2.8	13.3	102	<b>2.0</b>	0.6	6.2	129
35-54	<b>5.9</b>	4.7	7.4	1,781	<b>10.8</b>	7.4	15.6	267	<b>5.9</b>	3.6	9.6	263	<b>6.8</b>	3.9	11.5	231
55+	<b>3.5</b>	2.8	4.5	2,411	<b>3.6</b>	2.1	6.2	349	<b>4.3</b>	2.6	7.0	372	<b>2.5</b>	1.2	5.0	368
<b>SEX and AGE</b>																
Male																
18-34	<b>5.0</b>	2.7	9.2	283	*	*	*	*	*	*	*	*	<b>2.6</b>	0.6	11.6	52
35-54	<b>6.4</b>	4.5	9.0	713	<b>10.9</b>	6.0	19.2	102	<b>3.7</b>	1.5	9.3	125	<b>8.7</b>	4.2	17.3	87
55+	<b>4.3</b>	3.1	6.0	937	<b>3.4</b>	1.5	7.7	143	<b>5.5</b>	2.9	10.3	144	<b>3.2</b>	1.2	8.5	122
Female																
18-34	<b>3.5</b>	1.9	6.5	498	<b>7.0</b>	2.8	16.6	58	<b>7.6</b>	3.1	17.7	59	<b>1.4</b>	0.3	5.7	77
35-54	<b>5.4</b>	4.2	7.1	1,068	<b>10.6</b>	6.6	16.7	165	<b>8.1</b>	4.5	14.2	138	<b>4.8</b>	2.2	10.5	144
55+	<b>2.8</b>	2.0	3.9	1,474	<b>3.8</b>	1.8	7.7	206	<b>3.2</b>	1.5	6.9	228	<b>1.8</b>	0.7	4.4	246
<b>INCOME</b>																
Less than \$15,000	<b>5.9</b>	3.5	9.8	472	<b>5.9</b>	2.1	15.5	73	<b>13.8</b>	5.3	31.3	84	<b>4.9</b>	1.7	13.0	80
\$15,000 - \$24,999	<b>4.9</b>	2.9	8.1	811	<b>7.9</b>	3.7	16.0	118	<b>6.0</b>	2.9	11.8	129	<b>2.1</b>	0.5	7.9	145
\$25,000 - \$34,999	<b>2.9</b>	1.8	4.7	654	<b>8.0</b>	4.0	15.5	78	<b>4.5</b>	1.9	10.0	104	<b>3.2</b>	0.9	10.5	107
\$35,000 - \$49,999	<b>4.2</b>	2.7	6.3	840	<b>4.2</b>	1.8	9.9	115	<b>5.1</b>	2.0	12.7	127	<b>5.3</b>	2.2	12.4	128
\$50,000-\$74,999	<b>4.8</b>	3.0	7.8	787	<b>6.8</b>	3.2	14.1	112	<b>3.9</b>	1.4	10.5	109	<b>3.2</b>	1.2	8.5	105
\$75,000+	<b>6.1</b>	4.4	8.5	901	<b>10.5</b>	5.2	20.0	131	<b>4.5</b>	2.0	9.8	110	<b>7.3</b>	2.8	17.5	85
<b>EMPLOYMENT</b>																
Employed	<b>5.3</b>	4.3	6.6	2,669	<b>8.5</b>	5.6	12.5	366	<b>6.9</b>	4.3	10.9	380	<b>4.7</b>	2.8	7.9	378
Unemployed	<b>8.9</b>	3.9	19.4	173	*	*	*	*	*	*	*	*	*	*	*	*
Other**	<b>2.8</b>	2.1	3.7	2,161	<b>3.3</b>	1.8	6.0	314	<b>3.3</b>	1.8	6.0	328	<b>2.5</b>	1.2	5.0	326
<b>EDUCATION</b>																
K-11th Grade	<b>3.8</b>	1.6	8.7	393	<b>5.6</b>	1.4	19.5	52	<b>4.0</b>	1.0	14.0	53	<b>0.7</b>	0.2	3.2	91
12th Grade or GED	<b>5.5</b>	4.0	7.6	1,486	<b>7.2</b>	4.4	11.8	226	<b>8.3</b>	5.1	13.1	248	<b>4.4</b>	2.2	8.6	252
Some College	<b>5.4</b>	4.0	7.2	1,595	<b>9.8</b>	5.8	16.2	238	<b>4.3</b>	1.6	10.8	222	<b>5.9</b>	3.1	11.1	225
College Graduate+	<b>3.3</b>	2.4	4.7	1,527	<b>3.1</b>	1.4	6.7	191	<b>4.0</b>	1.8	8.8	215	<b>2.2</b>	0.8	6.0	168

\*Figure not reliable by BRFSS standards (n<50)

\*\*Other includes students, homemakers, retirees, and persons unable to work

Note: Definition of heavy drinking is males who consumed >60 drinks and females who consumed >30 drinks in the last 30 days.

This definition changed beginning in 2006.

# Heavy Drinking

## Idaho adults who were heavy drinkers, 2008

	Public Health District															
	District 4			District 5			District 6			District 7						
	%	95% CI		n	%	95% CI		n	%	95% CI		n	%	95% CI		n
<b>TOTAL</b>	<b>6.2</b>	4.2	9.0	734	<b>3.3</b>	2.2	4.9	709	<b>2.6</b>	1.4	4.8	695	<b>2.1</b>	1.1	4.0	687
<b>SEX</b>																
Male	<b>8.4</b>	5.1	13.4	285	<b>3.0</b>	1.6	5.6	273	<b>2.5</b>	1.1	5.6	251	<b>2.0</b>	0.9	4.6	281
Female	<b>3.9</b>	2.2	7.1	449	<b>3.6</b>	2.2	6.0	436	<b>2.7</b>	1.1	6.6	444	<b>2.2</b>	0.8	5.6	406
<b>AGE</b>																
18-24	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
25-34	<b>5.3</b>	1.9	14.3	87	<b>0.0</b>	0.0	0.0	76	<b>2.3</b>	0.3	14.5	91	<b>1.7</b>	0.4	6.9	109
35-44	<b>5.8</b>	2.6	12.4	146	<b>4.3</b>	1.7	10.3	105	<b>2.1</b>	0.6	6.6	87	<b>1.2</b>	0.3	5.1	111
45-54	<b>6.7</b>	3.6	12.2	145	<b>5.1</b>	2.2	11.2	147	<b>1.1</b>	0.3	3.5	139	<b>3.9</b>	1.7	8.7	140
55-64	<b>4.8</b>	2.3	9.8	161	<b>4.5</b>	2.0	10.1	138	<b>4.8</b>	2.3	9.8	160	<b>2.0</b>	0.3	12.7	119
65+	<b>3.2</b>	1.5	6.6	165	<b>5.2</b>	2.7	9.6	214	<b>2.4</b>	0.9	6.0	189	<b>1.0</b>	0.2	4.3	176
18-34	<b>7.7</b>	3.6	15.7	110	<b>0.0</b>	0.0	0.0	98	<b>2.9</b>	0.7	11.1	115	<b>2.1</b>	0.6	7.6	139
35-54	<b>6.2</b>	3.8	10.1	291	<b>4.7</b>	2.6	8.5	252	<b>1.6</b>	0.7	3.7	226	<b>2.6</b>	1.3	5.4	251
55+	<b>4.1</b>	2.4	6.8	326	<b>4.9</b>	2.9	8.0	352	<b>3.5</b>	2.0	6.2	349	<b>1.5</b>	0.4	5.6	295
<b>SEX and AGE</b>																
Male																
18-34	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
35-54	<b>8.1</b>	4.1	15.4	114	<b>3.2</b>	1.0	10.3	90	<b>0.7</b>	0.1	5.0	87	<b>2.2</b>	0.7	7.0	108
55+	<b>4.8</b>	2.2	10.1	132	<b>5.8</b>	2.9	11.2	147	<b>4.9</b>	2.2	10.2	124	<b>3.2</b>	0.8	11.5	125
Female																
18-34	<b>3.8</b>	0.7	17.4	72	<b>0.0</b>	0.0	0.0	64	<b>3.4</b>	0.5	20.8	76	<b>3.0</b>	0.6	14.2	92
35-54	<b>4.2</b>	2.1	8.0	177	<b>6.1</b>	3.1	11.8	162	<b>2.4</b>	0.9	6.1	139	<b>3.0</b>	1.2	7.4	143
55+	<b>3.4</b>	1.7	6.8	194	<b>4.1</b>	1.9	8.5	205	<b>2.3</b>	1.0	5.3	225	<b>0.0</b>	0.0	0.0	170
<b>INCOME</b>																
Less than \$15,000	<b>11.2</b>	3.7	29.1	53	<b>0.8</b>	0.1	5.5	64	<b>4.5</b>	1.5	12.9	66	<b>0.0</b>	0.0	0.0	52
\$15,000 - \$24,999	<b>9.0</b>	3.0	23.8	79	<b>3.3</b>	1.1	9.6	119	<b>2.3</b>	0.8	6.4	113	<b>0.0</b>	0.0	0.0	108
\$25,000 - \$34,999	<b>0.7</b>	0.2	2.9	88	<b>0.8</b>	0.2	3.4	101	<b>1.0</b>	0.2	4.8	99	<b>6.4</b>	1.5	23.2	77
\$35,000 - \$49,999	<b>4.4</b>	1.8	10.5	130	<b>0.3</b>	0.1	2.4	109	<b>4.8</b>	1.2	18.0	119	<b>4.6</b>	1.6	12.4	112
\$50,000-\$74,999	<b>8.1</b>	3.2	18.9	114	<b>4.4</b>	1.7	11.3	109	<b>1.1</b>	0.2	4.8	110	<b>1.3</b>	0.3	5.7	128
\$75,000+	<b>5.9</b>	3.1	11.1	197	<b>8.3</b>	4.7	14.4	119	<b>3.0</b>	0.7	11.5	118	<b>2.1</b>	0.8	5.8	141
<b>EMPLOYMENT</b>																
Employed	<b>6.2</b>	3.9	9.9	406	<b>3.8</b>	2.4	6.2	401	<b>2.9</b>	1.3	6.8	353	<b>3.0</b>	1.5	6.1	385
Unemployed	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Other**	<b>4.0</b>	2.1	7.4	294	<b>2.5</b>	1.2	5.0	292	<b>2.2</b>	1.1	4.4	325	<b>0.8</b>	0.2	2.5	282
<b>EDUCATION</b>																
K-11th Grade	*	*	*	*	<b>0.0</b>	0.0	0.0	68	*	*	*	*	*	*	*	*
12th Grade or GED	<b>8.4</b>	4.1	16.5	168	<b>1.7</b>	0.6	4.6	199	<b>2.9</b>	1.5	5.6	218	<b>2.5</b>	1.0	6.4	175
Some College	<b>5.8</b>	3.0	10.9	215	<b>4.1</b>	2.1	8.0	241	<b>2.8</b>	0.7	10.4	227	<b>3.0</b>	1.0	8.5	227
College Graduate+	<b>3.8</b>	1.9	7.3	311	<b>5.6</b>	3.2	9.7	199	<b>3.0</b>	1.0	8.4	201	<b>1.4</b>	0.5	3.8	242

\*Figure not reliable by BRFSS standards (n<50)

\*\*Other includes students, homemakers, retirees, and persons unable to work

Note: Definition of heavy drinking is males who consumed >60 drinks and females who consumed >30 drinks in the last 30 days.

This definition changed beginning in 2006.

# Illicit Drug Use

Idaho adults who used illicit drugs in the last 12 months, 2008

	Statewide			Public Health District								
				District 1			District 2			District 3		
	%	95% CI	n	%	95% CI	n	%	95% CI	n	%	95% CI	n
<b>TOTAL</b>	<b>4.1</b>	<b>3.2</b> <b>5.2</b>	<b>4,797</b>	<b>4.8</b>	<b>2.9</b> <b>7.8</b>	<b>679</b>	<b>2.7</b>	<b>1.5</b> <b>5.0</b>	<b>709</b>	<b>3.2</b>	<b>1.7</b> <b>5.7</b>	<b>699</b>
<b>SEX</b>												
Male	<b>5.4</b>	<b>3.8</b> <b>7.5</b>	<b>1,856</b>	<b>7.2</b>	<b>4.0</b> <b>12.6</b>	<b>264</b>	<b>2.6</b>	<b>1.0</b> <b>6.3</b>	<b>299</b>	<b>3.3</b>	<b>1.6</b> <b>6.9</b>	<b>250</b>
Female	<b>2.8</b>	<b>1.9</b> <b>4.0</b>	<b>2,941</b>	<b>2.5</b>	<b>1.1</b> <b>6.0</b>	<b>415</b>	<b>2.9</b>	<b>1.3</b> <b>6.4</b>	<b>410</b>	<b>3.0</b>	<b>1.1</b> <b>7.6</b>	<b>449</b>
<b>AGE</b>												
18-24	<b>7.7</b>	<b>4.0</b> <b>14.6</b>	<b>156</b>	*	*	*	*	*	*	*	*	*
25-34	<b>7.9</b>	<b>5.2</b> <b>11.8</b>	<b>597</b>	<b>8.1</b>	<b>3.0</b> <b>20.1</b>	<b>63</b>	<b>5.2</b>	<b>1.9</b> <b>13.3</b>	<b>81</b>	<b>5.5</b>	<b>2.2</b> <b>13.1</b>	<b>103</b>
35-44	<b>3.7</b>	<b>2.4</b> <b>5.9</b>	<b>723</b>	<b>7.3</b>	<b>3.2</b> <b>15.8</b>	<b>95</b>	<b>3.2</b>	<b>1.0</b> <b>9.9</b>	<b>103</b>	<b>3.1</b>	<b>1.1</b> <b>8.2</b>	<b>95</b>
45-54	<b>2.0</b>	<b>1.2</b> <b>3.4</b>	<b>975</b>	<b>2.0</b>	<b>0.7</b> <b>5.4</b>	<b>158</b>	<b>2.2</b>	<b>0.8</b> <b>6.0</b>	<b>144</b>	<b>1.1</b>	<b>0.3</b> <b>4.3</b>	<b>122</b>
55-64	<b>1.6</b>	<b>0.9</b> <b>2.9</b>	<b>1,020</b>	<b>3.5</b>	<b>1.4</b> <b>8.4</b>	<b>150</b>	<b>0.6</b>	<b>0.1</b> <b>3.9</b>	<b>161</b>	<b>4.3</b>	<b>1.5</b> <b>11.4</b>	<b>147</b>
65+	<b>0.9</b>	<b>0.4</b> <b>1.9</b>	<b>1,293</b>	<b>2.3</b>	<b>0.7</b> <b>7.5</b>	<b>185</b>	<b>0.5</b>	<b>0.1</b> <b>3.3</b>	<b>201</b>	<b>0.0</b>	<b>0.0</b> <b>0.0</b>	<b>204</b>
18-34	<b>7.8</b>	<b>5.5</b> <b>11.1</b>	<b>753</b>	<b>7.9</b>	<b>3.2</b> <b>18.2</b>	<b>84</b>	<b>5.0</b>	<b>2.0</b> <b>11.7</b>	<b>99</b>	<b>5.2</b>	<b>2.1</b> <b>12.6</b>	<b>124</b>
35-54	<b>2.9</b>	<b>2.0</b> <b>4.1</b>	<b>1,698</b>	<b>4.4</b>	<b>2.3</b> <b>8.5</b>	<b>253</b>	<b>2.7</b>	<b>1.2</b> <b>5.8</b>	<b>247</b>	<b>2.1</b>	<b>0.9</b> <b>4.8</b>	<b>217</b>
55+	<b>1.3</b>	<b>0.8</b> <b>2.0</b>	<b>2,313</b>	<b>2.9</b>	<b>1.4</b> <b>5.9</b>	<b>335</b>	<b>0.5</b>	<b>0.1</b> <b>2.0</b>	<b>362</b>	<b>2.0</b>	<b>0.7</b> <b>5.4</b>	<b>351</b>
<b>SEX and AGE</b>												
Male												
18-34	<b>9.7</b>	<b>6.0</b> <b>15.4</b>	<b>275</b>	*	*	*	*	*	*	*	*	*
35-54	<b>3.8</b>	<b>2.3</b> <b>6.1</b>	<b>669</b>	<b>5.5</b>	<b>2.1</b> <b>13.6</b>	<b>93</b>	<b>2.8</b>	<b>0.8</b> <b>8.9</b>	<b>115</b>	<b>2.4</b>	<b>0.7</b> <b>7.6</b>	<b>82</b>
55+	<b>1.9</b>	<b>1.1</b> <b>3.3</b>	<b>903</b>	<b>4.9</b>	<b>2.2</b> <b>10.4</b>	<b>140</b>	<b>0.5</b>	<b>0.1</b> <b>3.7</b>	<b>142</b>	<b>2.5</b>	<b>0.6</b> <b>9.7</b>	<b>116</b>
Female												
18-34	<b>5.8</b>	<b>3.5</b> <b>9.6</b>	<b>478</b>	<b>3.6</b>	<b>0.5</b> <b>21.2</b>	<b>54</b>	<b>5.9</b>	<b>1.8</b> <b>17.3</b>	<b>57</b>	<b>5.4</b>	<b>1.3</b> <b>20.0</b>	<b>75</b>
35-54	<b>1.9</b>	<b>1.3</b> <b>2.9</b>	<b>1,029</b>	<b>3.4</b>	<b>1.4</b> <b>8.0</b>	<b>160</b>	<b>2.6</b>	<b>0.9</b> <b>7.0</b>	<b>132</b>	<b>1.9</b>	<b>0.6</b> <b>5.7</b>	<b>135</b>
55+	<b>0.6</b>	<b>0.3</b> <b>1.5</b>	<b>1,410</b>	<b>1.0</b>	<b>0.1</b> <b>6.6</b>	<b>195</b>	<b>0.5</b>	<b>0.1</b> <b>3.4</b>	<b>220</b>	<b>1.5</b>	<b>0.3</b> <b>6.3</b>	<b>235</b>
<b>INCOME</b>												
Less than \$15,000	<b>4.1</b>	<b>2.1</b> <b>7.9</b>	<b>456</b>	<b>0.9</b>	<b>0.1</b> <b>6.0</b>	<b>74</b>	<b>5.8</b>	<b>1.4</b> <b>21.4</b>	<b>80</b>	<b>4.8</b>	<b>1.2</b> <b>17.5</b>	<b>75</b>
\$15,000 - \$24,999	<b>7.0</b>	<b>4.1</b> <b>11.7</b>	<b>778</b>	<b>12.8</b>	<b>6.1</b> <b>25.0</b>	<b>115</b>	<b>3.4</b>	<b>0.9</b> <b>12.3</b>	<b>123</b>	<b>2.4</b>	<b>0.7</b> <b>8.2</b>	<b>140</b>
\$25,000 - \$34,999	<b>5.2</b>	<b>2.7</b> <b>10.0</b>	<b>618</b>	<b>7.6</b>	<b>1.4</b> <b>32.8</b>	<b>69</b>	<b>4.3</b>	<b>1.0</b> <b>16.5</b>	<b>100</b>	<b>1.1</b>	<b>0.2</b> <b>7.1</b>	<b>100</b>
\$35,000 - \$49,999	<b>2.1</b>	<b>1.3</b> <b>3.5</b>	<b>815</b>	<b>1.4</b>	<b>0.3</b> <b>5.6</b>	<b>108</b>	<b>3.3</b>	<b>0.8</b> <b>12.0</b>	<b>126</b>	<b>1.5</b>	<b>0.4</b> <b>6.1</b>	<b>123</b>
\$50,000-\$74,999	<b>4.7</b>	<b>2.7</b> <b>8.0</b>	<b>752</b>	<b>1.0</b>	<b>0.2</b> <b>3.8</b>	<b>109</b>	<b>1.8</b>	<b>0.4</b> <b>7.5</b>	<b>105</b>	<b>9.9</b>	<b>3.6</b> <b>24.6</b>	<b>99</b>
\$75,000+	<b>3.3</b>	<b>1.7</b> <b>6.1</b>	<b>872</b>	<b>5.5</b>	<b>2.4</b> <b>12.1</b>	<b>126</b>	<b>0.9</b>	<b>0.1</b> <b>6.4</b>	<b>108</b>	<b>2.5</b>	<b>0.6</b> <b>10.3</b>	<b>85</b>
<b>EMPLOYMENT</b>												
Employed	<b>4.9</b>	<b>3.7</b> <b>6.6</b>	<b>2,563</b>	<b>5.8</b>	<b>3.2</b> <b>10.0</b>	<b>351</b>	<b>3.9</b>	<b>1.9</b> <b>7.9</b>	<b>366</b>	<b>4.4</b>	<b>2.2</b> <b>8.6</b>	<b>360</b>
Unemployed	<b>10.0</b>	<b>4.1</b> <b>22.2</b>	<b>165</b>	*	*	*	*	*	*	*	*	*
Other**	<b>1.7</b>	<b>1.0</b> <b>2.9</b>	<b>2,063</b>	<b>2.6</b>	<b>0.9</b> <b>7.6</b>	<b>299</b>	<b>0.9</b>	<b>0.3</b> <b>2.7</b>	<b>314</b>	<b>1.0</b>	<b>0.3</b> <b>3.3</b>	<b>309</b>
<b>EDUCATION</b>												
K-11th Grade	<b>4.7</b>	<b>2.2</b> <b>9.7</b>	<b>370</b>	<b>4.3</b>	<b>1.0</b> <b>16.9</b>	<b>51</b>	<b>3.8</b>	<b>0.5</b> <b>22.6</b>	<b>50</b>	<b>1.3</b>	<b>0.2</b> <b>8.9</b>	<b>81</b>
12th Grade or GED	<b>6.7</b>	<b>4.5</b> <b>9.9</b>	<b>1,402</b>	<b>9.0</b>	<b>4.6</b> <b>16.8</b>	<b>213</b>	<b>2.4</b>	<b>0.8</b> <b>6.9</b>	<b>235</b>	<b>3.1</b>	<b>1.3</b> <b>7.1</b>	<b>239</b>
Some College	<b>2.9</b>	<b>1.8</b> <b>4.5</b>	<b>1,544</b>	<b>2.8</b>	<b>1.0</b> <b>8.2</b>	<b>229</b>	<b>3.6</b>	<b>1.4</b> <b>8.9</b>	<b>213</b>	<b>3.4</b>	<b>1.0</b> <b>11.2</b>	<b>216</b>
College Graduate+	<b>2.6</b>	<b>1.5</b> <b>4.5</b>	<b>1,472</b>	<b>2.3</b>	<b>0.7</b> <b>7.2</b>	<b>183</b>	<b>1.9</b>	<b>0.4</b> <b>7.7</b>	<b>210</b>	<b>4.2</b>	<b>1.4</b> <b>11.5</b>	<b>163</b>

\*Figure not reliable by BRFSS standards (n<50)

\*\*Other includes students, homemakers, retirees, and persons unable to work

# Illicit Drug Use

Idaho adults who used illicit drugs in the last 12 months, 2008

	Public Health District															
	District 4				District 5				District 6				District 7			
	%	95% CI		n	%	95% CI		n	%	95% CI		n	%	95% CI		n
<b>TOTAL</b>	<b>6.3</b>	4.0	10.0	708	<b>2.8</b>	1.5	5.1	673	<b>3.8</b>	1.8	7.7	665	<b>1.3</b>	0.7	2.6	664
<b>SEX</b>																
Male	<b>9.0</b>	5.0	15.6	270	<b>3.4</b>	1.5	7.7	262	<b>5.2</b>	1.9	13.5	237	<b>1.0</b>	0.3	3.0	274
Female	<b>3.7</b>	1.7	7.7	438	<b>2.2</b>	1.0	5.1	411	<b>2.4</b>	1.0	5.7	428	<b>1.6</b>	0.7	3.7	390
<b>AGE</b>																
18-24	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
25-34	<b>11.6</b>	5.5	22.8	86	<b>7.4</b>	3.0	17.2	71	<b>5.8</b>	1.5	20.2	89	<b>4.2</b>	1.5	11.0	104
35-44	<b>4.7</b>	1.9	10.8	143	<b>3.0</b>	0.9	9.0	99	<b>1.0</b>	0.2	3.8	81	<b>1.1</b>	0.2	5.1	107
45-54	<b>2.0</b>	0.5	8.0	138	<b>2.5</b>	0.7	8.9	144	<b>2.8</b>	0.9	7.9	136	<b>1.8</b>	0.5	6.0	133
55-64	<b>0.7</b>	0.2	2.9	158	<b>0.8</b>	0.2	3.2	133	<b>0.9</b>	0.1	6.2	155	<b>0.0</b>	0.0	0.0	116
65+	<b>1.4</b>	0.3	5.4	156	<b>0.5</b>	0.1	3.5	199	<b>0.0</b>	0.0	0.0	176	<b>0.6</b>	0.1	4.4	172
18-34	<b>13.7</b>	7.5	23.5	107	<b>5.3</b>	2.1	12.7	92	<b>8.4</b>	3.4	19.1	113	<b>1.9</b>	0.7	5.1	134
35-54	<b>3.4</b>	1.6	7.1	281	<b>2.7</b>	1.1	6.3	243	<b>1.9</b>	0.8	4.6	217	<b>1.5</b>	0.6	3.8	240
55+	<b>1.0</b>	0.4	2.8	314	<b>0.6</b>	0.2	2.0	332	<b>0.4</b>	0.1	3.0	331	<b>0.3</b>	0.1	2.3	288
<b>SEX and AGE</b>																
Male																
18-34	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
35-54	<b>5.9</b>	2.6	13.1	106	<b>3.6</b>	1.1	10.9	87	<b>1.0</b>	0.3	4.0	82	<b>1.2</b>	0.2	5.8	104
55+	<b>1.4</b>	0.4	5.5	126	<b>1.0</b>	0.3	4.1	141	<b>0.9</b>	0.1	6.3	115	<b>0.7</b>	0.1	4.7	123
Female																
18-34	<b>9.8</b>	4.1	21.7	70	<b>5.1</b>	1.6	14.9	60	<b>4.3</b>	1.2	14.0	74	<b>2.6</b>	0.8	8.2	88
35-54	<b>0.9</b>	0.3	3.0	175	<b>1.8</b>	0.5	6.0	156	<b>2.8</b>	0.9	8.2	135	<b>1.7</b>	0.5	5.6	136
55+	<b>0.7</b>	0.2	2.8	188	<b>0.3</b>	0.0	2.1	191	<b>0.0</b>	0.0	0.0	216	<b>0.0</b>	0.0	0.0	165
<b>INCOME</b>																
Less than \$15,000	<b>4.5</b>	0.9	20.4	53	<b>0.9</b>	0.1	6.1	58	<b>12.7</b>	4.0	33.8	64	<b>0.0</b>	0.0	0.0	52
\$15,000 - \$24,999	<b>16.3</b>	6.6	34.7	73	<b>0.6</b>	0.1	4.2	114	<b>0.6</b>	0.1	4.3	109	<b>2.6</b>	0.8	8.5	104
\$25,000 - \$34,999	<b>8.0</b>	2.3	24.5	88	<b>4.1</b>	1.1	14.0	91	<b>6.2</b>	1.5	21.6	97	<b>2.7</b>	0.6	10.6	73
\$35,000 - \$49,999	<b>1.9</b>	0.7	5.2	129	<b>3.4</b>	1.1	10.1	108	<b>2.4</b>	0.3	15.0	114	<b>2.1</b>	0.5	8.4	107
\$50,000-\$74,999	<b>8.0</b>	3.2	18.3	106	<b>3.6</b>	0.8	14.1	106	<b>1.4</b>	0.2	9.4	102	<b>1.7</b>	0.4	6.7	125
\$75,000+	<b>3.3</b>	0.9	11.6	193	<b>5.0</b>	1.6	14.6	112	<b>4.6</b>	0.7	25.6	113	<b>0.0</b>	0.0	0.0	135
<b>EMPLOYMENT</b>																
Employed	<b>6.5</b>	3.6	11.5	395	<b>3.7</b>	1.9	7.1	381	<b>5.5</b>	2.4	12.0	339	<b>2.0</b>	1.0	4.1	371
Unemployed	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Other**	<b>3.0</b>	1.1	7.6	282	<b>1.3</b>	0.3	5.9	275	<b>1.1</b>	0.3	3.9	309	<b>0.2</b>	0.0	1.7	275
<b>EDUCATION</b>																
K-11th Grade	*	*	*	*	<b>2.8</b>	0.7	11.1	66	*	*	*	*	*	*	*	*
12th Grade or GED	<b>13.7</b>	7.2	24.8	165	<b>3.1</b>	0.8	11.4	184	<b>3.9</b>	1.5	9.6	202	<b>2.3</b>	0.8	6.5	164
Some College	<b>3.1</b>	1.4	7.0	209	<b>2.6</b>	0.9	6.8	232	<b>3.5</b>	0.7	15.7	221	<b>1.1</b>	0.3	3.8	224
College Graduate+	<b>2.9</b>	0.9	8.7	297	<b>2.9</b>	1.0	7.6	189	<b>3.0</b>	0.7	12.2	194	<b>0.6</b>	0.1	2.6	236

\*Figure not reliable by BRFSS standards (n<50)

\*\*Other includes students, homemakers, retirees, and persons unable to work

# Seat Belt Use

## Idaho adults who did not always wear a seat belt, 2008

	Statewide			Public Health District												
				District 1			District 2			District 3						
	%	95% CI		n	%	95% CI		n	%	95% CI		n	%	95% CI		n
<b>TOTAL</b>	<b>23.4</b>	21.8	25.1	5,039	<b>22.6</b>	18.8	26.9	720	<b>24.3</b>	20.5	28.5	746	<b>20.0</b>	16.1	24.7	739
<b>SEX</b>																
Male	<b>29.4</b>	26.7	32.2	1,952	<b>29.0</b>	22.7	36.3	277	<b>32.3</b>	26.1	39.3	315	<b>24.7</b>	18.9	31.6	265
Female	<b>17.6</b>	15.8	19.6	3,087	<b>16.4</b>	12.5	21.4	443	<b>16.2</b>	12.5	20.8	431	<b>15.4</b>	10.4	22.2	474
<b>AGE</b>																
18-24	<b>35.3</b>	27.2	44.3	163	*	*	*	*	*	*	*	*	*	*	*	*
25-34	<b>24.4</b>	20.6	28.5	625	<b>26.6</b>	16.1	40.6	67	<b>24.5</b>	15.9	35.8	85	<b>20.4</b>	13.2	30.1	107
35-44	<b>19.6</b>	16.6	23.0	768	<b>18.0</b>	11.1	28.0	103	<b>23.9</b>	16.6	33.2	110	<b>19.0</b>	11.9	28.8	103
45-54	<b>23.6</b>	20.7	26.7	1,021	<b>22.0</b>	15.9	29.7	167	<b>31.6</b>	24.0	40.4	150	<b>16.6</b>	10.7	24.9	131
55-64	<b>23.0</b>	20.0	26.2	1,056	<b>19.3</b>	13.7	26.4	161	<b>20.1</b>	14.2	27.8	164	<b>17.7</b>	11.7	25.8	150
65+	<b>18.2</b>	15.9	20.6	1,367	<b>22.4</b>	16.2	30.1	193	<b>16.1</b>	11.4	22.2	214	<b>16.2</b>	10.8	23.5	218
18-34	<b>28.3</b>	24.4	32.5	788	<b>28.4</b>	18.6	40.7	89	<b>27.4</b>	19.0	37.7	105	<b>25.1</b>	16.4	36.4	129
35-54	<b>21.6</b>	19.5	23.8	1,789	<b>20.2</b>	15.3	26.1	270	<b>28.1</b>	22.5	34.4	260	<b>17.9</b>	12.9	24.1	234
55+	<b>20.4</b>	18.6	22.4	2,423	<b>20.9</b>	16.5	26.0	354	<b>17.8</b>	13.9	22.5	378	<b>16.9</b>	12.7	22.1	368
<b>SEX and AGE</b>																
Male																
18-34	<b>34.2</b>	28.0	41.0	288	*	*	*	*	*	*	*	*	<b>25.5</b>	14.3	41.3	52
35-54	<b>26.6</b>	23.2	30.4	712	<b>24.5</b>	16.8	34.4	102	<b>36.3</b>	27.7	45.9	123	<b>22.1</b>	14.3	32.7	88
55+	<b>27.1</b>	24.0	30.5	942	<b>30.1</b>	22.7	38.7	144	<b>22.8</b>	16.4	30.8	147	<b>27.5</b>	19.6	37.1	121
Female																
18-34	<b>22.1</b>	17.6	27.3	500	<b>22.8</b>	12.8	37.3	59	<b>15.7</b>	8.5	27.4	60	<b>24.7</b>	13.0	41.9	77
35-54	<b>16.5</b>	14.2	19.1	1,077	<b>15.9</b>	10.7	23.1	168	<b>20.0</b>	13.7	28.2	137	<b>13.6</b>	8.5	21.2	146
55+	<b>14.4</b>	12.5	16.6	1,481	<b>12.6</b>	8.4	18.4	210	<b>13.2</b>	9.1	18.7	231	<b>7.3</b>	4.6	11.6	247
<b>INCOME</b>																
Less than \$15,000	<b>25.8</b>	20.1	32.5	479	<b>36.6</b>	23.0	52.7	75	<b>21.5</b>	12.3	34.8	85	<b>18.7</b>	10.3	31.7	80
\$15,000 - \$24,999	<b>26.5</b>	22.0	31.6	813	<b>27.6</b>	18.2	39.6	118	<b>28.5</b>	18.8	40.5	132	<b>29.9</b>	19.7	42.5	145
\$25,000 - \$34,999	<b>28.5</b>	23.3	34.4	652	<b>33.4</b>	20.6	49.1	77	<b>17.9</b>	11.3	27.2	104	<b>21.1</b>	12.4	33.7	107
\$35,000 - \$49,999	<b>24.2</b>	20.9	27.9	849	<b>23.0</b>	15.4	32.8	116	<b>25.2</b>	16.6	36.3	126	<b>18.6</b>	11.6	28.4	130
\$50,000-\$74,999	<b>22.1</b>	18.7	26.0	786	<b>19.5</b>	11.9	30.3	114	<b>26.9</b>	17.9	38.3	109	<b>9.0</b>	4.7	16.5	105
\$75,000+	<b>18.3</b>	15.3	21.7	902	<b>10.5</b>	5.3	19.6	133	<b>23.4</b>	15.7	33.3	111	<b>15.8</b>	9.2	25.9	86
<b>EMPLOYMENT</b>																
Employed	<b>25.2</b>	23.1	27.4	2,694	<b>22.4</b>	17.3	28.5	376	<b>27.5</b>	22.4	33.3	382	<b>18.9</b>	14.6	24.1	379
Unemployed	<b>26.6</b>	17.7	37.9	172	*	*	*	*	*	*	*	*	*	*	*	*
Other**	<b>19.8</b>	17.2	22.6	2,166	<b>22.8</b>	17.2	29.6	315	<b>16.1</b>	11.4	22.1	333	<b>18.0</b>	12.0	26.0	328
<b>EDUCATION</b>																
K-11th Grade	<b>28.5</b>	22.2	35.7	395	<b>32.6</b>	16.2	54.7	51	<b>17.6</b>	7.9	34.4	53	<b>25.2</b>	14.2	40.5	92
12th Grade or GED	<b>28.7</b>	25.3	32.3	1,504	<b>24.6</b>	18.0	32.6	231	<b>29.5</b>	22.5	37.7	251	<b>26.8</b>	18.8	36.7	253
Some College	<b>23.6</b>	20.9	26.5	1,605	<b>20.5</b>	14.9	27.6	243	<b>29.1</b>	22.2	37.1	223	<b>16.8</b>	11.7	23.7	225
College Graduate+	<b>16.7</b>	14.6	19.2	1,526	<b>19.7</b>	13.9	27.2	192	<b>15.7</b>	10.8	22.3	218	<b>11.1</b>	6.5	18.3	169

\*Figure not reliable by BRFSS standards (n<50)

\*\*Other includes students, homemakers, retirees, and persons unable to work

# Seat Belt Use

## Idaho adults who did not always wear a seat belt, 2008

	Public Health District															
	District 4			District 5			District 6			District 7						
	%	95% CI		n	%	95% CI		n	%	95% CI		n	%	95% CI		n
<b>TOTAL</b>	<b>14.7</b>	11.6	18.5	736	<b>27.2</b>	23.2	31.6	716	<b>38.5</b>	33.2	44.1	695	<b>31.3</b>	26.8	36.2	687
<b>SEX</b>																
Male	<b>19.9</b>	14.5	26.7	283	<b>34.8</b>	28.1	42.1	280	<b>43.2</b>	35.4	51.3	250	<b>38.6</b>	31.1	46.7	282
Female	<b>9.6</b>	6.9	13.1	453	<b>19.6</b>	15.6	24.4	436	<b>34.1</b>	26.9	42.0	445	<b>24.4</b>	19.6	30.0	405
<b>AGE</b>																
18-24	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
25-34	<b>10.2</b>	5.2	19.1	86	<b>34.1</b>	23.3	46.8	78	<b>41.9</b>	31.0	53.6	92	<b>43.4</b>	32.7	54.6	110
35-44	<b>14.9</b>	9.7	22.3	147	<b>21.2</b>	14.0	30.8	106	<b>29.0</b>	19.9	40.2	88	<b>23.3</b>	16.3	32.2	111
45-54	<b>13.5</b>	8.6	20.4	143	<b>29.0</b>	21.4	38.1	151	<b>41.0</b>	32.2	50.4	139	<b>31.7</b>	23.8	40.8	140
55-64	<b>20.3</b>	13.8	28.9	162	<b>23.5</b>	16.8	31.9	139	<b>30.1</b>	22.9	38.5	161	<b>36.4</b>	27.5	46.3	119
65+	<b>8.4</b>	5.2	13.3	168	<b>22.3</b>	16.9	28.9	213	<b>24.2</b>	18.0	31.5	186	<b>25.7</b>	19.4	33.3	175
18-34	<b>15.4</b>	8.8	25.5	109	<b>34.3</b>	24.6	45.4	100	<b>51.5</b>	39.5	63.4	116	<b>34.8</b>	25.2	46.0	140
35-54	<b>14.2</b>	10.4	19.1	290	<b>25.3</b>	19.8	31.8	257	<b>35.3</b>	28.7	42.5	227	<b>27.7</b>	22.2	34.0	251
55+	<b>14.5</b>	10.6	19.6	330	<b>22.9</b>	18.5	27.9	352	<b>27.0</b>	22.1	32.4	347	<b>30.9</b>	25.3	37.0	294
<b>SEX and AGE</b>																
Male																
18-34	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
35-54	<b>19.4</b>	13.1	27.8	111	<b>28.5</b>	19.9	38.9	94	<b>45.0</b>	34.1	56.4	86	<b>30.9</b>	22.6	40.7	108
55+	<b>18.3</b>	11.9	27.0	134	<b>27.8</b>	20.8	35.9	148	<b>31.9</b>	23.8	41.1	123	<b>39.8</b>	30.8	49.6	125
Female																
18-34	<b>8.4</b>	3.8	17.6	72	<b>17.8</b>	9.9	30.0	64	<b>53.0</b>	37.0	68.3	76	<b>25.7</b>	16.4	37.7	92
35-54	<b>8.9</b>	5.4	14.2	179	<b>22.1</b>	15.7	30.1	163	<b>25.8</b>	18.7	34.5	141	<b>24.5</b>	17.7	32.9	143
55+	<b>11.2</b>	7.0	17.3	196	<b>18.5</b>	13.3	25.0	204	<b>22.7</b>	17.4	29.0	224	<b>22.8</b>	16.6	30.5	169
<b>INCOME</b>																
Less than \$15,000	<b>10.1</b>	4.6	20.5	54	<b>21.1</b>	11.7	35.2	66	<b>42.2</b>	22.8	64.4	67	<b>36.3</b>	18.4	58.9	52
\$15,000 - \$24,999	<b>15.5</b>	6.2	33.6	79	<b>22.3</b>	13.7	34.0	122	<b>40.3</b>	29.0	52.8	113	<b>32.0</b>	20.9	45.7	104
\$25,000 - \$34,999	<b>21.0</b>	11.0	36.3	88	<b>34.6</b>	24.5	46.4	99	<b>44.9</b>	29.1	61.7	99	<b>33.6</b>	20.4	50.1	78
\$35,000 - \$49,999	<b>13.9</b>	8.6	21.6	132	<b>39.2</b>	28.5	51.0	113	<b>31.5</b>	22.4	42.3	120	<b>33.2</b>	24.0	44.0	112
\$50,000-\$74,999	<b>16.6</b>	9.7	26.9	112	<b>20.3</b>	12.5	31.3	109	<b>36.0</b>	26.5	46.8	109	<b>39.5</b>	30.0	49.9	128
\$75,000+	<b>11.9</b>	7.6	18.2	198	<b>26.1</b>	17.5	37.1	116	<b>42.1</b>	30.2	55.0	117	<b>20.7</b>	14.1	29.5	141
<b>EMPLOYMENT</b>																
Employed	<b>15.2</b>	11.3	20.0	409	<b>30.7</b>	25.4	36.7	407	<b>44.3</b>	37.5	51.2	354	<b>37.3</b>	31.4	43.7	387
Unemployed	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Other**	<b>13.8</b>	8.7	21.2	295	<b>20.1</b>	14.9	26.6	291	<b>30.6</b>	22.1	40.6	323	<b>21.9</b>	15.7	29.7	281
<b>EDUCATION</b>																
K-11th Grade	*	*	*	*	<b>21.2</b>	11.2	36.5	70	*	*	*	*	*	*	*	*
12th Grade or GED	<b>19.1</b>	11.4	30.2	172	<b>36.8</b>	28.4	46.0	206	<b>41.5</b>	32.7	50.8	216	<b>38.0</b>	28.5	48.5	175
Some College	<b>12.3</b>	8.4	17.8	217	<b>30.3</b>	23.7	37.8	242	<b>44.9</b>	35.2	54.9	227	<b>29.3</b>	21.7	38.4	228
College Graduate+	<b>12.0</b>	8.5	16.8	307	<b>15.4</b>	10.4	22.1	196	<b>23.6</b>	17.0	31.8	201	<b>28.4</b>	22.0	35.8	243

\*Figure not reliable by BRFSS standards (n<50)

\*\*Other includes students, homemakers, retirees, and persons unable to work

# Colorectal Cancer Screening

Idaho adults aged 50 and older who had never had a sigmoidoscopy or colonoscopy, 2008

	Statewide			Public Health District												
				District 1			District 2			District 3						
	%	95% CI		n	%	95% CI		n	%	95% CI		n	%	95% CI		n
<b>TOTAL</b>	<b>44.4</b>	42.3	46.6	2,946	<b>45.9</b>	40.8	51.1	437	<b>38.4</b>	33.4	43.6	450	<b>47.7</b>	42.3	53.2	428
<b>SEX</b>																
Male	<b>45.4</b>	42.1	48.8	1,169	<b>46.2</b>	38.6	53.9	185	<b>38.1</b>	30.6	46.2	187	<b>54.4</b>	45.5	63.0	145
Female	<b>43.4</b>	40.7	46.2	1,777	<b>45.7</b>	39.0	52.5	252	<b>38.7</b>	32.4	45.4	263	<b>41.7</b>	35.3	48.4	283
<b>AGE</b>																
50-64	<b>52.8</b>	49.9	55.7	1,597	<b>52.7</b>	45.9	59.3	243	<b>46.2</b>	39.3	53.2	241	<b>57.4</b>	49.8	64.6	215
65+	<b>31.4</b>	28.6	34.3	1,349	<b>36.2</b>	29.0	44.0	194	<b>28.4</b>	21.6	36.2	209	<b>34.6</b>	27.4	42.5	213
<b>SEX and AGE</b>																
Male																
50-64	<b>53.1</b>	48.6	57.5	650	<b>51.9</b>	42.4	61.4	118	<b>45.6</b>	35.7	55.9	106	<b>63.2</b>	51.4	73.6	79
65+	<b>32.3</b>	28.0	37.1	519	<b>37.1</b>	25.7	50.1	67	<b>27.0</b>	16.6	40.6	81	<b>40.6</b>	28.2	54.3	66
Female																
50-64	<b>52.6</b>	48.8	56.3	947	<b>53.5</b>	44.1	62.6	125	<b>46.8</b>	37.8	56.1	135	<b>51.5</b>	42.1	60.8	136
65+	<b>30.6</b>	27.2	34.3	830	<b>35.3</b>	27.0	44.6	127	<b>29.6</b>	21.7	39.0	128	<b>29.8</b>	22.2	38.8	147
<b>INCOME</b>																
Less than \$15,000	<b>57.4</b>	50.8	63.8	302	<b>46.1</b>	32.1	60.7	52	<b>58.1</b>	42.7	72.1	50	<b>65.8</b>	49.7	79.0	50
\$15,000 - \$24,999	<b>46.8</b>	41.5	52.2	527	<b>58.0</b>	45.6	69.5	78	<b>41.5</b>	29.6	54.5	87	<b>45.5</b>	33.3	58.3	89
\$25,000 - \$34,999	<b>40.9</b>	35.3	46.8	394	*	*	*	*	<b>31.5</b>	20.7	44.8	65	<b>42.7</b>	30.1	56.3	69
\$35,000 - \$49,999	<b>45.2</b>	40.0	50.5	486	<b>52.3</b>	39.7	64.7	65	<b>40.1</b>	28.8	52.6	77	<b>48.9</b>	36.7	61.2	76
\$50,000-\$74,999	<b>41.8</b>	36.4	47.4	394	<b>38.0</b>	26.2	51.5	62	<b>38.4</b>	26.0	52.5	58	<b>51.7</b>	37.3	65.9	55
\$75,000+	<b>44.3</b>	39.0	49.8	461	<b>36.3</b>	25.2	49.0	69	<b>31.7</b>	20.5	45.6	63	*	*	*	*
<b>EMPLOYMENT</b>																
Employed	<b>56.0</b>	52.6	59.3	1,197	<b>55.4</b>	47.4	63.2	174	<b>51.5</b>	43.4	59.4	179	<b>61.6</b>	52.8	69.6	158
Unemployed	<b>69.7</b>	57.5	79.7	78	*	*	*	*	*	*	*	*	*	*	*	*
Other**	<b>32.6</b>	30.1	35.3	1,669	<b>36.9</b>	30.5	43.8	246	<b>27.6</b>	21.6	34.6	261	<b>35.0</b>	28.6	42.1	258
<b>EDUCATION</b>																
K-11th Grade	<b>56.3</b>	48.3	64.1	216	*	*	*	*	*	*	*	*	*	*	*	*
12th Grade or GED	<b>48.5</b>	44.7	52.3	938	<b>50.8</b>	41.4	60.1	128	<b>41.0</b>	32.7	49.8	153	<b>57.0</b>	48.5	65.1	174
Some College	<b>44.9</b>	41.1	48.7	929	<b>44.5</b>	36.3	53.1	157	<b>43.0</b>	34.0	52.4	140	<b>42.4</b>	32.6	52.9	117
College Graduate+	<b>37.3</b>	33.5	41.3	858	<b>40.0</b>	30.7	50.2	115	<b>29.2</b>	21.3	38.7	127	<b>36.2</b>	25.7	48.1	91

\*Figure not reliable by BRFSS standards (n<50)

\*\*Other includes students, homemakers, retirees, and persons unable to work

# Colorectal Cancer Screening

Idaho adults aged 50 and older who had never had a sigmoidoscopy or colonoscopy, 2008

	Public Health District															
	District 4			District 5			District 6			District 7						
	%	95% CI		n	%	95% CI		n	%	95% CI		n	%	95% CI		n
<b>TOTAL</b>	<b>35.6</b>	30.5	41.1	406	<b>49.2</b>	44.0	54.4	433	<b>51.2</b>	45.8	56.5	425	<b>50.0</b>	44.3	55.6	367
<b>SEX</b>																
Male	<b>34.1</b>	26.3	43.0	160	<b>50.2</b>	42.2	58.2	178	<b>51.0</b>	42.4	59.5	156	<b>51.7</b>	43.1	60.3	158
Female	<b>36.9</b>	30.4	44.0	246	<b>48.2</b>	41.6	54.9	255	<b>51.4</b>	44.9	57.8	269	<b>48.3</b>	41.0	55.6	209
<b>AGE</b>																
50-64	<b>45.3</b>	38.4	52.5	240	<b>59.3</b>	52.1	66.1	226	<b>58.5</b>	51.4	65.3	240	<b>57.3</b>	49.5	64.8	192
65+	<b>17.9</b>	12.7	24.7	166	<b>34.3</b>	27.7	41.6	207	<b>39.8</b>	32.4	47.7	185	<b>38.3</b>	30.8	46.3	175
<b>SEX and AGE</b>																
Male																
50-64	<b>45.0</b>	34.1	56.3	88	<b>56.4</b>	45.3	66.9	88	<b>56.0</b>	44.8	66.7	91	<b>59.4</b>	47.3	70.4	80
65+	<b>13.0</b>	6.9	23.2	72	<b>40.1</b>	29.8	51.3	90	<b>42.1</b>	29.9	55.3	65	<b>38.0</b>	27.3	49.9	78
Female																
50-64	<b>45.7</b>	37.0	54.6	152	<b>62.4</b>	53.4	70.6	138	<b>61.2</b>	52.5	69.2	149	<b>55.0</b>	45.1	64.6	112
65+	<b>21.8</b>	14.4	31.7	94	<b>29.5</b>	21.5	38.9	117	<b>37.9</b>	29.2	47.5	120	<b>38.5</b>	28.6	49.5	97
<b>INCOME</b>																
Less than \$15,000	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
\$15,000 - \$24,999	*	*	*	*	<b>54.0</b>	42.8	64.9	88	<b>46.8</b>	33.7	60.3	76	<b>44.2</b>	31.1	58.2	60
\$25,000 - \$34,999	<b>29.4</b>	17.3	45.4	52	<b>56.1</b>	42.1	69.2	57	<b>50.1</b>	36.0	64.2	55	<b>39.2</b>	25.7	54.6	50
\$35,000 - \$49,999	<b>32.5</b>	22.1	44.9	82	<b>48.2</b>	34.0	62.6	53	<b>57.9</b>	44.8	70.0	70	<b>50.4</b>	36.2	64.5	63
\$50,000-\$74,999	*	*	*	*	<b>55.4</b>	41.7	68.3	59	<b>51.1</b>	37.6	64.5	58	<b>52.8</b>	38.5	66.6	53
\$75,000+	<b>44.5</b>	34.1	55.4	101	<b>37.7</b>	25.6	51.6	67	<b>53.1</b>	39.7	66.1	65	<b>54.7</b>	42.0	66.8	65
<b>EMPLOYMENT</b>																
Employed	<b>47.7</b>	39.5	56.1	176	<b>61.3</b>	53.4	68.7	189	<b>62.4</b>	53.9	70.1	166	<b>59.8</b>	51.3	67.8	155
Unemployed	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Other**	<b>21.8</b>	16.4	28.2	220	<b>34.3</b>	28.1	41.1	232	<b>41.4</b>	34.8	48.4	249	<b>40.3</b>	33.0	48.1	203
<b>EDUCATION</b>																
K-11th Grade	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
12th Grade or GED	<b>35.8</b>	26.2	46.6	99	<b>51.9</b>	42.5	61.2	131	<b>54.8</b>	45.4	63.9	139	<b>49.0</b>	39.3	58.8	114
Some College	<b>39.5</b>	29.9	50.0	120	<b>53.2</b>	44.2	62.0	143	<b>52.0</b>	42.8	61.1	133	<b>44.2</b>	34.3	54.5	119
College Graduate+	<b>31.7</b>	24.1	40.3	169	<b>36.3</b>	27.2	46.5	117	<b>43.2</b>	33.7	53.2	121	<b>52.0</b>	42.2	61.7	118

\*Figure not reliable by BRFSS standards (n<50)

\*\*Other includes students, homemakers, retirees, and persons unable to work

# Breast Cancer Screening

Idaho females aged 40 and older who had not had a mammogram and clinical breast exam in the last two years, 2008

	Statewide			Public Health District												
				District 1			District 2			District 3						
	%	95% CI		n	%	95% CI		n	%	95% CI		n	%	95% CI		n
<b>TOTAL FEMALES</b>	<b>36.8</b>	34.5	39.2	2,287	<b>39.4</b>	33.8	45.4	326	<b>39.2</b>	33.6	45.1	339	<b>42.7</b>	36.8	48.8	351
<b>AGE</b>																
40-49	<b>44.7</b>	39.8	49.7	521	<b>43.4</b>	32.0	55.5	80	<b>52.2</b>	40.1	64.0	77	<b>51.2</b>	38.8	63.5	70
50-64	<b>32.5</b>	29.1	36.2	946	<b>33.5</b>	25.1	43.0	124	<b>34.2</b>	25.9	43.5	135	<b>38.0</b>	29.2	47.8	135
65+	<b>34.5</b>	30.9	38.4	820	<b>43.8</b>	34.6	53.4	122	<b>33.5</b>	25.5	42.7	127	<b>39.4</b>	31.1	48.4	146
<b>INCOME</b>																
Less than \$15,000	<b>49.7</b>	42.8	56.8	272	*	*	*	*	<b>48.8</b>	33.5	64.4	50	*	*	*	*
\$15,000 - \$24,999	<b>50.3</b>	44.2	56.3	408	<b>54.5</b>	40.0	68.3	58	<b>43.5</b>	31.1	56.8	68	<b>52.6</b>	40.1	64.9	85
\$25,000 - \$34,999	<b>42.1</b>	35.6	48.9	295	*	*	*	*	*	*	*	*	<b>48.3</b>	33.4	63.5	51
\$35,000 - \$49,999	<b>34.7</b>	29.1	40.8	364	*	*	*	*	<b>43.7</b>	30.3	58.1	59	<b>35.4</b>	22.3	51.0	55
\$50,000-\$74,999	<b>32.0</b>	26.1	38.6	295	*	*	*	*	*	*	*	*	*	*	*	*
\$75,000+	<b>24.3</b>	19.4	29.9	340	*	*	*	*	*	*	*	*	*	*	*	*
<b>EMPLOYMENT</b>																
Employed	<b>36.9</b>	33.5	40.4	1,052	<b>37.6</b>	29.6	46.4	152	<b>42.0</b>	33.3	51.3	142	<b>44.2</b>	35.5	53.3	153
Unemployed	<b>51.3</b>	36.9	65.5	61	*	*	*	*	*	*	*	*	*	*	*	*
Other**	<b>35.8</b>	32.6	39.1	1,172	<b>40.4</b>	32.4	49.0	160	<b>36.0</b>	28.9	43.8	188	<b>39.5</b>	31.9	47.6	191
<b>EDUCATION</b>																
K-11th Grade	<b>53.2</b>	43.9	62.3	156	*	*	*	*	*	*	*	*	*	*	*	*
12th Grade or GED	<b>41.3</b>	37.1	45.5	772	<b>48.1</b>	37.9	58.4	106	<b>41.8</b>	33.0	51.2	132	<b>43.7</b>	34.4	53.6	143
Some College	<b>35.5</b>	31.6	39.7	763	<b>39.3</b>	30.0	49.5	120	<b>36.8</b>	27.1	47.7	99	<b>40.8</b>	30.6	51.8	107
College Graduate+	<b>29.8</b>	25.7	34.3	594	<b>22.8</b>	14.6	33.7	81	<b>35.6</b>	24.9	48.1	87	<b>36.9</b>	24.9	50.7	66

\*Figure not reliable by BRFSS standards (n<50)

\*\*Other includes students, homemakers, retirees, and persons unable to work

# Breast Cancer Screening

Idaho females aged 40 and older who had not had a mammogram and clinical breast exam in the last two years, 2008

	Public Health District															
	District 4			District 5			District 6			District 7						
	%	95% CI		n	%	95% CI		n	%	95% CI		n	%	95% CI		n
<b>TOTAL FEMALES</b>	<b>26.8</b>	21.8	32.5	330	<b>40.9</b>	35.2	46.9	331	<b>39.0</b>	33.3	45.0	336	<b>42.0</b>	35.7	48.5	274
<b>AGE</b>																
40-49	<b>34.1</b>	24.4	45.5	85	<b>50.6</b>	38.6	62.6	76	<b>51.8</b>	39.0	64.4	66	<b>47.9</b>	35.4	60.7	67
50-64	<b>23.8</b>	16.9	32.4	152	<b>40.9</b>	32.2	50.2	138	<b>28.8</b>	21.6	37.2	151	<b>40.4</b>	31.0	50.6	111
65+	<b>22.3</b>	14.3	33.0	93	<b>32.0</b>	23.8	41.4	117	<b>40.7</b>	31.7	50.4	119	<b>37.6</b>	27.9	48.3	96
<b>INCOME</b>																
Less than \$15,000	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
\$15,000 - \$24,999	*	*	*	*	<b>44.5</b>	31.2	58.6	58	<b>52.2</b>	38.0	66.0	58	*	*	*	*
\$25,000 - \$34,999	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
\$35,000 - \$49,999	<b>27.6</b>	17.4	41.0	70	*	*	*	*	<b>27.0</b>	15.5	42.7	52	*	*	*	*
\$50,000-\$74,999	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
\$75,000+	<b>12.3</b>	6.2	23.1	76	*	*	*	*	*	*	*	*	<b>45.9</b>	31.8	60.8	50
<b>EMPLOYMENT</b>																
Employed	<b>27.2</b>	20.3	35.4	163	<b>44.1</b>	35.9	52.6	167	<b>33.6</b>	25.2	43.0	140	<b>41.2</b>	32.5	50.6	135
Unemployed	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Other**	<b>25.2</b>	18.4	33.6	157	<b>35.7</b>	28.2	43.9	156	<b>43.9</b>	36.2	51.9	186	<b>41.8</b>	33.1	51.0	134
<b>EDUCATION</b>																
K-11th Grade	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
12th Grade or GED	<b>38.6</b>	27.6	51.0	84	<b>42.2</b>	31.6	53.5	100	<b>35.4</b>	26.6	45.4	121	<b>37.0</b>	26.7	48.5	86
Some College	<b>24.6</b>	17.0	34.3	114	<b>38.8</b>	30.0	48.4	123	<b>37.2</b>	27.5	48.0	107	<b>45.7</b>	34.9	56.8	93
College Graduate+	<b>20.3</b>	13.4	29.4	119	<b>35.1</b>	24.5	47.4	77	<b>38.6</b>	27.7	50.8	83	<b>44.2</b>	32.9	56.2	81

\*Figure not reliable by BRFSS standards (n<50)

\*\*Other includes students, homemakers, retirees, and persons unable to work

# Cervical Cancer Screening

Idaho females who had not had a Pap test in the last three years, 2008

	Statewide			Public Health District												
				District 1			District 2			District 3						
	%	95% CI		n	%	95% CI		n	%	95% CI		n	%	95% CI		n
<b>TOTAL FEMALES</b>	<b>23.1</b>	20.5	26.1	1,980	<b>16.2</b>	12.3	21.0	307	<b>17.7</b>	13.2	23.3	274	<b>28.9</b>	21.3	38.1	281
<b>AGE</b>																
18-24	<b>46.6</b>	35.2	58.3	97	*	*	*	*	*	*	*	*	*	*	*	*
25-34	<b>11.9</b>	8.4	16.7	388	*	*	*	*	<b>6.5</b>	2.1	18.7	50	<b>14.6</b>	6.8	28.5	62
35-44	<b>16.9</b>	13.3	21.2	394	<b>14.4</b>	7.8	25.0	60	*	*	*	*	<b>16.0</b>	8.2	28.8	58
45-54	<b>19.2</b>	15.1	24.0	403	<b>24.6</b>	15.5	36.6	74	<b>25.9</b>	15.6	39.8	55	*	*	*	*
55-64	<b>21.8</b>	17.2	27.2	336	*	*	*	*	<b>20.1</b>	10.9	34.2	51	*	*	*	*
65+	<b>40.3</b>	34.5	46.3	348	<b>35.3</b>	23.8	48.6	63	<b>33.9</b>	22.7	47.3	62	<b>48.0</b>	34.1	62.1	57
18-34	<b>24.5</b>	19.3	30.6	485	<b>5.9</b>	1.9	17.0	58	<b>8.7</b>	3.2	21.6	57	<b>34.6</b>	20.8	51.6	74
35-54	<b>18.0</b>	15.2	21.1	797	<b>19.5</b>	13.5	27.4	134	<b>21.4</b>	14.3	30.7	104	<b>18.6</b>	11.6	28.4	102
55+	<b>30.5</b>	26.7	34.5	684	<b>26.1</b>	18.1	36.1	110	<b>28.3</b>	20.4	37.9	113	<b>35.2</b>	26.1	45.6	103
<b>INCOME</b>																
Less than \$15,000	<b>37.6</b>	27.2	49.4	195	*	*	*	*	*	*	*	*	*	*	*	*
\$15,000 - \$24,999	<b>27.3</b>	20.9	34.7	325	<b>21.9</b>	11.8	36.8	52	*	*	*	*	<b>45.4</b>	26.9	65.2	61
\$25,000 - \$34,999	<b>31.1</b>	22.5	41.4	249	*	*	*	*	*	*	*	*	*	*	*	*
\$35,000 - \$49,999	<b>22.1</b>	16.8	28.4	315	*	*	*	*	*	*	*	*	*	*	*	*
\$50,000-\$74,999	<b>11.1</b>	7.0	17.1	307	<b>6.8</b>	2.7	16.1	57	*	*	*	*	*	*	*	*
\$75,000+	<b>14.0</b>	9.4	20.4	364	<b>0.6</b>	0.1	4.4	56	*	*	*	*	*	*	*	*
<b>EMPLOYMENT</b>																
Employed	<b>18.1</b>	15.4	21.2	1,077	<b>15.4</b>	10.6	21.8	172	<b>14.8</b>	9.3	22.8	137	<b>23.2</b>	15.4	33.3	148
Unemployed	<b>28.5</b>	15.6	46.2	71	*	*	*	*	*	*	*	*	*	*	*	*
Other**	<b>29.6</b>	24.7	35.0	829	<b>13.7</b>	9.0	20.3	119	<b>21.1</b>	14.2	30.3	126	<b>38.8</b>	25.3	54.4	122
<b>EDUCATION</b>																
K-11th Grade	<b>31.4</b>	21.2	43.6	135	*	*	*	*	*	*	*	*	*	*	*	*
12th Grade or GED	<b>26.9</b>	21.8	32.7	555	<b>19.0</b>	12.3	28.0	101	<b>23.7</b>	15.6	34.4	97	<b>35.2</b>	20.9	52.9	94
Some College	<b>28.6</b>	23.5	34.4	652	<b>20.9</b>	13.9	30.3	101	<b>19.9</b>	10.8	33.7	74	<b>32.8</b>	20.9	47.4	94
College Graduate+	<b>11.8</b>	9.1	15.1	636	<b>6.1</b>	2.8	12.6	88	<b>10.6</b>	5.9	18.2	88	<b>6.9</b>	2.7	16.7	65

\*Figure not reliable by BRFSS standards (n<50)

\*\*Other includes students, homemakers, retirees, and persons unable to work

# Cervical Cancer Screening

Idaho females who had not had a Pap test in the last three years, 2008

	Public Health District															
	District 4			District 5			District 6			District 7						
	%	95% CI		n	%	95% CI		n	%	95% CI		n	%	95% CI		n
<b>TOTAL FEMALES</b>	<b>17.6</b>	13.0	23.4	304	<b>22.4</b>	17.3	28.6	279	<b>31.9</b>	23.1	42.2	274	<b>33.7</b>	24.8	44.0	261
<b>AGE</b>																
18-24	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
25-34	<b>14.5</b>	6.9	27.9	54	*	*	*	*	<b>11.8</b>	5.2	24.6	62	<b>14.0</b>	7.3	25.1	71
35-44	<b>11.8</b>	6.3	21.2	81	*	*	*	*	*	*	*	*	<b>14.4</b>	7.4	26.1	54
45-54	<b>12.3</b>	5.8	24.3	64	<b>21.5</b>	12.4	34.7	65	*	*	*	*	<b>15.2</b>	7.7	27.8	53
55-64	<b>20.2</b>	11.6	32.8	54	*	*	*	*	<b>20.5</b>	11.8	33.3	58	*	*	*	*
65+	*	*	*	*	<b>35.0</b>	22.7	49.7	53	*	*	*	*	*	*	*	*
18-34	<b>19.7</b>	11.4	31.8	69	<b>17.1</b>	9.0	30.3	61	<b>31.2</b>	16.1	51.8	76	<b>45.2</b>	30.4	60.8	90
35-54	<b>12.1</b>	7.4	19.0	145	<b>22.6</b>	15.3	31.9	114	<b>30.0</b>	20.3	41.9	91	<b>14.8</b>	9.3	22.8	107
55+	<b>25.6</b>	17.2	36.4	87	<b>31.7</b>	22.8	42.0	102	<b>36.6</b>	27.4	46.8	106	<b>37.6</b>	25.7	51.3	63
<b>INCOME</b>																
Less than \$15,000	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
\$15,000 - \$24,999	*	*	*	*	*	*	*	*	<b>45.0</b>	28.8	62.4	51	*	*	*	*
\$25,000 - \$34,999	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
\$35,000 - \$49,999	<b>21.4</b>	10.4	39.1	50	*	*	*	*	<b>18.9</b>	8.6	36.7	53	*	*	*	*
\$50,000-\$74,999	<b>7.8</b>	2.6	20.9	50	*	*	*	*	*	*	*	*	<b>9.5</b>	4.0	20.6	54
\$75,000+	<b>10.0</b>	4.8	19.7	89	*	*	*	*	*	*	*	*	<b>29.7</b>	12.6	55.2	53
<b>EMPLOYMENT</b>																
Employed	<b>12.2</b>	7.8	18.6	180	<b>19.8</b>	13.9	27.4	160	<b>28.5</b>	18.7	40.9	134	<b>21.7</b>	13.9	32.3	146
Unemployed	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Other**	<b>24.5</b>	16.1	35.4	113	<b>27.0</b>	18.3	37.9	112	<b>36.7</b>	22.2	53.9	129	<b>45.2</b>	30.2	61.0	108
<b>EDUCATION</b>																
K-11th Grade	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
12th Grade or GED	<b>22.4</b>	12.5	36.7	64	<b>27.0</b>	16.6	40.8	65	<b>29.7</b>	17.9	45.2	82	<b>40.4</b>	21.7	62.4	52
Some College	<b>17.0</b>	9.6	28.2	92	<b>25.8</b>	17.0	37.0	105	<b>42.8</b>	25.1	62.6	87	<b>45.2</b>	30.4	60.9	99
College Graduate+	<b>11.7</b>	6.8	19.4	135	<b>12.2</b>	6.8	21.0	78	<b>23.1</b>	12.7	38.2	84	<b>14.4</b>	8.7	22.9	98

\*Figure not reliable by BRFSS standards (n<50)

\*\*Other includes students, homemakers, retirees, and persons unable to work

# Prostate Cancer Screening

Idaho males aged 40 and older who had not had a Prostate Specific Antigen (PSA) test in the last two years, 2008

	Statewide			Public Health District												
				District 1			District 2			District 3						
	%	95% CI		n	%	95% CI		n	%	95% CI		n	%	95% CI		n
<b>TOTAL MALES</b>	<b>47.1</b>	44.0	50.2	1,450	<b>49.7</b>	42.5	56.8	224	<b>48.5</b>	41.5	55.6	233	<b>48.0</b>	39.8	56.2	180
<b>AGE</b>																
40-49	<b>74.5</b>	68.3	79.8	322	*	*	*	*	<b>85.0</b>	72.1	92.6	51	*	*	*	*
50-59	<b>46.6</b>	41.1	52.1	427	<b>49.3</b>	38.0	60.7	82	<b>52.0</b>	39.7	64.0	72	*	*	*	*
60-69	<b>26.3</b>	21.8	31.4	374	<b>27.1</b>	17.3	39.9	58	<b>28.7</b>	17.9	42.6	60	<b>30.4</b>	18.8	45.1	50
70+	<b>23.2</b>	18.5	28.7	327	*	*	*	*	<b>23.8</b>	13.3	39.0	50	*	*	*	*
40-64	<b>54.7</b>	51.0	58.4	947	<b>56.5</b>	48.1	64.6	162	<b>57.5</b>	48.9	65.6	154	<b>56.4</b>	46.4	65.9	115
65+	<b>25.0</b>	21.0	29.4	503	<b>30.4</b>	19.6	44.0	62	<b>29.1</b>	19.2	41.5	79	<b>25.4</b>	15.5	38.7	65
<b>INCOME</b>																
Less than \$15,000	<b>64.1</b>	52.0	74.6	90	*	*	*	*	*	*	*	*	*	*	*	*
\$15,000 - \$24,999	<b>47.1</b>	39.0	55.3	207	*	*	*	*	*	*	*	*	*	*	*	*
\$25,000 - \$34,999	<b>47.3</b>	38.5	56.2	189	*	*	*	*	*	*	*	*	*	*	*	*
\$35,000 - \$49,999	<b>47.9</b>	40.8	55.1	262	*	*	*	*	*	*	*	*	*	*	*	*
\$50,000-\$74,999	<b>46.7</b>	39.7	53.9	254	*	*	*	*	*	*	*	*	*	*	*	*
\$75,000+	<b>44.0</b>	37.8	50.5	340	<b>42.6</b>	28.6	57.8	52	*	*	*	*	*	*	*	*
<b>EMPLOYMENT</b>																
Employed	<b>56.0</b>	52.0	60.0	814	<b>57.2</b>	47.4	66.4	118	<b>55.6</b>	46.4	64.4	131	<b>56.1</b>	44.7	66.9	89
Unemployed	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Other**	<b>27.1</b>	23.2	31.3	593	<b>36.2</b>	26.6	47.0	98	<b>34.9</b>	24.9	46.4	95	<b>30.1</b>	20.4	42.1	83
<b>EDUCATION</b>																
K-11th Grade	<b>54.5</b>	43.0	65.5	98	*	*	*	*	*	*	*	*	*	*	*	*
12th Grade or GED	<b>52.2</b>	46.3	58.0	389	<b>60.6</b>	46.9	72.9	62	<b>60.0</b>	46.5	72.1	66	<b>58.3</b>	44.1	71.2	58
Some College	<b>48.5</b>	42.8	54.2	421	<b>44.9</b>	32.8	57.6	72	<b>51.9</b>	39.5	64.1	71	<b>39.5</b>	25.5	55.5	51
College Graduate+	<b>41.5</b>	36.5	46.7	539	<b>44.8</b>	32.5	57.8	73	<b>35.7</b>	25.3	47.7	80	<b>44.3</b>	29.7	60.0	51

\*Figure not reliable by BRFSS standards (n<50)

\*\*Other includes students, homemakers, retirees, and persons unable to work

# Prostate Cancer Screening

Idaho males aged 40 and older who had not had a Prostate Specific Antigen (PSA) test in the last two years, 2008

	Public Health District															
	District 4			District 5			District 6			District 7						
	%	95% CI		n	%	95% CI		n	%	95% CI		n	%	95% CI		n
<b>TOTAL MALES</b>	<b>41.6</b>	34.2	49.4	205	<b>44.8</b>	37.2	52.7	212	<b>48.9</b>	40.9	56.9	190	<b>54.3</b>	46.7	61.6	206
<b>AGE</b>																
40-49	<b>65.2</b>	50.2	77.7	51	*	*	*	*	*	*	*	*	<b>78.9</b>	65.0	88.2	54
50-59	<b>46.1</b>	32.3	60.5	53	<b>51.3</b>	38.0	64.4	60	<b>43.7</b>	30.5	57.9	60	<b>42.2</b>	28.7	57.0	51
60-69	*	*	*	*	<b>18.2</b>	10.3	30.1	62	<b>22.8</b>	13.5	35.9	52	*	*	*	*
70+	<b>8.9</b>	3.9	19.2	53	<b>16.9</b>	8.5	30.8	52	*	*	*	*	<b>37.7</b>	25.3	52.0	57
40-64	<b>50.1</b>	41.0	59.3	133	<b>54.7</b>	45.0	64.0	124	<b>54.9</b>	45.2	64.1	128	<b>58.8</b>	49.5	67.5	131
65+	<b>13.0</b>	7.2	22.6	72	<b>18.0</b>	11.0	28.1	88	<b>30.9</b>	20.3	44.0	62	<b>40.0</b>	28.9	52.2	75
<b>INCOME</b>																
Less than \$15,000	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
\$15,000 - \$24,999	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
\$25,000 - \$34,999	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
\$35,000 - \$49,999	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
\$50,000-\$74,999	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
\$75,000+	<b>43.3</b>	30.4	57.1	62	*	*	*	*	*	*	*	*	<b>58.7</b>	45.4	70.9	59
<b>EMPLOYMENT</b>																
Employed	<b>51.2</b>	41.2	61.2	113	<b>57.8</b>	48.1	67.0	126	<b>57.9</b>	47.7	67.4	112	<b>61.2</b>	51.8	69.8	125
Unemployed	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Other**	<b>17.3</b>	10.1	28.0	82	<b>10.8</b>	6.0	18.8	83	<b>29.1</b>	19.7	40.7	76	<b>34.8</b>	23.9	47.6	76
<b>EDUCATION</b>																
K-11th Grade	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
12th Grade or GED	*	*	*	*	<b>50.2</b>	36.0	64.3	59	<b>54.5</b>	39.5	68.7	54	<b>48.4</b>	33.8	63.3	50
Some College	*	*	*	*	<b>56.2</b>	41.7	69.7	61	<b>53.2</b>	39.5	66.4	60	<b>59.6</b>	44.7	72.9	60
College Graduate+	<b>41.6</b>	31.8	52.2	109	<b>33.2</b>	22.2	46.3	76	<b>33.1</b>	21.6	46.9	66	<b>51.8</b>	40.4	63.0	84

\*Figure not reliable by BRFSS standards (n<50)

\*\*Other includes students, homemakers, retirees, and persons unable to work

# Dental Visits

Idaho adults who had not visited the dentist in the last 12 months, 2008

	Statewide			Public Health District												
				District 1			District 2			District 3						
	%	95% CI		n	%	95% CI		n	%	95% CI		n	%	95% CI		n
<b>TOTAL</b>	<b>32.0</b>	30.2	33.9	5,091	<b>33.7</b>	29.6	38.1	725	<b>35.9</b>	31.7	40.4	748	<b>38.6</b>	33.9	43.6	748
<b>SEX</b>																
Male	<b>35.2</b>	32.3	38.3	1,978	<b>33.7</b>	27.3	40.7	281	<b>38.6</b>	32.2	45.5	316	<b>42.3</b>	35.0	50.0	266
Female	<b>28.9</b>	26.9	31.0	3,113	<b>33.7</b>	28.8	39.1	444	<b>33.3</b>	28.1	38.9	432	<b>35.0</b>	29.4	41.1	482
<b>AGE</b>																
18-24	<b>35.8</b>	27.5	45.0	162	*	*	*	*	*	*	*	*	*	*	*	*
25-34	<b>33.5</b>	28.9	38.4	628	<b>33.2</b>	22.0	46.7	66	<b>37.4</b>	27.1	49.1	85	<b>36.8</b>	27.5	47.2	108
35-44	<b>29.6</b>	26.0	33.6	773	<b>30.8</b>	22.2	41.0	104	<b>33.8</b>	24.9	43.9	111	<b>35.9</b>	26.5	46.6	103
45-54	<b>31.7</b>	28.4	35.2	1,035	<b>36.5</b>	29.1	44.6	169	<b>43.3</b>	35.0	52.0	153	<b>38.7</b>	30.0	48.2	130
55-64	<b>30.6</b>	27.4	33.9	1,065	<b>32.5</b>	25.4	40.5	161	<b>29.0</b>	21.9	37.3	165	<b>41.0</b>	32.3	50.4	151
65+	<b>32.1</b>	29.4	34.9	1,386	<b>36.7</b>	29.6	44.5	196	<b>37.9</b>	30.9	45.5	211	<b>32.0</b>	25.6	39.2	224
18-34	<b>34.3</b>	30.1	38.8	790	<b>32.3</b>	22.2	44.3	88	<b>34.7</b>	25.6	45.2	105	<b>42.4</b>	32.1	53.4	130
35-54	<b>30.7</b>	28.2	33.3	1,808	<b>33.9</b>	28.1	40.2	273	<b>39.0</b>	32.8	45.5	264	<b>37.3</b>	30.6	44.4	233
55+	<b>31.4</b>	29.3	33.5	2,451	<b>34.7</b>	29.6	40.2	357	<b>34.0</b>	28.9	39.6	376	<b>36.0</b>	30.6	41.8	375
<b>SEX and AGE</b>																
Male																
18-34	<b>40.1</b>	33.3	47.3	288	*	*	*	*	*	*	*	*	<b>49.6</b>	34.1	65.1	52
35-54	<b>32.8</b>	28.9	36.9	725	<b>33.8</b>	25.1	43.8	104	<b>41.8</b>	32.9	51.3	126	<b>37.1</b>	27.0	48.6	87
55+	<b>32.7</b>	29.4	36.2	955	<b>34.5</b>	26.8	43.1	147	<b>33.9</b>	26.1	42.7	146	<b>39.4</b>	30.4	49.2	123
Female																
18-34	<b>28.2</b>	23.7	33.3	502	<b>32.6</b>	20.9	47.0	59	<b>29.0</b>	18.3	42.7	61	<b>34.9</b>	22.5	49.7	78
35-54	<b>28.5</b>	25.5	31.7	1,083	<b>33.9</b>	26.6	42.0	169	<b>36.1</b>	27.8	45.2	138	<b>37.4</b>	29.3	46.2	146
55+	<b>30.2</b>	27.6	32.9	1,496	<b>35.0</b>	28.3	42.2	210	<b>34.1</b>	27.7	41.2	230	<b>33.0</b>	26.9	39.6	252
<b>INCOME</b>																
Less than \$15,000	<b>48.5</b>	41.7	55.4	483	<b>65.1</b>	50.3	77.5	75	<b>38.7</b>	26.6	52.3	86	<b>50.1</b>	35.2	64.9	80
\$15,000 - \$24,999	<b>51.7</b>	46.2	57.1	824	<b>47.7</b>	36.5	59.2	119	<b>53.9</b>	42.7	64.7	131	<b>56.6</b>	45.4	67.2	149
\$25,000 - \$34,999	<b>41.5</b>	35.7	47.5	660	<b>41.3</b>	28.2	55.8	78	<b>41.9</b>	31.3	53.3	104	<b>39.5</b>	27.7	52.7	109
\$35,000 - \$49,999	<b>30.0</b>	26.1	34.2	852	<b>25.7</b>	17.9	35.4	116	<b>32.9</b>	23.6	43.9	128	<b>33.8</b>	25.0	44.0	130
\$50,000-\$74,999	<b>21.3</b>	17.7	25.3	791	<b>24.1</b>	16.3	34.0	114	<b>20.9</b>	13.7	30.4	109	<b>30.2</b>	20.0	42.8	105
\$75,000+	<b>15.4</b>	12.6	18.8	910	<b>19.2</b>	12.1	29.0	135	<b>20.9</b>	13.4	31.1	111	<b>20.0</b>	10.9	34.0	86
<b>EMPLOYMENT</b>																
Employed	<b>32.1</b>	29.7	34.7	2,719	<b>31.6</b>	26.0	37.7	378	<b>38.6</b>	32.9	44.7	388	<b>38.8</b>	32.8	45.1	381
Unemployed	<b>45.3</b>	35.1	55.9	176	*	*	*	*	*	*	*	*	*	*	*	*
Other**	<b>30.0</b>	27.5	32.7	2,188	<b>35.1</b>	29.3	41.3	317	<b>31.0</b>	25.5	37.1	329	<b>35.0</b>	28.1	42.6	335
<b>EDUCATION</b>																
K-11th Grade	<b>54.8</b>	47.8	61.7	401	<b>61.4</b>	42.9	77.2	52	<b>26.9</b>	16.2	41.2	53	<b>50.7</b>	36.9	64.4	92
12th Grade or GED	<b>37.9</b>	34.3	41.7	1,511	<b>36.3</b>	29.2	44.0	231	<b>45.5</b>	37.7	53.5	248	<b>42.5</b>	34.1	51.5	256
Some College	<b>33.1</b>	29.9	36.4	1,622	<b>35.3</b>	28.5	42.8	244	<b>39.7</b>	32.2	47.7	226	<b>38.5</b>	30.2	47.6	228
College Graduate+	<b>17.9</b>	15.6	20.5	1,546	<b>19.5</b>	14.0	26.3	195	<b>23.8</b>	17.9	31.0	220	<b>23.5</b>	17.1	31.4	170

\*Figure not reliable by BRFSS standards (n<50)

\*\*Other includes students, homemakers, retirees, and persons unable to work

# Dental Visits

Idaho adults who had not visited the dentist in the last 12 months, 2008

	Public Health District															
	District 4				District 5				District 6				District 7			
	%	95% CI		n	%	95% CI		n	%	95% CI		n	%	95% CI		n
<b>TOTAL</b>	<b>26.8</b>	22.7	31.3	750	<b>36.2</b>	31.8	40.8	726	<b>30.4</b>	25.9	35.3	699	<b>28.8</b>	24.3	33.8	695
<b>SEX</b>																
Male	<b>33.5</b>	26.6	41.1	293	<b>36.5</b>	29.4	44.2	283	<b>33.4</b>	26.2	41.4	253	<b>30.5</b>	23.5	38.6	286
Female	<b>19.9</b>	16.0	24.5	457	<b>35.9</b>	31.0	41.2	443	<b>27.5</b>	22.4	33.1	446	<b>27.1</b>	21.7	33.2	409
<b>AGE</b>																
18-24	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
25-34	<b>25.9</b>	16.5	38.1	87	<b>47.0</b>	34.7	59.6	78	<b>32.8</b>	22.5	44.9	93	<b>33.0</b>	23.1	44.6	111
35-44	<b>25.1</b>	17.9	33.9	149	<b>32.9</b>	23.9	43.4	107	<b>33.2</b>	23.5	44.5	88	<b>23.1</b>	16.1	32.0	111
45-54	<b>30.8</b>	23.2	39.6	150	<b>24.0</b>	17.5	31.9	153	<b>28.0</b>	20.2	37.4	140	<b>23.1</b>	16.4	31.5	140
55-64	<b>24.2</b>	17.6	32.3	163	<b>33.6</b>	25.8	42.4	141	<b>28.1</b>	21.4	35.9	162	<b>29.8</b>	21.6	39.6	122
65+	<b>19.9</b>	14.3	26.9	171	<b>43.8</b>	36.9	50.9	219	<b>33.9</b>	27.0	41.7	187	<b>29.9</b>	23.1	37.7	178
18-34	<b>29.2</b>	20.1	40.5	110	<b>43.4</b>	32.6	54.8	99	<b>29.7</b>	20.2	41.5	117	<b>33.1</b>	23.3	44.6	141
35-54	<b>27.8</b>	22.5	33.9	299	<b>28.2</b>	22.6	34.6	260	<b>30.4</b>	24.1	37.6	228	<b>23.1</b>	18.1	29.1	251
55+	<b>22.1</b>	17.6	27.4	334	<b>39.3</b>	34.0	44.8	360	<b>31.2</b>	26.2	36.6	349	<b>29.9</b>	24.4	36.0	300
<b>SEX and AGE</b>																
Male																
18-34	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
35-54	<b>33.6</b>	25.1	43.2	118	<b>24.7</b>	16.7	34.8	95	<b>35.8</b>	25.6	47.4	87	<b>23.8</b>	16.5	33.1	108
55+	<b>25.5</b>	18.3	34.3	136	<b>41.5</b>	33.4	50.1	150	<b>25.8</b>	18.5	34.6	125	<b>32.5</b>	24.1	42.3	128
Female																
18-34	<b>18.1</b>	10.5	29.5	72	<b>40.6</b>	28.9	53.4	63	<b>22.6</b>	13.1	36.3	77	<b>30.9</b>	19.8	44.8	92
35-54	<b>21.6</b>	15.7	28.9	181	<b>31.8</b>	24.5	40.1	165	<b>25.1</b>	18.2	33.6	141	<b>22.4</b>	16.1	30.4	143
55+	<b>19.1</b>	14.0	25.5	198	<b>37.4</b>	30.7	44.5	210	<b>36.0</b>	29.6	42.9	224	<b>27.4</b>	20.8	35.2	172
<b>INCOME</b>																
Less than \$15,000	<b>56.7</b>	38.3	73.4	55	<b>53.0</b>	37.7	67.8	67	<b>34.4</b>	20.2	52.1	68	<b>35.1</b>	18.5	56.4	52
\$15,000 - \$24,999	<b>56.1</b>	40.5	70.6	81	<b>59.8</b>	47.0	71.5	123	<b>38.4</b>	27.5	50.7	112	<b>42.0</b>	28.3	57.1	109
\$25,000 - \$34,999	<b>36.1</b>	22.6	52.1	88	<b>45.0</b>	33.8	56.8	100	<b>45.5</b>	30.6	61.2	101	<b>49.7</b>	34.1	65.4	80
\$35,000 - \$49,999	<b>30.6</b>	21.4	41.7	133	<b>32.8</b>	23.1	44.2	113	<b>32.3</b>	22.7	43.8	120	<b>21.6</b>	14.3	31.2	112
\$50,000-\$74,999	<b>16.8</b>	9.6	27.6	116	<b>26.1</b>	17.2	37.5	109	<b>18.8</b>	11.7	28.7	110	<b>15.5</b>	10.0	23.3	128
\$75,000+	<b>11.7</b>	7.2	18.5	200	<b>14.4</b>	8.6	23.1	119	<b>16.6</b>	9.1	28.4	118	<b>16.2</b>	10.2	24.8	141
<b>EMPLOYMENT</b>																
Employed	<b>27.1</b>	21.7	33.3	416	<b>34.0</b>	28.2	40.4	410	<b>31.7</b>	25.5	38.6	356	<b>31.4</b>	25.5	38.0	390
Unemployed	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Other**	<b>23.3</b>	17.5	30.4	300	<b>39.6</b>	33.3	46.3	297	<b>28.6</b>	22.7	35.3	325	<b>23.4</b>	17.0	31.4	285
<b>EDUCATION</b>																
K-11th Grade	*	*	*	*	<b>61.2</b>	44.8	75.4	70	<b>32.0</b>	17.0	52.0	51	*	*	*	*
12th Grade or GED	<b>31.9</b>	22.6	42.9	173	<b>44.1</b>	35.6	53.0	208	<b>35.1</b>	26.8	44.5	216	<b>36.9</b>	27.8	47.0	179
Some College	<b>31.2</b>	23.7	39.9	220	<b>32.6</b>	26.4	39.6	247	<b>32.8</b>	24.8	41.9	228	<b>25.1</b>	18.2	33.6	229
College Graduate+	<b>13.5</b>	9.6	18.8	316	<b>19.3</b>	13.2	27.4	199	<b>21.3</b>	15.1	29.2	202	<b>17.6</b>	12.2	24.7	244

\*Figure not reliable by BRFSS standards (n<50)

\*\*Other includes students, homemakers, retirees, and persons unable to work



## Appendix A

### Trends of Selected Risk Variables by Public Health District From 2001 Through 2008

The following tables present data for each health district for selected risk variables from the years 2001 through 2008. The following table will help with interpretation of the variable headings.

<b>Variable Heading</b>	<b>Definition</b>	<b>Question/Page in Appendix B</b>
General Health	Adults who reported their general health status as "fair" or "poor"	Q1.1 p. 96
Health Care Coverage	Adults without health care coverage	Q3.1 p. 96
Dental Insurance	Adults without dental insurance	ID2.4 p. 106
Diabetes	Adults who had ever been told they had diabetes	Q6.1 p. 96
Asthma	Adults who were diagnosed and still had asthma	Q9.1-Q9.2 p. 98
Physical Activity	Adults who did not participate in leisure time physical activity	Q5.1 p. 96
Overweight	Adults who were overweight (BMI $\geq$ 25)	Q12.11, Q12.12 p. 99
Obesity	Adults who were obese (BMI $\geq$ 30)	
Cigarette Smoking	Adults who had smoked 100 or cigarettes in their lifetimes and who currently smoke every day or some days.	Q11.1-Q11.2 p. 98
Binge Drinking	Males who consumed 5+ drinks and females who consumed 4+ drinks on an occasion	Q13.1-Q13.5 p. 100
Heavy Drinking	Males who consumed 60+ drinks and females who consumed 30+ drinks in the last 30 days	Q13.1-Q13.3 p. 100
Illicit Drug Use	Adults who used illicit drugs in the last 12 months	ID8.1-ID8.2 p. 107
Seat Belt Use	Adults who did not always wear a seat belt	Q16.1 p. 101
Colorectal Cancer Screening	Adults aged 50 and older who had never had a sigmoidoscopy or colonoscopy.	Q20.1-Q20.3 p. 102
Breast Cancer Screening	Females aged 40 and older who had not had a mammogram and clinical breast exam in the last two years	Q18.1-Q18.4 p. 101
Cervical Cancer Screening	Females who had not had a Pap smear in the last three years	Q18.5-Q18.6 p. 101
Prostate Cancer Screening	Males aged 40 and older who had not had a PSA test in the last two years	Q19.1-Q19.2 p. 102
Dental Visits	Adults who had not visited the dentist in the last 12 months	Q7.1 p. 97

# Trends of Selected Risk Factors

## By Public Health District, 2001-2008

	Year													
	2001			2002			2003			2004				
	%	95% CI	n											
<b>Idaho</b>														
General Health	13.0	12.0 14.1	4,822	13.4	12.4 14.6	4,850	13.6	12.6 14.8	4,813	12.5	11.5 13.6	5,060		
Health Care Coverage	14.8	13.6 16.1	4,819	16.8	15.6 18.2	4,845	16.3	15.0 17.7	4,808	15.5	14.3 16.8	5,051		
Dental Insurance	44.9	43.2 46.7	4,693	45.0	43.3 46.7	4,791	44.6	42.9 46.3	4,771	43.5	41.8 45.2	5,013		
Diabetes	5.4	4.8 6.2	4,830	6.1	5.4 6.8	4,853	6.3	5.5 7.1	4,823	6.2	5.5 6.9	5,070		
Asthma <sup>1</sup>	8.0	7.1 8.9	4,821	7.8	6.9 8.7	4,836	8.0	7.1 9.0	4,800	7.9	7.0 8.9	5,041		
Physical Activity	21.0	19.6 22.5	4,831	19.4	18.1 20.7	4,855	18.3	17.1 19.7	4,817	18.8	17.5 20.1	5,061		
Overweight <sup>2</sup>	59.3	57.6 61.1	4,588	57.3	55.6 59.1	4,661	59.3	57.5 61.1	4,583	58.2	56.4 60.0	4,889		
Obesity	20.5	19.2 21.9	4,588	20.4	19.1 21.8	4,661	22.0	20.5 23.5	4,583	21.1	19.7 22.5	4,889		
Cigarette Smoking	19.6	18.3 21.0	4,823	20.7	19.3 22.2	4,846	18.9	17.6 20.3	4,813	17.4	16.2 18.8	5,054		
Binge Drinking <sup>3,4</sup>	*	* *	*	*	* *	*	*	* *	*	*	* *	*		
Heavy Drinking <sup>3,4</sup>	*	* *	*	*	* *	*	*	* *	*	*	* *	*		
Illicit Drug Use <sup>5</sup>	*	* *	*	5.8	4.9 6.9	4,830	6.5	5.6 7.5	4,802	5.8	4.9 6.8	5,046		
Seat Belt Use	34.8	33.2 36.4	4,713	34.7	33.1 36.4	4,847	32.7	31.1 34.4	4,812	*	* *	*		
Colorectal Cancer Screening	56.6	53.8 59.3	1,931	53.7	51.2 56.1	2,165	*	* *	*	52.5	50.2 54.9	2,435		
Breast Cancer Screening <sup>6</sup>	*	* *	*	37.6	35.1 40.2	1,848	*	* *	*	40.7	38.2 43.2	2,019		
Cervical Cancer Screening <sup>2</sup>	*	* *	*	16.2	14.3 18.2	2,026	*	* *	*	21.0	18.9 23.2	2,041		
Prostate Cancer Screening	52.1	48.6 55.5	1,209	50.3	47.0 53.5	1,237	*	* *	*	52.0	48.8 55.2	1,305		
Dental Visits	34.6	33.0 36.2	4,714	32.5	30.9 34.1	4,836	33.8	32.2 35.5	4,795	33.4	31.8 35.0	5,025		
<b>District 1</b>														
General Health	14.3	11.8 17.3	690	17.4	14.7 20.6	721	16.8	13.9 20.1	681	13.1	10.7 15.9	706		
Health Care Coverage	15.9	13.1 19.3	687	20.7	17.4 24.5	722	20.2	17.0 23.9	679	19.6	16.3 23.5	702		
Dental Insurance	50.7	46.5 54.9	667	53.5	49.3 57.5	714	50.9	46.7 55.1	678	51.8	47.6 56.0	697		
Diabetes	5.4	3.9 7.5	690	6.8	5.1 9.0	723	7.0	5.2 9.4	683	4.6	3.3 6.5	707		
Asthma <sup>1</sup>	7.7	5.8 10.1	688	10.2	8.0 12.9	721	8.3	6.4 10.7	676	7.9	6.0 10.4	701		
Physical Activity	20.8	17.7 24.3	690	19.5	16.5 22.9	722	19.0	15.9 22.5	683	19.1	16.0 22.6	705		
Overweight <sup>2</sup>	58.1	53.9 62.2	658	59.0	54.9 63.1	695	59.6	55.3 63.7	649	60.0	55.8 64.1	678		
Obesity	17.6	14.6 21.1	658	22.8	19.5 26.4	695	20.4	17.2 24.0	649	21.5	18.2 25.2	678		
Cigarette Smoking	20.6	17.4 24.1	688	27.0	23.3 31.0	721	21.7	18.5 25.3	682	20.9	17.7 24.6	703		
Binge Drinking <sup>3,4</sup>	*	* *	*	*	* *	*	*	* *	*	*	* *	*		
Heavy Drinking <sup>3,4</sup>	*	* *	*	*	* *	*	*	* *	*	*	* *	*		
Illicit Drug Use <sup>5</sup>	*	* *	*	7.3	5.2 10.3	718	6.7	4.8 9.2	677	5.2	3.5 7.6	701		
Seat Belt Use	32.1	28.3 36.2	674	28.5	24.8 32.4	720	30.0	26.2 34.1	683	*	* *	*		
Colorectal Cancer Screening	57.8	51.7 63.7	300	58.7	53.2 64.1	360	*	* *	*	55.3	49.6 60.8	376		
Breast Cancer Screening <sup>6</sup>	*	* *	*	35.3	29.6 41.6	288	*	* *	*	46.6	40.5 52.9	302		
Cervical Cancer Screening <sup>2</sup>	*	* *	*	14.8	10.8 19.9	302	*	* *	*	24.7	19.6 30.6	305		
Prostate Cancer Screening	55.8	47.8 63.5	174	54.5	47.1 61.8	205	*	* *	*	56.5	48.6 64.0	186		
Dental Visits	40.4	36.4 44.6	673	37.5	33.6 41.6	719	37.9	33.9 42.0	681	37.5	33.6 41.7	696		

\* Indicates data are not available for that year

1 Prior to 2002, Idaho BRFSS reported adults who have ever been told they have asthma. Beginning in 2002, Idaho reports current asthma status.

2 Results may differ from previous publications due to revisions in risk factor calculation.

3 Question wording changes make data prior to 2005 not comparable with 2005 and later.

4 Prior to 2005, Idaho BRFSS reported drinking status among all adults. Beginning in 2005, Idaho reports gender specific drinking status.

5 Question wording changes make data prior to 2002 not comparable with 2002 and later.

6 Prior to 2004, Idaho BRFSS reported all ages. Beginning in 2004, Idaho reports only women aged 40 and older.

# Trends of Selected Risk Factors

## By Public Health District, 2001-2008

	Year															
	2005				2006				2007				2008			
	%	95% CI	n	%	95% CI	n	%	95% CI	n	%	95% CI	n				
<b>Idaho</b>																
General Health	14.9	13.7	16.2	5,724	13.9	12.8	15.1	5,337	14.9	13.8	16.2	5,303	14.0	12.9	15.2	5,099
Health Care Coverage	18.9	17.3	20.6	5,716	18.8	17.2	20.4	5,326	18.9	17.3	20.6	5,299	18.5	16.9	20.4	5,099
Dental Insurance	45.7	43.9	47.5	5,390	45.7	43.9	47.5	5,081	43.0	41.2	44.9	4,992	43.0	41.0	44.9	4,814
Diabetes	6.8	6.2	7.6	5,726	6.8	6.1	7.7	5,339	7.9	7.1	8.7	5,306	7.0	6.3	7.8	5,108
Asthma <sup>1</sup>	7.3	6.5	8.2	5,697	9.2	8.2	10.3	5,295	8.7	7.7	9.8	5,273	8.9	7.8	10.1	5,079
Physical Activity	21.6	20.2	23.1	5,722	20.8	19.4	22.2	5,331	19.6	18.2	21.0	5,304	21.1	19.7	22.7	5,104
Overweight <sup>2</sup>	61.4	59.5	63.1	5,439	59.7	57.9	61.6	5,091	63.1	61.2	64.9	5,042	62.2	60.2	64.2	4,851
Obesity	24.5	23.1	26.1	5,439	24.1	22.7	25.6	5,091	25.1	23.6	26.7	5,042	25.2	23.6	26.8	4,851
Cigarette Smoking	17.9	16.6	19.4	5,719	16.8	15.5	18.2	5,328	19.1	17.6	20.7	5,302	16.9	15.4	18.4	5,095
Binge Drinking <sup>3,4</sup>	14.2	12.7	15.8	5,683	14.8	13.4	16.3	5,258	14.7	13.3	16.3	5,227	13.1	11.8	14.6	5,028
Heavy Drinking <sup>3,4</sup>	4.2	3.6	5.0	5,668	4.9	4.0	5.9	5,223	5.0	4.1	5.9	5,180	4.6	3.9	5.6	5,010
Illicit Drug Use <sup>5</sup>	5.3	4.3	6.5	5,332	4.1	3.2	5.1	5,084	5.1	4.1	6.5	4,945	4.1	3.2	5.2	4,797
Seat Belt Use	23.8	22.3	25.4	5,422	26.8	25.2	28.4	5,245	24.6	22.9	26.4	5,026	23.4	21.8	25.1	5,039
Colorectal Cancer Screening	*	*	*	*	45.8	43.6	48.0	2,686	*	*	*	*	44.4	42.3	46.6	2,946
Breast Cancer Screening <sup>6</sup>	*	*	*	*	37.8	35.4	40.2	2,182	*	*	*	*	36.8	34.5	39.2	2,287
Cervical Cancer Screening <sup>2</sup>	*	*	*	*	22.5	20.2	24.9	2,143	*	*	*	*	23.1	20.5	26.1	1,980
Prostate Cancer Screening	*	*	*	*	48.7	45.5	51.8	1,405	*	*	*	*	47.1	44.0	50.2	1,450
Dental Visits	33.5	31.9	35.3	5,481	34.6	32.9	36.3	5,321	32.3	30.6	34.1	5,018	32.0	30.2	33.9	5,091
<b>District 1</b>																
General Health	13.4	10.5	16.8	902	15.9	13.2	19.1	748	13.1	10.7	16.0	769	16.5	13.5	19.9	728
Health Care Coverage	21.2	17.1	25.8	902	20.6	16.8	24.9	750	18.4	15.0	22.4	767	20.5	16.7	25.0	727
Dental Insurance	51.9	47.5	56.3	857	46.9	42.4	51.5	718	49.2	44.6	53.8	715	48.6	43.9	53.3	685
Diabetes	6.3	4.9	8.0	901	7.0	5.4	9.2	750	7.0	5.3	9.2	770	6.4	4.8	8.4	729
Asthma <sup>1</sup>	7.4	5.6	9.8	894	11.0	8.0	14.9	739	8.4	6.4	11.1	760	9.5	7.2	12.5	724
Physical Activity	20.8	17.7	24.4	900	16.8	13.8	20.3	749	18.1	15.2	21.3	769	24.5	20.9	28.6	729
Overweight <sup>2</sup>	59.4	54.9	63.7	858	62.7	58.1	67.1	723	60.8	56.3	65.2	735	63.3	58.6	67.8	685
Obesity	21.7	18.5	25.2	858	23.7	20.3	27.6	723	23.7	20.2	27.5	735	24.3	20.7	28.3	685
Cigarette Smoking	20.7	17.0	25.0	900	19.2	15.8	23.1	747	23.6	19.9	27.8	765	20.5	16.9	24.7	729
Binge Drinking <sup>3,4</sup>	13.2	10.5	16.4	893	17.4	13.9	21.5	740	17.4	14.1	21.4	753	15.8	12.2	20.1	713
Heavy Drinking <sup>3,4</sup>	5.2	3.8	7.1	887	6.7	4.8	9.1	731	5.1	3.5	7.4	744	6.8	4.9	9.4	710
Illicit Drug Use <sup>5</sup>	6.6	4.0	10.7	845	4.6	2.8	7.7	717	4.7	2.8	7.9	705	4.8	2.9	7.8	679
Seat Belt Use	17.5	14.7	20.8	864	22.2	18.5	26.3	737	22.2	18.3	26.7	717	22.6	18.8	26.9	720
Colorectal Cancer Screening	*	*	*	*	45.6	40.3	51.0	423	*	*	*	*	45.9	40.8	51.1	437
Breast Cancer Screening <sup>6</sup>	*	*	*	*	38.2	32.5	44.3	320	*	*	*	*	39.4	33.8	45.4	326
Cervical Cancer Screening <sup>2</sup>	*	*	*	*	20.9	15.4	27.8	294	*	*	*	*	16.2	12.3	21.0	307
Prostate Cancer Screening	*	*	*	*	49.6	42.4	56.8	225	*	*	*	*	49.7	42.5	56.8	224
Dental Visits	38.3	33.9	42.8	869	37.8	33.6	42.2	747	36.7	32.4	41.2	716	33.7	29.6	38.1	725

1 Prior to 2002, Idaho BRFSS reported adults who have ever been told they have asthma. Beginning in 2002, Idaho reports current asthma.

2 Results may differ from previous publications due to revisions in risk factor calculation.

3 Question wording changes make data prior to 2005 not comparable with 2005 and later.

4 Prior to 2005, Idaho BRFSS reported drinking status among all adults. Beginning in 2005, Idaho reports gender specific drinking status.

5 Question wording changes make data prior to 2002 not comparable with 2002 and later.

6 Prior to 2004, Idaho BRFSS reported all ages. Beginning in 2004, Idaho reports only women aged 40 and older.

# Trends of Selected Risk Factors

## By Public Health District, 2001-2008

	Year											
	2001			2002			2003			2004		
	%	95% CI	n									
<b>District 2</b>												
General Health	11.7	9.3 14.6	681	15.2	12.1 18.8	705	15.0	12.3 18.1	656	16.8	13.9 20.1	737
Health Care Coverage	14.4	11.5 17.9	684	16.1	13.2 19.4	704	15.9	12.8 19.5	658	15.9	12.9 19.4	736
Dental Insurance	45.4	41.1 49.7	663	46.5	42.5 50.6	698	49.5	45.1 53.9	652	49.8	45.7 54.0	729
Diabetes	4.2	2.9 6.0	683	6.3	4.6 8.4	705	5.8	4.2 8.0	658	6.7	4.9 9.1	737
Asthma <sup>1</sup>	8.3	6.2 11.0	681	9.2	6.6 12.7	699	7.3	5.4 9.8	655	7.7	5.8 10.1	732
Physical Activity	22.5	19.3 26.2	684	16.8	14.0 20.0	706	16.7	13.9 20.1	657	16.3	13.7 19.4	734
Overweight <sup>2</sup>	58.6	54.3 62.9	644	58.8	54.4 63.0	678	59.4	54.9 63.8	625	59.1	54.8 63.2	710
Obesity	21.6	18.2 25.4	644	19.1	16.2 22.5	678	20.8	17.6 24.5	625	20.9	17.8 24.3	710
Cigarette Smoking	18.2	15.1 21.8	683	20.1	17.0 23.7	704	17.5	14.4 21.1	658	18.6	15.7 21.9	737
Binge Drinking <sup>3,4</sup>	*	* *	*	*	* *	*	*	* *	*	*	* *	*
Heavy Drinking <sup>3,4</sup>	*	* *	*	*	* *	*	*	* *	*	*	* *	*
Illicit Drug Use <sup>5</sup>	*	* *	*	6.0	4.3 8.4	698	7.3	5.1 10.4	655	5.5	3.7 8.2	733
Seat Belt Use	38.3	34.3 42.6	662	34.5	30.7 38.6	705	30.7	26.7 35.1	655	*	* *	*
Colorectal Cancer Screening	50.8	44.4 57.2	276	49.1	43.0 55.2	316	*	* *	*	46.1	40.3 51.9	353
Breast Cancer Screening <sup>6</sup>	*	* *	*	38.8	32.6 45.4	265	*	* *	*	39.4	33.5 45.6	289
Cervical Cancer Screening <sup>2</sup>	*	* *	*	15.9	11.6 21.3	281	*	* *	*	21.8	16.9 27.7	293
Prostate Cancer Screening	48.9	41.3 56.5	184	48.4	40.6 56.3	182	*	* *	*	50.7	43.1 58.3	209
Dental Visits	37.0	32.9 41.3	667	35.8	31.8 40.1	699	37.4	33.2 41.8	655	36.4	32.5 40.5	731
<b>District 3</b>												
General Health	16.2	13.4 19.4	681	14.0	11.4 17.0	688	16.8	13.9 20.2	677	15.7	13.0 18.7	708
Health Care Coverage	16.0	13.1 19.3	679	17.3	14.3 20.7	686	22.6	19.0 26.7	677	18.9	15.6 22.6	708
Dental Insurance	48.0	43.8 52.3	661	45.9	41.6 50.2	670	48.8	44.5 53.1	674	46.7	42.5 51.0	704
Diabetes	6.1	4.3 8.4	681	7.5	5.6 10.0	689	7.7	5.8 10.3	679	7.2	5.5 9.5	710
Asthma <sup>1</sup>	7.9	6.0 10.4	680	6.3	4.6 8.7	687	7.9	5.9 10.6	678	9.5	7.4 12.2	707
Physical Activity	23.0	19.9 26.6	681	22.0	18.8 25.7	689	20.6	17.4 24.2	676	27.3	23.4 31.5	710
Overweight <sup>2</sup>	63.1	58.9 67.1	652	60.6	56.1 64.9	655	61.5	57.1 65.8	637	60.1	55.7 64.4	682
Obesity	24.6	21.2 28.4	652	22.8	19.4 26.5	655	24.4	21.0 28.3	637	24.0	20.6 27.8	682
Cigarette Smoking	24.0	20.6 27.9	680	20.2	17.0 23.7	687	21.8	18.5 25.5	676	19.7	16.6 23.2	707
Binge Drinking <sup>3,4</sup>	*	* *	*	*	* *	*	*	* *	*	*	* *	*
Heavy Drinking <sup>3,4</sup>	*	* *	*	*	* *	*	*	* *	*	*	* *	*
Illicit Drug Use <sup>5</sup>	*	* *	*	3.0	1.7 5.0	681	6.3	4.3 9.1	676	5.5	3.7 8.1	708
Seat Belt Use	35.6	31.8 39.7	662	35.8	31.8 40.1	689	31.7	27.7 35.9	677	*	* *	*
Colorectal Cancer Screening	58.1	51.6 64.3	263	58.6	52.2 64.7	308	*	* *	*	56.3	50.5 62.0	338
Breast Cancer Screening <sup>6</sup>	*	* *	*	42.9	36.3 49.8	260	*	* *	*	42.3	36.0 48.9	280
Cervical Cancer Screening <sup>2</sup>	*	* *	*	16.8	12.2 22.6	278	*	* *	*	16.5	11.5 23.0	264
Prostate Cancer Screening	56.9	48.5 65.0	155	50.4	41.9 58.8	164	*	* *	*	55.9	47.9 63.7	168
Dental Visits	40.9	36.9 45.1	665	37.4	33.3 41.7	686	41.5	37.3 45.8	673	38.6	34.5 42.9	707

\* Indicates data are not available for that year

1 Prior to 2002, Idaho BRFSS reported adults who have ever been told they have asthma. Beginning in 2002, Idaho reports current asthma status.

2 Results may differ from previous publications due to revisions in risk factor calculation.

3 Question wording changes make data prior to 2005 not comparable with 2005 and later.

4 Prior to 2005, Idaho BRFSS reported drinking status among all adults. Beginning in 2005, Idaho reports gender specific drinking status.

5 Question wording changes make data prior to 2002 not comparable with 2002 and later.

6 Prior to 2004, Idaho BRFSS reported all ages. Beginning in 2004, Idaho reports only women aged 40 and older.

# Trends of Selected Risk Factors

## By Public Health District, 2001-2008

	Year											
	2005			2006			2007			2008		
	%	95% CI	n									
<b>District 2</b>												
General Health	16.7	14.0 19.8	791	15.9	13.0 19.3	746	17.9	14.5 21.7	776	16.5	13.7 19.8	751
Health Care Coverage	22.5	18.6 26.9	785	21.8	17.1 27.3	743	15.5	12.0 19.9	776	16.2	12.8 20.2	750
Dental Insurance	54.6	50.2 58.9	744	52.5	47.7 57.3	709	44.5	39.6 49.4	732	46.1	41.5 50.7	708
Diabetes	6.9	5.1 9.1	791	6.8	5.2 8.9	746	8.2	6.3 10.4	777	7.9	6.1 10.1	754
Asthma <sup>1</sup>	7.4	5.4 10.0	785	7.0	5.3 9.3	742	8.0	5.9 10.9	772	11.5	9.0 14.6	752
Physical Activity	23.9	20.2 28.0	791	22.2	18.5 26.3	745	20.7	17.3 24.6	777	21.2	17.9 25.0	752
Overweight <sup>2</sup>	61.1	56.8 65.2	744	61.7	56.5 66.7	717	64.6	59.5 69.4	741	63.5	58.8 67.9	717
Obesity	24.9	21.4 28.7	744	23.1	19.6 27.0	717	27.1	23.1 31.4	741	28.1	24.3 32.2	717
Cigarette Smoking	19.3	15.7 23.4	785	21.8	17.3 27.0	745	19.4	15.5 23.9	778	18.3	15.0 22.0	748
Binge Drinking <sup>3,4</sup>	13.6	10.8 17.0	783	19.6	15.2 24.9	727	15.8	12.1 20.3	766	17.5	14.1 21.5	741
Heavy Drinking <sup>3,4</sup>	4.5	3.1 6.5	786	5.3	3.4 8.2	719	4.6	3.0 6.9	752	5.5	3.8 7.9	739
Illicit Drug Use <sup>5</sup>	4.4	2.7 7.0	737	4.0	2.0 7.8	717	6.0	3.4 10.5	729	2.7	1.5 5.0	709
Seat Belt Use	27.3	23.5 31.5	750	25.9	21.5 30.8	730	23.6	19.6 28.0	737	24.3	20.5 28.5	746
Colorectal Cancer Screening	*	*	*	39.6	34.5 44.9	398	*	*	*	38.4	33.4 43.6	450
Breast Cancer Screening <sup>6</sup>	*	*	*	36.0	30.2 42.1	298	*	*	*	39.2	33.6 45.1	339
Cervical Cancer Screening <sup>2</sup>	*	*	*	25.3	19.1 32.8	287	*	*	*	17.7	13.2 23.3	274
Prostate Cancer Screening	*	*	*	53.0	46.0 59.8	233	*	*	*	48.5	41.5 55.6	233
Dental Visits	37.3	33.0 41.8	754	39.6	35.0 44.4	741	41.0	36.2 46.0	737	35.9	31.7 40.4	748
<b>District 3</b>												
General Health	17.9	14.8 21.5	783	18.4	15.6 21.7	788	16.0	13.3 19.3	747	16.5	13.5 20.1	749
Health Care Coverage	25.7	21.5 30.4	785	23.5	19.9 27.5	783	24.1	19.9 29.0	745	25.4	20.9 30.5	750
Dental Insurance	48.4	43.9 52.9	735	46.0	41.8 50.2	751	46.4	41.7 51.2	709	48.5	43.5 53.5	702
Diabetes	8.4	6.5 10.8	785	7.9	6.1 10.2	788	9.3	7.3 11.8	746	7.1	5.5 9.1	749
Asthma <sup>1</sup>	8.1	6.2 10.5	780	8.3	6.3 10.8	786	8.5	6.5 11.0	744	10.3	7.7 13.6	743
Physical Activity	27.7	23.9 31.8	785	25.0	21.7 28.6	786	23.2	19.7 27.2	748	25.6	21.9 29.7	748
Overweight <sup>2</sup>	68.0	63.5 72.2	749	61.1	56.8 65.4	739	67.1	62.3 71.6	713	67.2	62.3 71.7	712
Obesity	29.9	26.1 34.0	749	27.5	24.0 31.3	739	29.5	25.6 33.7	713	31.0	26.5 35.9	712
Cigarette Smoking	21.8	18.0 26.2	786	20.9	17.7 24.5	785	20.6	16.8 25.0	749	17.7	14.3 21.7	748
Binge Drinking <sup>3,4</sup>	13.3	10.4 17.0	777	10.4	8.1 13.2	775	13.4	10.0 17.8	734	11.2	8.4 14.7	735
Heavy Drinking <sup>3,4</sup>	3.4	2.1 5.5	775	3.1	1.9 5.0	764	4.9	3.4 7.0	731	3.8	2.5 5.8	736
Illicit Drug Use <sup>5</sup>	6.7	3.9 11.2	728	3.6	2.2 5.9	754	3.8	1.9 7.3	702	3.2	1.7 5.7	699
Seat Belt Use	21.4	18.0 25.3	739	23.9	20.4 27.8	772	22.3	18.6 26.6	716	20.0	16.1 24.7	739
Colorectal Cancer Screening	*	*	*	50.5	45.0 56.0	370	*	*	*	47.7	42.3 53.2	428
Breast Cancer Screening <sup>6</sup>	*	*	*	39.5	33.5 45.8	301	*	*	*	42.7	36.8 48.8	351
Cervical Cancer Screening <sup>2</sup>	*	*	*	23.7	18.2 30.3	308	*	*	*	28.9	21.3 38.1	281
Prostate Cancer Screening	*	*	*	46.8	39.3 54.5	193	*	*	*	48.0	39.8 56.2	180
Dental Visits	39.9	35.5 44.5	753	41.7	37.6 45.9	786	32.7	28.6 37.1	715	38.6	33.9 43.5	748

1 Prior to 2002, Idaho BRFSS reported adults who have ever been told they have asthma. Beginning in 2002, Idaho reports current asthma.

2 Results may differ from previous publications due to revisions in risk factor calculation.

3 Question wording changes make data prior to 2005 not comparable with 2005 and later.

4 Prior to 2005, Idaho BRFSS reported drinking status among all adults. Beginning in 2005, Idaho reports gender specific drinking status.

5 Question wording changes make data prior to 2002 not comparable with 2002 and later.

6 Prior to 2004, Idaho BRFSS reported all ages. Beginning in 2004, Idaho reports only women aged 40 and older.

# Trends of Selected Risk Factors

## By Public Health District, 2001-2008

	Year											
	2001			2002			2003			2004		
	%	95% CI	n									
<b>District 4</b>												
General Health	11.4	9.1 14.1	692	10.3	8.2 12.9	713	11.0	8.6 13.8	681	9.1	7.1 11.5	753
Health Care Coverage	14.3	11.5 17.7	692	13.8	11.0 17.2	714	11.2	8.7 14.3	680	11.7	9.3 14.7	753
Dental Insurance	38.1	33.8 42.6	672	36.1	32.2 40.3	705	32.7	28.8 36.7	672	33.8	29.9 38.0	748
Diabetes	5.3	3.8 7.4	693	4.2	2.9 6.1	714	4.9	3.3 7.1	683	4.3	3.0 6.1	754
Asthma <sup>1</sup>	7.7	5.8 10.1	692	6.7	4.9 9.1	709	8.9	6.7 11.7	681	7.9	5.9 10.5	748
Physical Activity	19.2	15.8 23.1	692	15.8	13.1 18.9	714	16.7	13.8 20.0	683	14.0	11.5 16.9	753
Overweight <sup>2</sup>	58.7	54.2 63.0	658	54.4	50.1 58.6	680	56.6	52.2 60.9	646	54.3	50.0 58.6	737
Obesity	20.0	16.9 23.5	658	19.4	16.4 22.9	680	19.9	16.5 23.7	646	16.8	14.0 20.1	737
Cigarette Smoking	20.9	17.7 24.6	693	21.1	17.8 24.9	713	18.2	15.2 21.7	681	15.6	12.7 19.0	753
Binge Drinking <sup>3,4</sup>	*	*	*	*	*	*	*	*	*	*	*	*
Heavy Drinking <sup>3,4</sup>	*	*	*	*	*	*	*	*	*	*	*	*
Illicit Drug Use <sup>5</sup>	*	*	*	8.0	5.6 11.3	711	7.9	5.6 10.9	681	7.5	5.3 10.6	748
Seat Belt Use	27.6	23.8 31.7	670	27.6	23.9 31.6	713	26.0	22.2 30.1	683	*	*	*
Colorectal Cancer Screening	52.8	45.6 59.9	256	44.5	38.1 51.1	264	*	*	*	47.7	41.6 53.9	313
Breast Cancer Screening <sup>6</sup>	*	*	*	26.0	20.6 32.3	243	*	*	*	30.6	25.0 36.9	286
Cervical Cancer Screening <sup>2</sup>	*	*	*	11.7	8.3 16.3	313	*	*	*	12.8	9.1 17.7	316
Prostate Cancer Screening	45.8	37.4 54.4	170	50.4	42.3 58.4	172	*	*	*	46.0	38.1 54.0	178
Dental Visits	30.3	26.6 34.3	673	25.5	22.0 29.4	714	25.9	22.4 29.8	680	28.8	25.1 32.8	752
<b>District 5</b>												
General Health	15.4	12.7 18.5	683	14.8	12.0 18.1	624	11.9	9.4 14.9	710	13.5	11.0 16.6	685
Health Care Coverage	17.6	14.6 21.1	685	21.3	17.7 25.5	622	20.7	17.2 24.6	707	16.8	13.6 20.6	683
Dental Insurance	54.9	50.7 59.0	664	55.8	51.2 60.2	619	55.4	51.1 59.7	700	51.9	47.5 56.2	679
Diabetes	5.0	3.5 7.0	685	5.6	4.0 7.8	624	5.8	4.1 8.2	711	7.0	5.2 9.4	687
Asthma <sup>1</sup>	8.6	6.7 11.1	685	7.3	5.4 9.8	622	7.2	5.4 9.4	706	8.4	6.3 11.1	686
Physical Activity	27.6	24.0 31.5	685	23.7	20.0 27.9	624	20.5	17.3 24.0	711	24.0	20.5 27.8	687
Overweight <sup>2</sup>	57.0	52.8 61.1	638	57.8	53.2 62.2	603	59.7	55.3 64.0	672	60.6	56.3 64.8	661
Obesity	19.1	15.9 22.7	638	19.5	16.2 23.3	603	22.3	18.9 26.2	672	21.6	18.3 25.4	661
Cigarette Smoking	21.3	18.2 24.8	683	18.8	15.4 22.8	622	21.0	17.7 24.8	709	20.0	16.6 23.9	685
Binge Drinking <sup>3,4</sup>	*	*	*	*	*	*	*	*	*	*	*	*
Heavy Drinking <sup>3,4</sup>	*	*	*	*	*	*	*	*	*	*	*	*
Illicit Drug Use <sup>5</sup>	*	*	*	6.1	4.0 9.3	624	6.2	4.2 9.1	709	5.3	3.4 8.1	684
Seat Belt Use	35.8	31.9 39.9	669	41.8	37.3 46.3	624	35.8	31.7 40.1	709	*	*	*
Colorectal Cancer Screening	59.1	52.7 65.3	287	57.8	51.6 63.7	303	*	*	*	50.2	44.4 56.0	353
Breast Cancer Screening <sup>6</sup>	*	*	*	45.8	39.2 52.6	252	*	*	*	44.3	38.1 50.7	287
Cervical Cancer Screening <sup>2</sup>	*	*	*	22.3	16.7 29.0	258	*	*	*	27.7	22.0 34.2	272
Prostate Cancer Screening	55.0	46.7 63.0	163	42.0	34.0 50.5	163	*	*	*	54.9	46.9 62.6	178
Dental Visits	37.1	33.2 41.3	669	37.1	32.9 41.5	621	37.3	33.2 41.5	705	38.2	34.0 42.5	678

\* Indicates data are not available for that year

1 Prior to 2002, Idaho BRFSS reported adults who have ever been told they have asthma. Beginning in 2002, Idaho reports current asthma status.

2 Results may differ from previous publications due to revisions in risk factor calculation.

3 Question wording changes make data prior to 2005 not comparable with 2005 and later.

4 Prior to 2005, Idaho BRFSS reported drinking status among all adults. Beginning in 2005, Idaho reports gender specific drinking status.

5 Question wording changes make data prior to 2002 not comparable with 2002 and later.

6 Prior to 2004, Idaho BRFSS reported all ages. Beginning in 2004, Idaho reports only women aged 40 and older.

# Trends of Selected Risk Factors

## By Public Health District, 2001-2008

	Year											
	2005			2006			2007			2008		
	%	95% CI	n									
<b>District 4</b>												
General Health	13.5	10.8 16.7	784	10.4	8.0 13.5	776	11.7	9.4 14.4	752	11.5	9.3 14.3	751
Health Care Coverage	12.9	9.4 17.4	783	16.3	12.8 20.7	775	15.9	12.3 20.2	751	16.5	12.8 21.1	750
Dental Insurance	34.0	29.6 38.6	735	38.0	33.6 42.6	736	33.5	29.3 37.9	716	31.6	27.4 36.2	710
Diabetes	6.0	4.5 7.9	786	6.6	4.9 8.9	776	7.2	5.5 9.2	752	6.0	4.6 8.0	751
Asthma <sup>1</sup>	5.7	4.1 7.7	782	9.5	7.2 12.3	769	9.4	7.0 12.5	747	9.9	7.3 13.2	750
Physical Activity	17.3	14.0 21.0	784	17.3	14.3 20.9	774	16.4	13.3 19.9	752	15.5	12.6 18.9	751
Overweight <sup>2</sup>	58.3	53.8 62.8	750	57.4	52.9 61.8	737	64.2	59.7 68.6	715	59.4	54.7 64.0	720
Obesity	22.6	19.2 26.4	750	22.1	18.8 25.7	737	22.9	19.3 27.0	715	21.1	17.8 24.9	720
Cigarette Smoking	16.2	13.3 19.7	785	15.3	12.4 18.9	775	17.5	14.1 21.6	751	18.3	14.8 22.3	749
Binge Drinking <sup>3,4</sup>	19.6	15.7 24.3	782	18.0	14.5 22.1	762	16.8	13.2 21.1	734	15.0	11.8 18.8	738
Heavy Drinking <sup>3,4</sup>	5.2	3.7 7.2	780	5.7	3.6 9.0	763	5.4	3.5 8.3	729	6.2	4.2 9.0	734
Illicit Drug Use <sup>5</sup>	4.8	2.9 7.9	725	5.3	3.3 8.4	728	6.6	4.0 10.7	700	6.3	4.0 10.0	708
Seat Belt Use	19.0	15.6 23.0	741	19.7	16.3 23.5	766	20.7	16.8 25.2	720	14.7	11.6 18.5	736
Colorectal Cancer Screening	*	*	*	40.0	34.5 45.8	366	*	*	*	35.6	30.5 41.1	406
Breast Cancer Screening <sup>6</sup>	*	*	*	32.7	27.3 38.7	332	*	*	*	26.8	21.8 32.5	330
Cervical Cancer Screening <sup>2</sup>	*	*	*	15.9	11.9 20.8	345	*	*	*	17.6	13.0 23.4	304
Prostate Cancer Screening	*	*	*	48.7	40.7 56.8	179	*	*	*	41.6	34.2 49.4	205
Dental Visits	26.2	22.5 30.3	750	27.1	23.3 31.4	773	25.3	21.4 29.7	716	26.8	22.7 31.3	750
<b>District 5</b>												
General Health	16.7	14.0 19.8	922	15.1	12.1 18.8	753	19.1	15.9 22.7	755	13.1	10.7 15.9	724
Health Care Coverage	24.7	21.1 28.6	923	24.2	19.9 29.1	752	24.7	20.7 29.1	755	21.2	17.2 26.0	727
Dental Insurance	57.3	53.2 61.2	867	59.7	55.1 64.1	717	55.0	50.5 59.5	700	49.5	44.7 54.2	680
Diabetes	7.3	5.8 9.3	924	5.5	4.0 7.5	754	10.0	7.8 12.7	755	7.7	5.9 9.8	726
Asthma <sup>1</sup>	8.4	6.2 11.1	920	8.7	6.7 11.4	749	7.6	5.6 10.2	755	7.1	5.3 9.4	723
Physical Activity	24.4	21.2 28.0	922	25.7	21.7 30.1	753	24.4	20.8 28.3	755	23.3	19.5 27.6	725
Overweight <sup>2</sup>	62.1	58.2 65.9	873	58.9	54.2 63.4	717	62.9	58.3 67.4	709	62.8	58.0 67.2	679
Obesity	23.1	19.9 26.7	873	24.5	20.9 28.5	717	26.3	22.7 30.2	709	22.8	19.4 26.7	679
Cigarette Smoking	20.5	17.4 24.1	921	18.3	14.8 22.3	753	23.0	19.3 27.1	754	16.1	12.6 20.3	725
Binge Drinking <sup>3,4</sup>	12.5	9.7 15.9	917	18.2	14.4 22.8	747	13.5	10.6 17.2	744	11.9	8.6 16.2	715
Heavy Drinking <sup>3,4</sup>	4.0	2.5 6.3	915	7.6	5.1 11.2	742	5.8	3.7 9.0	740	3.3	2.2 4.9	709
Illicit Drug Use <sup>5</sup>	5.3	3.6 7.7	862	3.8	2.3 6.2	713	4.9	2.9 8.2	694	2.8	1.5 5.1	673
Seat Belt Use	25.0	21.6 28.6	875	34.8	30.4 39.5	742	25.7	21.7 30.1	706	27.2	23.2 31.6	716
Colorectal Cancer Screening	*	*	*	48.1	42.8 53.6	414	*	*	*	49.2	44.0 54.4	433
Breast Cancer Screening <sup>6</sup>	*	*	*	40.9	35.2 46.8	345	*	*	*	40.9	35.2 46.9	331
Cervical Cancer Screening <sup>2</sup>	*	*	*	20.1	14.6 26.9	315	*	*	*	22.4	17.3 28.6	279
Prostate Cancer Screening	*	*	*	49.7	42.0 57.4	192	*	*	*	44.8	37.2 52.7	212
Dental Visits	39.5	35.7 43.5	879	43.2	38.6 47.9	752	34.1	29.9 38.6	703	36.2	31.8 40.8	726

1 Prior to 2002, Idaho BRFSS reported adults who have ever been told they have asthma. Beginning in 2002, Idaho reports current asthma.

2 Results may differ from previous publications due to revisions in risk factor calculation.

3 Question wording changes make data prior to 2005 not comparable with 2005 and later.

4 Prior to 2005, Idaho BRFSS reported drinking status among all adults. Beginning in 2005, Idaho reports gender specific drinking status.

5 Question wording changes make data prior to 2002 not comparable with 2002 and later.

6 Prior to 2004, Idaho BRFSS reported all ages. Beginning in 2004, Idaho reports only women aged 40 and older.

# Trends of Selected Risk Factors

## By Public Health District, 2001-2008

	Year											
	2001			2002			2003			2004		
	%	95% CI	n									
<b>District 6</b>												
General Health	11.9	9.5 14.8	697	14.0	11.5 16.9	704	14.4	11.8 17.5	693	14.1	11.7 17.0	756
Health Care Coverage	11.5	9.2 14.4	697	15.7	12.7 19.2	704	13.8	11.1 17.1	692	14.0	11.2 17.5	755
Dental Insurance	41.3	37.3 45.4	684	42.4	38.3 46.6	692	46.2	41.9 50.4	684	41.5	37.5 45.7	751
Diabetes	6.4	4.7 8.6	699	7.9	6.0 10.2	704	8.2	6.2 10.7	693	8.9	7.1 11.2	759
Asthma <sup>1</sup>	7.4	5.6 9.6	697	7.5	5.7 10.0	703	8.8	6.7 11.5	690	6.7	5.1 8.7	753
Physical Activity	18.2	15.4 21.5	700	23.6	20.3 27.2	705	20.1	16.8 23.9	692	17.2	14.6 20.3	757
Overweight <sup>2</sup>	59.9	55.9 63.8	668	58.5	54.2 62.6	681	61.3	57.0 65.5	665	57.8	53.5 62.1	726
Obesity	24.1	20.7 27.8	668	20.2	17.1 23.7	681	26.6	22.9 30.8	665	21.7	18.7 25.1	726
Cigarette Smoking	15.8	13.2 18.9	698	19.1	16.0 22.5	704	17.7	14.8 21.1	693	16.6	13.7 19.9	754
Binge Drinking <sup>3,4</sup>	*	* *	*	*	* *	*	*	* *	*	*	* *	*
Heavy Drinking <sup>3,4</sup>	*	* *	*	*	* *	*	*	* *	*	*	* *	*
Illicit Drug Use <sup>5</sup>	*	* *	*	5.1	3.4 7.5	703	5.8	4.0 8.3	690	3.1	2.0 4.9	755
Seat Belt Use	43.9	39.9 48.0	688	45.1	40.9 49.3	702	41.0	36.8 45.3	692	*	* *	*
Colorectal Cancer Screening	64.7	58.5 70.5	279	59.1	52.7 65.2	300	*	* *	*	63.8	58.3 68.9	367
Breast Cancer Screening <sup>6</sup>	*	* *	*	41.2	35.0 47.7	271	*	* *	*	42.1	36.1 48.3	302
Cervical Cancer Screening <sup>2</sup>	*	* *	*	18.6	14.1 24.2	302	*	* *	*	23.8	18.2 30.5	296
Prostate Cancer Screening	52.8	44.8 60.7	174	49.7	41.4 57.9	175	*	* *	*	55.6	48.2 62.9	200
Dental Visits	30.8	27.2 34.6	684	30.9	27.2 34.8	704	35.6	31.6 39.8	689	32.3	28.6 36.3	751
<b>District 7</b>												
General Health	10.6	8.4 13.3	698	12.1	9.5 15.1	695	12.2	10.0 14.8	715	10.5	8.4 13.0	715
Health Care Coverage	13.5	10.8 16.9	695	15.8	12.6 19.5	693	13.6	10.8 17.0	715	14.9	12.0 18.3	714
Dental Insurance	42.8	38.7 47.0	682	44.6	40.1 49.1	693	43.0	38.7 47.4	711	41.1	37.0 45.4	705
Diabetes	5.7	4.1 7.7	699	6.3	4.6 8.5	694	5.8	4.1 8.0	716	7.2	5.4 9.4	716
Asthma <sup>1</sup>	8.7	6.7 11.3	698	8.9	6.7 11.8	695	6.0	4.4 8.1	714	6.5	4.8 8.7	714
Physical Activity	17.7	14.7 21.2	699	17.7	14.8 21.0	695	15.7	13.1 18.7	715	16.6	13.7 20.0	715
Overweight <sup>2</sup>	59.8	55.3 64.0	670	55.6	50.9 60.3	669	60.0	55.5 64.3	689	60.4	55.9 64.7	695
Obesity	17.6	14.8 20.9	670	18.8	15.7 22.3	669	21.4	17.8 25.4	689	25.9	22.2 30.0	695
Cigarette Smoking	12.4	10.1 15.3	698	16.4	13.3 20.0	695	13.5	10.9 16.7	714	12.2	9.8 15.1	715
Binge Drinking <sup>3,4</sup>	*	* *	*	*	* *	*	*	* *	*	*	* *	*
Heavy Drinking <sup>3,4</sup>	*	* *	*	*	* *	*	*	* *	*	*	* *	*
Illicit Drug Use <sup>5</sup>	*	* *	*	2.7	1.7 4.4	695	3.4	2.0 5.7	714	5.6	3.7 8.5	717
Seat Belt Use	41.3	37.3 45.5	688	40.5	36.0 45.1	694	43.0	38.7 47.5	713	*	* *	*
Colorectal Cancer Screening	54.8	48.3 61.2	270	52.1	46.1 58.2	314	*	* *	*	49.9	44.1 55.8	335
Breast Cancer Screening <sup>6</sup>	*	* *	*	47.5	41.0 54.0	269	*	* *	*	49.6	43.1 56.1	273
Cervical Cancer Screening <sup>2</sup>	*	* *	*	19.4	14.7 25.1	292	*	* *	*	30.9	24.5 38.2	295
Prostate Cancer Screening	55.1	47.3 62.7	189	55.2	47.3 62.9	176	*	* *	*	49.0	41.2 56.7	186
Dental Visits	28.8	25.0 32.8	683	31.1	27.0 35.5	693	29.8	25.9 34.1	712	26.8	23.2 30.6	710

\* Indicates data are not available for that year

1 Prior to 2002, Idaho BRFSS reported adults who have ever been told they have asthma. Beginning in 2002, Idaho reports current asthma status.

2 Results may differ from previous publications due to revisions in risk factor calculation.

3 Question wording changes make data prior to 2005 not comparable with 2005 and later.

4 Prior to 2005, Idaho BRFSS reported drinking status among all adults. Beginning in 2005, Idaho reports gender specific drinking status.

5 Question wording changes make data prior to 2002 not comparable with 2002 and later.

6 Prior to 2004, Idaho BRFSS reported all ages. Beginning in 2004, Idaho reports only women aged 40 and older.

# Trends of Selected Risk Factors

## By Public Health District, 2001-2008

	Year											
	2005			2006			2007			2008		
	%	95% CI	n									
<b>District 6</b>												
General Health	15.7	13.0 18.9	757	12.6	10.3 15.2	768	18.0	14.9 21.6	755	14.2	11.5 17.3	704
Health Care Coverage	15.1	11.9 18.9	755	13.2	10.7 16.3	765	18.4	14.8 22.6	754	14.2	10.6 18.6	703
Dental Insurance	44.1	39.8 48.6	723	40.6	36.5 44.7	726	41.9	37.4 46.4	713	43.7	38.4 49.1	664
Diabetes	7.1	5.4 9.1	757	6.9	5.3 8.8	768	8.3	6.5 10.5	756	8.9	6.8 11.5	704
Asthma <sup>1</sup>	6.7	5.1 8.7	754	8.8	6.9 11.1	757	7.5	5.4 10.2	748	6.5	4.6 9.2	699
Physical Activity	21.0	17.8 24.5	757	21.8	18.7 25.2	765	21.5	18.3 25.1	751	22.3	18.5 26.6	704
Overweight <sup>2</sup>	63.5	58.9 67.8	713	61.7	57.3 65.8	736	61.7	57.0 66.2	722	65.1	59.6 70.2	674
Obesity	28.5	24.4 33.0	713	27.2	23.7 31.0	736	24.6	21.0 28.7	722	32.9	28.2 38.1	674
Cigarette Smoking	15.9	13.1 19.2	758	15.0	12.3 18.2	764	19.1	15.7 23.2	754	12.0	9.3 15.4	701
Binge Drinking <sup>3,4</sup>	12.6	9.6 16.5	752	10.4	7.9 13.5	754	15.0	11.4 19.5	747	8.6	6.2 11.9	698
Heavy Drinking <sup>3,4</sup>	2.5	1.6 4.0	752	2.9	1.8 4.7	757	6.0	3.5 10.0	741	2.6	1.4 4.8	695
Illicit Drug Use <sup>5</sup>	6.2	3.7 10.3	707	3.8	2.2 6.6	730	5.7	3.1 10.3	709	3.8	1.8 7.7	665
Seat Belt Use	32.5	28.3 37.0	721	37.1	33.1 41.4	750	32.8	28.4 37.5	716	38.5	33.2 44.1	695
Colorectal Cancer Screening	*	*	*	53.8	48.3 59.2	379	*	*	*	51.2	45.8 56.5	425
Breast Cancer Screening <sup>6</sup>	*	*	*	43.3	37.3 49.4	308	*	*	*	39.0	33.3 45.0	336
Cervical Cancer Screening <sup>2</sup>	*	*	*	25.9	19.5 33.5	269	*	*	*	31.9	23.1 42.2	274
Prostate Cancer Screening	*	*	*	45.5	38.3 52.8	202	*	*	*	48.9	41.0 56.9	190
Dental Visits	32.1	28.0 36.5	735	32.4	28.6 36.4	764	35.3	31.0 39.9	717	30.4	25.9 35.3	699
<b>District 7</b>												
General Health	12.6	10.3 15.4	785	12.2	9.8 15.1	758	14.9	11.6 18.7	749	12.6	10.0 15.8	692
Health Care Coverage	16.3	13.4 19.6	783	14.0	11.0 17.5	758	16.7	13.2 21.0	751	14.6	11.0 19.0	692
Dental Insurance	46.0	41.6 50.4	729	47.9	43.4 52.3	724	42.0	37.4 46.8	707	46.6	41.2 52.1	665
Diabetes	6.7	5.1 8.7	782	7.0	5.3 9.1	757	6.3	4.8 8.1	750	7.1	5.5 9.1	695
Asthma <sup>1</sup>	9.1	6.7 12.2	782	9.6	7.3 12.7	753	10.5	7.7 14.1	747	6.0	4.3 8.3	688
Physical Activity	21.1	18.0 24.7	783	21.4	17.9 25.3	759	16.9	14.0 20.3	752	21.1	17.4 25.4	695
Overweight <sup>2</sup>	59.7	55.2 64.0	752	57.5	52.8 62.0	722	58.4	53.3 63.3	707	56.8	51.1 62.3	664
Obesity	23.2	19.9 26.8	752	22.7	19.4 26.4	722	24.4	20.8 28.4	707	21.4	17.9 25.4	664
Cigarette Smoking	12.1	9.8 14.8	784	9.3	7.3 11.8	759	11.5	9.1 14.5	751	12.2	9.1 16.1	695
Binge Drinking <sup>3,4</sup>	7.6	5.8 9.9	779	7.5	5.7 9.8	753	8.7	6.5 11.4	749	10.8	7.7 15.0	688
Heavy Drinking <sup>3,4</sup>	3.6	2.4 5.3	773	2.1	1.3 3.2	747	2.3	1.4 3.7	743	2.1	1.1 4.0	687
Illicit Drug Use <sup>5</sup>	3.2	2.0 5.1	728	1.8	0.8 3.7	725	3.3	1.8 5.8	706	1.3	0.7 2.6	664
Seat Belt Use	34.0	29.8 38.4	732	35.9	31.7 40.2	748	31.5	27.1 36.3	714	31.3	26.8 36.2	687
Colorectal Cancer Screening	*	*	*	46.4	40.6 52.3	336	*	*	*	50.0	44.3 55.6	367
Breast Cancer Screening <sup>6</sup>	*	*	*	39.4	33.3 45.7	278	*	*	*	42.0	35.7 48.5	274
Cervical Cancer Screening <sup>2</sup>	*	*	*	35.0	28.1 42.7	325	*	*	*	33.7	24.8 44.0	261
Prostate Cancer Screening	*	*	*	48.2	40.3 56.2	181	*	*	*	54.3	46.7 61.6	206
Dental Visits	29.6	25.8 33.7	741	29.0	25.1 33.3	758	32.9	28.4 37.8	714	28.8	24.3 33.8	695

1 Prior to 2002, Idaho BRFSS reported adults who have ever been told they have asthma. Beginning in 2002, Idaho reports current asthma.

2 Results may differ from previous publications due to revisions in risk factor calculation.

3 Question wording changes make data prior to 2005 not comparable with 2005 and later.

4 Prior to 2005, Idaho BRFSS reported drinking status among all adults. Beginning in 2005, Idaho reports gender specific drinking status.

5 Question wording changes make data prior to 2002 not comparable with 2002 and later.

6 Prior to 2004, Idaho BRFSS reported all ages. Beginning in 2004, Idaho reports only women aged 40 and older.



## Appendix B

---

### 2008 Behavioral Risk Factor Surveillance System Questionnaire

In the following pages, the 2008 Idaho BRFSS questionnaire is presented in its entirety. In addition to the text of each question, the results for most of the questions are included. The number of people who responded to the question (n) is listed below the text for each question. The percent for each response category can be found in parentheses to the right of the response categories. Note also that the percentages are weighted results (see Methodology on pages 1-2) and do not directly correspond to the n's for each question. The results have not been provided for all questions.

**NOTE:** Care should be exercised when interpreting these results. In order to interpret the results correctly, it is important to understand what group has been asked the question. For example, in question ID2.3 on page 106, 17.8% of the respondents have Medicare. However this question is asked only of those who respond “yes” to question Q3.1. The exclusion of certain respondents is indicated by the bold text at the beginning of question ID2.3. These are called “skip patterns” and are used extensively throughout the survey. In order to properly identify what group has been asked a specific question, the skip pattern must be considered. In addition, there may be some confusion when trying to compare results from the appendix to those in the previous sections. The risk factors reported in the main body of the report are often constructed from the responses of multiple questions and may not be comparable to the results of one particular question.

Response categories displayed in capital letters were not read aloud. Responses of “Don't Know” and “Refused” are excluded from the n's, percentage, and means calculations. Other responses may also be excluded and are indicating by showing the response without any percentage. If there are fewer than 50 respondents for a question the results are suppressed as the results are considered unreliable. Also the results of some questions, such as Q12.15 about county of residence, are suppressed due to length of resulting lists.

For help in interpreting these results, contact the Idaho BRFSS Project Director at (208) 332-7326.

## SECTION 1: HEALTH STATUS

Q1.1 Would you say that in general your health is ...  
(n=5099)

- 1 Excellent (22.1%)
- 2 Very good (35.8%)
- 3 Good (28.1%)
- 4 Fair (10.5%)
- 5 Poor (3.6%)
  
- 7 DON'T KNOW/NOT SURE
- 9 REFUSED

## SECTION 2: HEALTHY DAYS AND HEALTH-RELATED QUALITY OF LIFE

Q2.1 Now thinking about your physical health, which includes physical illness and injury, for how many days during the past 30 days was your physical health not good?  
(n=5015)

Mean number of days not good overall: 3.3  
Reported any days not good: 36.1%  
Mean number of days for those with any: 9.2

- 88 NONE
- 77 DON'T KNOW/NOT SURE
- 99 REFUSED

Q2.2 Now thinking about your mental health, which includes stress, depression, and problems with emotions, for how many days during the past 30 days was your mental health not good?  
(n=5032)

Mean number of days not good overall: 3.3  
Reported any days not good: 35.5%  
Mean number of days for those with any: 9.2

- 88 NONE
- 77 DON'T KNOW/NOT SURE
- 99 REFUSED

Q2.3 **[IF Q2.1<>88 OR Q2.2<>88]** During the past 30 days, for about how many days did poor physical or mental health keep you from doing your usual activities, such as self-care, work, or recreation?  
(n=2770)

Mean number of days limited overall: 3.8  
Reported any days limited: 40.5%  
Mean number of days for those with any: 9.5

- 88 NONE
- 77 DON'T KNOW/NOT SURE
- 99 REFUSED

## SECTION 3: HEALTH CARE ACCESS

Q3.1 Do you have any kind of health care coverage, including health insurance, prepaid plans such as HMOs, or government plans such as Medicare?  
(n=5099)

- 1 Yes (81.5%)
- 2 No (18.5%)
  
- 7 DON'T KNOW/NOT SURE
- 9 REFUSED

Q3.2 Do you have one person you think of as your personal doctor or health care provider?  
(n=5096)

- 1 Yes, only one (64.3%)
- 2 More than one (8.4%)
- 3 No (27.3%)
  
- 7 DON'T KNOW/NOT SURE
- 9 REFUSED

Q3.3 Was there a time in the past 12 months when you needed to see a doctor but could not because of cost?  
(n=5100)

- 1 Yes (16.4%)
- 2 No (83.7%)
  
- 7 DON'T KNOW/NOT SURE
- 9 REFUSED

Q3.4 About how long has it been since you last visited a doctor for a routine checkup? A routine checkup is a general physical exam, not an exam for a specific injury, illness, or condition.  
(n=5089)

- 1 Within past year (anytime less than 12 months ago) (55.6%)
- 2 Within past 2 years (1 year but less than 2 years ago) (15.6%)
- 3 Within past 5 years (2 years but less than 5 years ago) (11.0%)
- 4 5 or more years ago (16.3%)
- 8 Never (1.5%)
  
- 7 DON'T KNOW/NOT SURE
- 9 REFUSED

## SECTION 4: SLEEP

Q4.1 The next question is about getting enough rest or sleep. During the past 30 days, for about how many days have you felt you did not get enough rest or sleep?  
(n=5013)

Mean number of days not enough sleep overall: 8.3  
Reported any days not enough sleep: 70.5%  
Mean number of days for those with any: 11.8

- 88 NONE
- 77 DON'T KNOW/NOT SURE
- 99 REFUSED

## SECTION 5: EXERCISE

Q5.1 During the past month, other than your regular job, did you participate in any physical activities or exercises such as running, calisthenics, golf, gardening, or walking for exercise?  
(n=5104)

- 1 Yes (78.9%)
- 2 No (21.1%)
  
- 7 DON'T KNOW/NOT SURE
- 9 REFUSED

## SECTION 6: DIABETES

Q6.1 Have you ever been told by a doctor that you have diabetes?  
(n=5108)

- 1 Yes (7.0%)
- 2 Yes, but female told only during pregnancy (0.9%)
- 3 No (90.8%)
- 4 No, pre-diabetes or borderline diabetes (1.3%)
  
- 7 DON'T KNOW/NOT SURE
- 9 REFUSED

## MODULE 2: DIABETES

M2.1 **[IF Q6.1=1]** How old were you when you were told you have diabetes?  
(n=498)

Mean age in years: 51.4

98 DON'T KNOW/NOT SURE  
99 REFUSED

M2.2 **[IF Q6.1=1]** Are you now taking insulin?  
(n=526)

1 Yes (28.4%)  
2 No (71.6%)

9 REFUSED

M2.3 **[IF Q6.1=1]** About how often do you check your blood for glucose or sugar? Include times when checked by a family member or friend, but do NOT include times when checked by a health professional.  
(n=519)

Mean number of times per day: 1.5

888 NEVER  
777 DON'T KNOW/NOT SURE  
999 REFUSED

M2.4 **[IF Q6.1=1]** About how often do you check your feet for any sores or irritations? Include times when checked by a family member or friend, but do NOT include times when checked by a health professional.  
(n=518)

Mean number of times per day: 0.7

555 NO FEET  
888 NEVER  
777 DON'T KNOW/NOT SURE  
999 REFUSED

M2.5 **[IF Q6.1=1]** About how many times in the past 12 months have you seen a doctor, nurse, or other health professional for your diabetes?  
(n=516)

Mean number of times per day: 3.2

88 NONE  
77 DON'T KNOW/NOT SURE  
99 REFUSED

M2.6 **[IF Q6.1=1]** A test for "A one C" measures the average level of blood sugar over the past three months. About how many times in the past 12 months has a doctor, nurse, or other health professional checked you for "A one C"?  
(n=501)

Mean number of times in past 12 months: 2.4

98 NEVER HEARD OF "A ONE C" TEST (4.5%)

88 NEVER  
77 DON'T KNOW/NOT SURE  
99 REFUSED

M2.7 **[IF Q6.1=1 AND M2.4<=555]** About how many times in the past 12 months has a health professional checked your feet for any sores or irritations?  
(n=503)

Mean number of times in past 12 months: 2.1

88 NEVER  
77 DON'T KNOW/NOT SURE  
99 REFUSED

M2.8 **[IF Q6.1=1]** When was the last time you had an eye exam in which the pupils were dilated? This would have made you temporarily sensitive to bright light.  
(n=523)

1 Within the past month (anytime less than 1 month ago) (9.4%)  
2 Within the past year (1 month but less than 12 months ago) (53.7%)  
3 Within the past 2 years (1 year but less than 2 years ago) (13.0%)  
4 2 or more years ago (19.8%)  
8 Never (4.1%)  
7 DON'T KNOW/NOT SURE  
9 REFUSED

M2.9 **[IF Q6.1=1]** Has a doctor ever told you that diabetes has affected your eyes or that you had retinopathy?  
(n=520)

1 Yes (20.0%)  
2 No (80.0%)  
7 DON'T KNOW/NOT SURE  
9 REFUSED

M2.10 **[IF Q6.1=1]** Have you ever taken a course or class in how to manage your diabetes yourself?  
(n=525)

1 Yes (56.8%)  
2 No (43.2%)  
7 DON'T KNOW/NOT SURE  
9 REFUSED

## SECTION 7: ORAL HEALTH

Q7.1 How long has it been since you last visited a dentist or a dental clinic for any reason?  
(n=5091)

1 Within the past year (anytime less than 12 months ago) (68.0%)  
2 Within the past 2 years (1 year but less than 2 years ago) (11.6%)  
3 Within the past 5 years (2 years but less than 5 years ago) (9.7%)  
4 5 or more years ago (10.1%)  
8 Never (0.7%)  
7 DON'T KNOW/NOT SURE  
9 REFUSED

Q7.2 How many of your permanent teeth have been removed because of tooth decay or gum disease? Include teeth lost to infection, but do not include teeth lost for other reasons, such as injury or orthodontics.  
(n=5047)

1 1 to 5 (29.2%)  
2 6 or more, but not all (7.9%)  
3 All (4.6%)  
8 None (58.3%)  
7 DON'T KNOW/NOT SURE  
9 REFUSED

Q7.3 **[IF Q7.1<=>8 AND Q7.2<=>3]** How long has it been since you had your teeth cleaned by a dentist or dental hygienist?  
(n=4672)

- 1 Within the past year (anytime less than 12 months ago) (67.4%)
- 2 Within the past 2 years (1 year but less than 2 years ago) (10.6%)
- 3 Within the past 5 years (2 years but less than 5 years ago) (10.3%)
- 4 5 or more years ago (10.3%)
- 8 Never (1.4%)
- 7 DON'T KNOW/NOT SURE
- 9 REFUSED

#### SECTION 8: CARDIOVASCULAR DISEASE PREVALENCE

Q8.1 Has a doctor, nurse, or other health professional ever told you that you had any of the following? (Ever told) you had a heart attack, also called a myocardial infarction?  
(n=5088)

- 1 Yes (4.3%)
- 2 No (95.7%)
- 7 DON'T KNOW/NOT SURE
- 9 REFUSED

Q8.2 (Ever told) you had angina or coronary heart disease?  
(n=5056)

- 1 Yes (4.0%)
- 2 No (96.0%)
- 7 DON'T KNOW/NOT SURE
- 9 REFUSED

Q8.3 (Ever told) you had a stroke?  
(n=5101)

- 1 Yes (2.3%)
- 2 No (97.7%)
- 7 DON'T KNOW/NOT SURE
- 9 REFUSED

#### SECTION 9: ASTHMA

Q9.1 Have you ever been told by a doctor, nurse, or other health professional that you had asthma?  
(n=5094)

- 1 Yes (12.7%)
- 2 No (87.3%)
- 7 DON'T KNOW/NOT SURE
- 9 REFUSED

Q9.2 **[IF Q9.1=1]** Do you still have asthma?  
(n=632)

- 1 Yes (72.0%)
- 2 No (28.0%)
- 7 DON'T KNOW/NOT SURE
- 9 REFUSED

#### SECTION 10: DISABILITY

Q10.1 Are you limited in any way in any activities because of physical, mental, or emotional problems?  
(n=5083)

- 1 Yes (22.6%)
- 2 No (77.4%)
- 7 DON'T KNOW/NOT SURE
- 9 REFUSED

Q10.2 Do you now have any health problem that requires you to use special equipment, such as a cane, a wheelchair, a special bed, or a special telephone?  
(n=5107)

- 1 Yes (6.7%)
- 2 No (93.3%)
- 7 DON'T KNOW/NOT SURE
- 9 REFUSED

#### SECTION 11: TOBACCO USE

Q11.1 Have you smoked at least 100 cigarettes in your entire life?  
(n=5097)

- 1 Yes (38.8%)
- 2 No (61.2%)
- 7 DON'T KNOW/NOT SURE
- 9 REFUSED

Q11.2 **[IF Q11.1=1]** Do you now smoke cigarettes every day, some days, or not at all?  
(n=2122)

- 1 Every day (32.2%)
- 2 Some days (11.4%)
- 3 Not at all (56.5%)
- 7 DON'T KNOW/NOT SURE
- 9 REFUSED

Q11.3 **[IF Q11.1=1 AND Q11.2=1 OR 2]** During the past 12 months, have you stopped smoking for one day or longer because you were trying to quit smoking?  
(n=798)

- 1 Yes (56.2%)
- 2 No (43.8%)
- 7 DON'T KNOW/NOT SURE
- 9 REFUSED

#### SECTION 12: DEMOGRAPHICS

Q12.1 What is your age?  
(n=5069)

Mean age in years: 45.6

- 7 DON'T KNOW/NOT SURE
- 9 REFUSED

Q12.2 Are you Hispanic or Latino?  
(n=5093)

- 1 Yes (7.6%)
- 2 No (92.4%)
- 7 DON'T KNOW/NOT SURE
- 9 REFUSED

Q12.3 Which one or more of the following would you say is your race?

[SEE RESULTS FOLLOWING Q12.4]

Q12.4 Which one of these groups would you say best represents your race?  
(n=5049)

- 1 White (94.0%)
- 2 Black or African American (0.6%)
- 3 Asian (0.5%)
- 4 Native Hawaiian or Other Pacific Islander (0.4%)
- 5 American Indian, Alaska Native (1.8%)
- 6 Other (2.8%)
- 7 DON'T KNOW/NOT SURE
- 9 REFUSED

Q12.5 Have you ever served on active duty in the United States Armed Forces, either in the regular military or in a National Guard or military reserve unit? Active duty does not include training for the Reserves or National Guard, but DOES include activation, for example, the Persian Gulf War.  
(n=5109)

- 1 Yes (12.7%)
- 2 No (87.4%)
- 7 DON'T KNOW/NOT SURE
- 9 REFUSED

Q12.6 Are you ... ?  
(n=5095)

- 1 Married (68.1%)
- 2 Divorced (8.9%)
- 3 Widowed (5.3%)
- 4 Separated (1.1%)
- 5 Never married (13.2%)
- 6 A member of an unmarried couple (3.4%)
- 9 REFUSED

Q12.7 How many children less than 18 years of age live in your household?  
(n=5107)

Mean number of children: 1.1

- 88 NONE
- 99 REFUSED

Q12.8 What is the highest grade or year of school you completed?  
(n=5100)

- 1 Never attended school or only kindergarten (0.1%)
- 2 Grades 1 through 8 (Elementary) (2.6%)
- 3 Grades 9 through 11 (Some high school) (7.5%)
- 4 Grade 12 or GED (High school graduate) (28.9%)
- 5 College 1 year to 3 years (Some college or technical school) (30.8%)
- 6 College 4 years or more (College graduate) (30.2%)
- 9 REFUSED

Q12.9 Are you currently ... ?  
(n=5103)

- 1 Employed for wages (49.7%)
- 2 Self-employed (11.6%)
- 3 Out of work for more than 1 year (1.6%)
- 4 Out of work for less than 1 year (3.0%)
- 5 A homemaker (9.9%)
- 6 A student (4.4%)
- 7 Retired (15.6%)
- 8 Unable to work (4.2%)
- 9 REFUSED

Q12.10 Is your annual household income from all sources ...  
(n=4534)

- 1 Less than \$10,000 (4.1%)
- 2 \$10,000 to less than \$15,000 (4.3%)
- 3 \$15,000 to less than \$20,000 (7.5%)
- 4 \$20,000 to less than \$25,000 (9.9%)
- 5 \$25,000 to less than \$35,000 (14.8%)
- 6 \$35,000 to less than \$50,000 (18.2%)
- 7 \$50,000 to less than \$75,000 (18.0%)
- 8 \$75,000 or more (23.4%)
- 77 DON'T KNOW/NOT SURE
- 99 REFUSED

Q12.11 About how much do you weigh without shoes?  
(n=4879)

Mean weight in pounds (Male): 194.4  
Mean weight in pounds (Female): 159.6

- 7777 DON'T KNOW/NOT SURE
- 9999 REFUSED

Q12.12 [IF Q12.1<=>7777 OR 9999] About how tall are you without shoes?  
(n=5062)

Mean height in feet and inches (Male): 5 ft. 11 in.  
Mean height in feet and inches (Female): 5 ft. 5 in.

- 7777 DON'T KNOW/NOT SURE
- 9999 REFUSED

Q12.13 How much did you weigh a year ago?  
(n=4814)

Mean weight in pounds (Male): 195.0  
Mean weight in pounds (Female): 160.0

- 7777 DON'T KNOW/NOT SURE
- 9999 REFUSED

Q12.14 Was the change between your current weight and your weight a year ago intentional?  
(n=2164)

- 1 Yes (42.6%)
- 2 No (57.4%)

- 7 DON'T KNOW/NOT SURE
- 9 REFUSED

Q12.15 What county do you live in?

(Due to the size of the table, these data are not printed.)

Q12.16 What is your ZIP Code where you live?

(ZIP codes are not available.)

Q12.17 Do you have more than one telephone number in your household? Do not include cell phones or numbers that are only used by a computer or fax machine.  
(n=5108)

- 1 Yes (4.4%)
- 2 No (95.6%)

- 7 DON'T KNOW/NOT SURE
- 9 REFUSED

Q12.18 **[IF Q12.7=1]** How many of these telephone numbers are residential numbers?  
(n=237)

- 1 One (67.3%)
- 2 Two (31.3%)
- 3 Three (1.1%)
- 4 Four (0.3%)
- 5 Five (0.6%)
  
- 7 DON'T KNOW/NOT SURE
- 9 REFUSED

Q12.19 During the past 12 months, has your household been without telephone service for 1 week or more? Do not include interruptions of telephone service because of weather or natural disasters.  
(n=5092)

- 1 Yes (4.2%)
- 2 No (95.8%)
  
- 7 DON'T KNOW/NOT SURE
- 9 REFUSED

Q12.20 Indicate sex of respondent.  
(n=5111)

- 1 Male (49.8%)
- 2 Female (50.2%)

Q12.21 **[IF Q12.0=2 AND Q12.1<45]** To your knowledge, are you now pregnant?  
(n=1000)

- 1 Yes (4.5%)
- 2 No (95.5%)
  
- 7 DON'T KNOW/NOT SURE
- 9 REFUSED

### SECTION 13: ALCOHOL CONSUMPTION

Q13.1 During the past 30 days, have you had at least one drink of any alcoholic beverage such as beer, wine, a malt beverage or liquor?  
(n=5079)

- 1 Yes (46.9%)
- 2 No (53.1%)
  
- 7 DON'T KNOW/NOT SURE
- 9 REFUSED

Q13.2 **[IF Q13.1=1]** During the past 30 days, how many days per week or per month did you have at least one drink of any alcoholic beverage?  
(n=2191)

Mean number of days in the last 30: 9.2

- 888 NO DRINKS IN LAST 30 DAYS
- 777 DON'T KNOW/NOT SURE
- 999 REFUSED

Q13.3 **[IF Q13.1=1 AND Q13.2<=888]** One drink is equivalent to a 12-ounce beer, a 5-ounce glass of wine, or a drink with one shot of liquor. During the past 30 days, on the days when you drank, about how many drinks did you drink on the average?  
(n=2192)

Mean number of drinks: 2.3

- 77 DON'T KNOW/NOT SURE
- 99 REFUSED

Q13.4 **[IF Q13.1=1 AND Q13.2<=888]** Considering all types of alcoholic beverages, how many times during the past 30 days did you have (IF MALE) 5 or more drinks on an occasion? (IF FEMALE) 4 or more drinks on an occasion?  
(n=2183)

Mean number of times overall: 1.3  
Reported any occasions: 28.5%  
Mean number of days for those that had any: 4.8

- 88 NONE
- 77 DON'T KNOW/NOT SURE
- 99 REFUSED

Q13.5 **[IF Q13.1=1 AND Q13.2<=888]** During the past 30 days, what is the largest number of drinks you had on any occasion?  
(n=2136)

Mean number of most drinks: 3.5

- 77 DON'T KNOW/NOT SURE
- 99 REFUSED

### SECTION 14: IMMUNIZATION

Q14.1 A flu shot is an influenza vaccine injected into your arm. During the past 12 months, have you had a flu shot?  
(n=5058)

- 1 Yes (33.0%)
- 2 No (67.0%)

- 7 DON'T KNOW/NOT SURE
- 9 REFUSED

Q14.2 **[IF Q14.1=1]** During what month and year did you receive your most recent flu shot?  
(n=1989)

Mode month of most recent flu shot: October, 2007

- 77/7777 DON'T KNOW/NOT SURE
- 99/9999 REFUSED

Q14.3 **[IF Q14.1<=1]** During the past 12 months, have you had a flu vaccine that was sprayed in your nose? The flu vaccine sprayed in the nose is also called FluMist®.  
(n=5063)

- 1 Yes (0.7%)
- 2 No (99.3%)

- 7 DON'T KNOW/NOT SURE
- 9 REFUSED

Q14.4 **[IF Q14.3=1]** During what month and year did you receive your most recent flu vaccine that was sprayed in your nose?  
(n<50)

Figure not reliable by BRFSS standards (n<50).

Q14.5 A pneumonia shot or pneumococcal vaccine is usually given only once or twice in a person's lifetime and is different from the flu shot. Have you ever had a pneumonia shot?  
(n=4713)

- 1 Yes (20.9%)
- 2 No (79.1%)

- 7 DON'T KNOW/NOT SURE
- 9 REFUSED

## SECTION 15: FALLS

Q15.1 **[IF Q12.1>= 45]** The next question asks about recent falls. By a fall, we mean when a person unintentionally comes to rest on the ground or another lower level. In the past three months, how many times have you fallen?  
(n=3483)

Mean number of falls overall: 0.5  
Reported any falls: 20.8%  
Mean number of falls for those with any: 2.5

77 DON'T KNOW/NOT SURE  
99 REFUSED

Q15.2 **[IF C15Q01=1]** Did this fall cause an injury? **[IF C15Q01>1]** How many of these falls caused an injury?  
(n=732)

Mean number of injuries from falls overall: 0.4  
Reported any injury from falls: 29.5%  
Mean number of injuries from falls for those with any: 1.5

88 NONE  
77 DON'T KNOW/NOT SURE  
99 REFUSED

## SECTION 16: SEATBELT USE

Q16.1 How often do you use seat belts when you drive or ride in a car? Would you say ...  
(n=5044)

1 Always (76.5%)  
2 Nearly always (15.1%)  
3 Sometimes (4.8%)  
4 Seldom (1.9%)  
5 Never (1.7%)  
8 Never drive or ride in a car (0.2%)

7 DON'T KNOW/NOT SURE  
9 REFUSED

## SECTION 17: DRINKING AND DRIVING

Q17.1 **[IF Q13.1<>2 AND Q16.1<>8]** During the past 30 days, how many times have you driven when you've had perhaps too much to drink?  
(n=2223)

Mean number of times driven when too much to drink: 0.1  
Reported any driving when too much to drink: 4.3%  
Mean number of times driving when too much to drink for those reporting any: 1.9

88 NONE  
77 DON'T KNOW/NOT SURE  
99 REFUSED

## SECTION 18: WOMEN'S HEALTH

Q18.1 **[IF Q12.20=2]** A mammogram is an x-ray of each breast to look for breast cancer. Have you ever had a mammogram?  
(n=3078)

1 Yes (59.4%)  
2 No (40.6%)

7 DON'T KNOW/NOT SURE  
9 REFUSED

Q18.2 **[IF Q12.20=2 AND Q18.1=1]** How long has it been since you had your last mammogram?  
(n=2218)

1 Within the past year (< 12 months ago) (59.5%)  
2 Within the past 2 years (1 year but < 2 years ago) (16.3%)  
3 Within the past 3 years (2 years but < 3 years ago) (7.3%)  
4 Within the past 5 years (3 years but < 5 years ago) (6.2%)  
5 5 or more years ago (10.8%)

7 DON'T KNOW/NOT SURE  
9 REFUSED

Q18.3 **[IF Q12.20=2]** A clinical breast exam is when a doctor, nurse, or other health professional feels the breasts for lumps. Have you ever had a clinical breast exam?  
(n=3073)

1 Yes (89.4%)  
2 No (10.6%)

7 DON'T KNOW/NOT SURE  
9 REFUSED

Q18.4 **[IF Q12.20=2 AND Q18.3=1]** How long has it been since your last breast exam?  
(n=2807)

1 Within the past year (< 12 months ago) (63.6%)  
2 Within the past 2 years (1 year but < 2 years ago) (15.8%)  
3 Within the past 3 years (2 years but < 3 years ago) (6.1%)  
4 Within the past 5 years (3 years but < 5 years ago) (6.1%)  
5 5 or more years ago (8.4%)

7 DON'T KNOW/NOT SURE  
9 REFUSED

Q18.5 **[IF Q12.20=2]** A Pap test is a test for cancer of the cervix. Have you ever had a Pap test?  
(n=3069)

1 Yes (92.9%)  
2 No (7.1%)

7 DON'T KNOW/NOT SURE  
9 REFUSED

Q18.6 **[IF Q12.20=2 AND Q18.5=1]** How long has it been since you had your last Pap test?  
(n=2919)

1 Within the past year (< 12 months ago) (51.4%)  
2 Within the past 2 years (1 year but < 2 years ago) (16.0%)  
3 Within the past 3 years (2 years but < 3 years ago) (7.4%)  
4 Within the past 5 years (3 years but < 5 years ago) (7.3%)  
5 5 or more years ago (17.9%)

7 DON'T KNOW/NOT SURE  
9 REFUSED

Q18.7 **[IF Q12.20=2 AND Q12.21<=>1]** Have you had a hysterectomy?  
(n=3027)

- 1 Yes (27.1%)
- 2 No (72.9%)
  
- 7 DON'T KNOW/NOT SURE
- 9 REFUSED

#### SECTION 19: PROSTATE CANCER SCREENING

Q19.1 **[IF Q12.20=1 AND Q12.1>= 40]** Now, I will ask you some questions about prostate cancer screening. A Prostate-Specific Antigen test, also called a PSA test, is a blood test used to check men for prostate cancer. Have you ever had a PSA test?  
(n=1469)

- 1 Yes (63.8%)
- 2 No (36.2%)
  
- 7 DON'T KNOW/NOT SURE
- 9 REFUSED

Q19.2 **[IF Q12.20=1 AND Q12.1>= 40 AND Q19.1=1]** How long has it been since you had your last PSA test?  
(n=976)

- 1 Within the past year (< 12 months ago) (66.2%)
- 2 Within the past 2 years (1 year but < 2 years ago) (17.0%)
- 3 Within the past 3 years (2 years but < 3 years ago) (7.2%)
- 4 Within the past 5 years (3 years but < 5 years ago) (4.6%)
- 5 5 or more years ago (5.0%)
  
- 7 DON'T KNOW/NOT SURE
- 9 REFUSED

Q19.3 **[IF Q12.20=1 AND Q12.1>= 40]** A digital rectal exam is an exam in which a doctor, nurse, or other health professional places a gloved finger into the rectum to feel the size, shape, and hardness of the prostate gland. Have you ever had a digital rectal exam?  
(n=1507)

- 1 Yes (75.2%)
- 2 No (24.8%)
  
- 7 DON'T KNOW/NOT SURE
- 9 REFUSED

Q19.4 **[IF Q12.20=1 AND Q12.1>= 40 AND Q19.3=1]** How long has it been since your last digital rectal exam?  
(n=1153)

- 1 Within the past year (< 12 months ago) (48.3%)
- 2 Within the past 2 years (1 year but < 2 years ago) (16.4%)
- 3 Within the past 3 years (2 years but < 3 years ago) (9.4%)
- 4 Within the past 5 years (3 years but < 5 years ago) (9.0%)
- 5 5 or more years ago (16.9%)
  
- 7 DON'T KNOW/NOT SURE
- 9 REFUSED

Q19.5 **[IF Q12.20=1 AND Q12.1>=40]** Have you ever been told by a doctor, nurse, or other health professional that you had prostate cancer?  
(n=1521)

- 1 Yes (4.5%)
- 2 No (95.5%)
  
- 7 DON'T KNOW/NOT SURE
- 9 REFUSED

#### SECTION 20: COLORECTAL CANCER SCREENING

Q20.1 **[IF Q12.1>= 50]** A blood stool test is a test that may use a special kit at home to determine whether the stool contains blood. Have you ever had this test using a home kit?  
(n=2981)

- 1 Yes (38.2%)
- 2 No (61.8%)
  
- 7 DON'T KNOW/NOT SURE
- 9 REFUSED

Q20.2 **[IF Q12.1>= 50 AND Q20.1=1]** How long has it been since you had your last blood stool test using a home kit?  
(n=1159)

- 1 Within the past year (< 12 months ago) (30.9%)
- 2 Within the past 2 years (1 year but < 2 years ago) (18.9%)
- 3 Within the past 3 years (2 years but < 3 years ago) (10.1%)
- 4 Within the past 5 years (3 years but < 5 years ago) (12.8%)
- 5 5 or more years ago (27.2%)
  
- 7 DON'T KNOW/NOT SURE
- 9 REFUSED

Q20.3 **[IF Q12.1>= 50]** Sigmoidoscopy and colonoscopy are exams in which a tube is inserted in the rectum to view the colon for signs of cancer or other health problems. Have you ever had either of these exams?  
(n=2984)

- 1 Yes (55.5%)
- 2 No (44.5%)
  
- 7 DON'T KNOW/NOT SURE
- 9 REFUSED

Q20.4 **[IF Q12.1>= 50 AND Q20.3=1]** For a SIGMOIDOSCOPY, a flexible tube is inserted into the rectum to look for problems. A COLONOSCOPY is similar, but uses a longer tube, and you are usually given medication through a needle in your arm to make you sleepy and told to have someone else drive you home after the test. Was your MOST RECENT exam a sigmoidoscopy or a colonoscopy?  
(n=1645)

- 1 Sigmoidoscopy (10.9%)
- 2 Colonoscopy (89.1%)
  
- 7 DON'T KNOW/NOT SURE
- 9 REFUSED

Q20.5 **[IF Q12.1>= 50 AND Q20.3=1]** How long has it been since you had your last sigmoidoscopy or colonoscopy?  
(n=1687)

- 1 Within the past year (< 12 months ago) (26.0%)
- 2 Within the past 2 years (1 year but < 2 years ago) (21.0%)
- 3 Within the past 3 years (2 years but < 3 years ago) (13.4%)
- 4 Within the past 5 years (3 years but < 5 years ago) (21.6%)
- 5 Within the past 10 years (5 years but < 10 years ago) (12.0%)
- 6 10 or more years ago (6.0%)
  
- 7 DON'T KNOW/NOT SURE
- 9 REFUSED

**SECTION 21: HIV/AIDS**

Q21.1 **[IF Q12.1<65]** The next few questions are about the national health problem of HIV, the virus that causes AIDS. Please remember that your answers are strictly confidential and that you don't have to answer every question if you do not want to. Although we will ask you about testing, we will not ask you about the results of any test you may have had.

Have you ever been tested for HIV? Do not count tests you may have had as part of a blood donation. Include testing fluid from your mouth.

(n=3549)

- 1 Yes (31.2%)
- 2 No (68.8%)

- 7 DON'T KNOW/NOT SURE
- 9 REFUSED

Q21.2 **[IF Q12.1<65 AND Q21.1=1]** Not including blood donations, in what month and year was your last HIV test?

(n=884)

Mode month/year of last HIV test: Unknown month/2007

- 77/7777 DON'T KNOW/NOT SURE
- 99/9999 REFUSED

Q21.3 **[IF Q12.1<65 AND Q21.1=1]** Where did you have your last HIV test - at a private doctor or HMO office, at a counseling and testing site, at a hospital, at a clinic, in a jail or prison, at a drug treatment facility, at home, or somewhere else?

(n=1027)

- 1 Private doctor or HMO (40.8%)
- 2 Counseling and testing site (2.2%)
- 3 Hospital (16.5%)
- 4 Clinic (23.9%)
- 5 Jail or prison (or other correctional facility) (2.3%)
- 6 Drug treatment facility (0.1%)
- 7 At home (4.4%)
- 8 Somewhere else (9.8%)

- 77 DON'T KNOW/NOT SURE
- 99 REFUSED

Q21.4 **[IF Q12.1<65 AND Q21.2=WITHIN LAST 12 MONTHS]** Was it a rapid test where you could get your results within a couple of hours?

(n=161)

- 1 Yes (23.4%)
- 2 No (76.6%)

- 7 DON'T KNOW/NOT SURE
- 9 REFUSED

Q21.5 **[IF Q12.1<65]** I'm going to read you a list. When I'm done, please tell me if any of the situations apply to you. You do not need to tell me which one: you have used intravenous drugs in the past year; you have been treated for a sexually transmitted disease or venereal disease in the past year; you have given or received money or drugs in exchange for sex in the past year; you had anal sex without a condom in the past year. Do any of these situations apply to you?

(n=3631)

- 1 Yes (2.5%)
- 2 No (97.5%)

- 7 DON'T KNOW/NOT SURE
- 9 REFUSED

**SECTION 22: EMOTIONAL SUPPORT AND LIFE SATISFACTION**

Q22.1 The next two questions are about emotional support and your satisfaction with life.

How often do you get the social and emotional support you need? (n=4913)

- 1 Always (49.8%)
- 2 Usually (34.6%)
- 3 Sometimes (10.0%)
- 4 Rarely (2.3%)
- 5 Never (3.4%)

- 7 DON'T KNOW/NOT SURE
- 9 REFUSED

Q22.2 In general, how satisfied are you with your life? (n=4955)

- 1 Very satisfied (47.5%)
- 2 Satisfied (48.1%)
- 3 Dissatisfied (4.0%)
- 4 Very dissatisfied (0.5%)

- 7 DON'T KNOW/NOT SURE
- 9 REFUSED

**MODULE 9: ADULT ASTHMA HISTORY**

M9.1 **[IF Q9.1=1]** Previously you said you were told by a doctor, nurse or other health professional that you had asthma. How old were you when you were first told by a doctor, nurse, or other health professional that you had asthma?

(n=606)

Mean age in years (older than 10 years): 32.1

- 97 AGE 10 OR YOUNGER (0.0%)

- 98 DON'T KNOW/NOT SURE
- 99 REFUSED

M9.2 **[IF Q9.2=1]** During the past 12 months, have you had an episode of asthma or an asthma attack?

(n=461)

- 1 Yes (53.7%)
- 2 No (46.3%)

- 7 DON'T KNOW/NOT SURE
- 9 REFUSED

M9.3 **[IF Q9.2=1 AND M9.2=1]** During the past 12 months, how many times did you visit an emergency room or urgent care center because of your asthma?

(n=227)

Mean number of emergency/urgent care visits overall: 0.5

Reported any emergency/urgent care visit: 24.7%

Mean number of visits for those with any: 2.2

- 88 NONE
- 98 DON'T KNOW/NOT SURE
- 99 REFUSED

M9.4 **[IF Q9.2=1 AND IF M9.3>0 AND<88]** Besides those emergency room or urgent care visits, during the past 12 months, how many times did you see a doctor, nurse or other health professional for urgent treatment of worsening asthma symptoms?

(n=227)

Mean number of times had urgent treatment overall: 0.6

Reported any urgent treatment: 28.2%

Mean number of times for those with any: 2.0

- 88 NONE
- 98 DON'T KNOW/NOT SURE
- 99 REFUSED

M9.5 **[IF Q9.2=1]** During the past 12 months, how many times did you see a doctor, nurse, or other health professional for a routine checkup for your asthma?  
(n=459)

Mean number of times saw health professional overall: 0.8  
Reported any times saw health professional: 49.0%  
Mean number of times for those who saw any: 1.8

88 NONE  
98 DON'T KNOW/NOT SURE  
99 REFUSED

M9.6 **[IF Q9.2=1]** During the past 12 months, how many days were you unable to work or carry out your usual activities because of your asthma?  
(n=451)

Mean number of days asthma-limited activity overall: 3.1  
Reported any days asthma-limited activity: 20.9%  
Mean number of days for those with any: 15.0

888 NONE  
777 DON'T KNOW/NOT SURE  
999 REFUSED

M9.7 **[IF Q9.2=1]** Symptoms of asthma include cough, wheezing, shortness of breath, chest tightness and phlegm production when you don't have a cold or respiratory infection. During the past 30 days, how often did you have any symptoms of asthma?  
(n=450)

1 Less than once a week (25.3%)  
2 Once or twice a week (20.7%)  
3 More that 2 times a week, but not every day (7.9%)  
4 Every day, but not all the time (10.5%)  
5 Every day, all the time (6.1%)  
6 Not at any time (29.5%)

7 DON'T KNOW/NOT SURE  
9 REFUSED

M9.8 **[IF Q9.2=1 AND M9.7<=8]** During the past 30 days, how many days did symptoms of asthma make it difficult for you to stay asleep? Would you say ...  
(n=325)

1 One or two (17.4%)  
2 Three to four (7.4%)  
3 Five (2.8%)  
4 Six to ten (7.0%)  
5 More than ten (4.1%)  
6 None (61.3%)

7 DON'T KNOW/NOT SURE  
9 REFUSED

M9.9 **[IF Q9.2=1]** During the past 30 days, how many days did you take a prescription asthma medication to PREVENT an asthma attack from occurring?  
(n=454)

1 1 to 14 days (21.1%)  
2 15 to 24 days (3.4%)  
3 25 to 30 days (27.2%)  
8 Never (48.3%)

7 DON'T KNOW/NOT SURE  
9 REFUSED

M9.10 **[IF Q9.2=1]** During the past 30 days, how often did you use a prescription asthma inhaler DURING AN ASTHMA ATTACK to stop it?  
(n=450)

1 1 to 4 times (in the past 30 days) (27.3%)  
2 5 to 14 times (in the past 30 days) (8.5%)  
3 15 to 29 times (in the past 30 days) (2.8%)  
4 30 to 59 times (in the past 30 days) (3.0%)  
5 60 to 99 times (in the past 30 days) (1.5%)  
6 100 or more times (in the past 30 days) (0.6%)  
7 Never (include no attack in past 30 days) (56.3%)

7 DON'T KNOW/NOT SURE  
9 REFUSED

### MODULE 13: ANXIETY AND DEPRESSION

M13.1 Over the last 2 weeks, how many days have you had little interest or pleasure in doing things?  
(n=4813)

Mean number of days little interest or pleasure overall: 1.5  
Reported any days with little interest or pleasure: 33.8%  
Mean number of days for those with any: 4.3

88 NONE  
77 DON'T KNOW/NOT SURE  
99 REFUSED

M13.2 Over the last 2 weeks, how many days have you felt down, depressed OR hopeless?  
(n=4898)

Mean number of days down, etc. overall: 1.1  
Reported any days down/depressed/hopeless: 29.2%  
Mean number of days for those with any: 3.8

88 NONE  
77 DON'T KNOW/NOT SURE  
99 REFUSED

M13.3 Over the last 2 weeks, how many days have you had trouble falling asleep OR staying asleep OR sleeping too much?  
(n=4874)

Mean number of days with sleeping problems overall: 2.9  
Reported any days with sleeping problems: 50.7%  
Mean number of days for those with any: 5.8

88 NONE  
77 DON'T KNOW/NOT SURE  
99 REFUSED

M13.4 Over the last 2 weeks, how many days have you felt tired OR had little energy?  
(n=4852)

Mean number of days tired or little energy overall: 3.7  
Reported any days tired or little energy: 68.3%  
Mean number of days for those with any: 5.4

88 NONE  
77 DON'T KNOW/NOT SURE  
99 REFUSED

M13.5 Over the last 2 weeks, how many days have you had a poor appetite OR eaten too much?  
(n=4857)

Mean number of days poor/excessive appetite overall: 2.0  
Reported any days poor/excessive appetite: 38.6%  
Mean number of days for those with any: 5.1

88 NONE  
77 DON'T KNOW/NOT SURE  
99 REFUSED

M13.6 Over the last 2 weeks, how many days have you felt bad about yourself OR that you were a failure or had let yourself or your family down?

(n=4880)

Mean number of days felt bad, etc. overall: 0.9  
Reported any days felt bad/failure/let others down: 19.8%  
Mean number of days for those with any: 4.6

88 NONE  
77 DON'T KNOW/NOT SURE  
99 REFUSED

M13.7 Over the last 2 weeks, how many days have you had trouble concentrating on things, such as reading the newspaper OR watching the T.V.?

(n=4877)

Mean number of days with trouble concentrating overall: 1.0  
Reported any days with trouble concentrating: 19.3%  
Mean number of days for those with any: 5.1

88 NONE  
77 DON'T KNOW/NOT SURE  
99 REFUSED

M13.8 Over the last 2 weeks, how many days have you moved or spoken so slowly that other people could have noticed? OR THE OPPOSITE - being so fidgety or restless that you were moving around a lot more than usual?

(n=4824)

Mean number of days unusually slow or restless overall: 0.5  
Reported any days unusually slow or restless: 9.9%  
Mean number of days for those with any: 5.1

88 NONE  
77 DON'T KNOW/NOT SURE  
99 REFUSED

M13.9 Has a doctor or other healthcare provider EVER told you that you had an anxiety disorder (including acute stress disorder, anxiety, generalized anxiety disorder, obsessive-compulsive disorder, panic disorder, phobia, posttraumatic stress disorder, or social anxiety disorder)?

(n=4896)

1 Yes (11.6%)  
2 No (88.4%)

7 DON'T KNOW/NOT SURE  
9 REFUSED

M13.10 Has a doctor or other healthcare provider EVER told you that you have a depressive disorder (including depression, major depression, dysthymia, or minor depression)?

(n=4893)

1 Yes (18.9%)  
2 No (81.2%)

7 DON'T KNOW/NOT SURE  
9 REFUSED

#### MODULE 15: RANDOM CHILD SELECTION

M15.1 **[IF Q12.7>=1 AND Q12.7<88 RANDOMLY SELECT ONE OF THE CHILDREN]** What is the birth month and year of the **[SELECTED]** child?

(Due to the size of the table, these data are not printed.)

M15.2 **[IF Q12.7<88]** Is the child a boy or a girl?  
(n=1619)

1 Boy (51.2%)  
2 Girl (48.8%)

9 REFUSED

M15.3 **[IF Q12.7<88]** Is the child Hispanic or Latino?  
(n=1623)

1 Yes (12.3%)  
2 No (87.7%)

7 DON'T KNOW/NOT SURE  
9 REFUSED

M15.4 **[IF Q12.7<88]** Which one or more of the follow would you say is the race of the child?  
(n=1656)

[SEE RESULTS FOLLOWING M15.5]

M15.5 **[IF Q12.7<88]** Which one of these groups would you say best represents the child's race?  
(n=1608)

1 White (92.1%)  
2 Black or African American (1.4%)  
3 Asian (0.6%)  
4 Native Hawaiian or Other Pacific Islander (0.6%)  
5 American Indian or Alaskan Native (1.7%)  
6 Other (3.8%)

7 DON'T KNOW/NOT SURE  
9 REFUSED

M15.6 **[IF Q12.7<88]** How are you related to the child?  
(n=1634)

1 Parent (include biologic, step, or adoptive parent) (91.4%)  
2 Grandparent (4.0%)  
3 Foster parent or guardian (1.0%)  
4 Sibling (include biologic, step, and adoptive sibling) (2.1%)  
5 Other relative (0.7%)  
6 Not related in any way (0.9%)

7 DON'T KNOW/NOT SURE  
9 REFUSED

#### STATE ADDED 1: CHILD SAFETY

ID1.1 **[IF Q12.7<>88 OR 99 AND CHILDAGE<16]** How often does the **[CHILDAGE]**-month/year old child in your household use a **[IF CHILDAGE<5 "car safety seat", IF CHILDAGE>=5 "seatbelt"]** when they ride in a car?  
(n=1276)

1 Always (90.2%)  
2 Nearly always (8.2%)  
3 Sometimes (1.2%)  
4 Seldom (0.2%)  
5 Never (0.2%)

7 DON'T KNOW/NOT SURE  
9 REFUSED

**MODULE 16: CHILDHOOD ASTHMA PREVALENCE**

M16.1 **[IF Q12.7<88]** Has a doctor or other health professional ever said that the child has asthma?  
(n=1623)

- 1 Yes (8.8%)
- 2 No (91.2%)
  
- 7 DON'T KNOW/NOT SURE
- 9 REFUSED

M16.2 **[IF Q12.7<88 AND M16.1=1]** Does the child still have asthma?  
(n=138)

- 1 Yes (75.8%)
- 2 No (24.2%)
  
- 7 DON'T KNOW/NOT SURE
- 9 REFUSED

**STATE ADDED 2: HEALTH CARE COVERAGE**

ID2.1 **[IF Q3.1=2]** Previously you said you did not have any health care coverage. What is the main reason you are without health care coverage?  
(n=610)

- 1 Lost job or changed employers (10.8%)
- 2 Spouse/parent lost job/changed employers (0.6%)
- 3 Became divorced or separated (0.6%)
- 4 Became ineligible because of age or left school (0.5%)
- 5 Employer doesn't offer or stopped offering coverage (10.8%)
- 6 Cut back to part time or became temporary employee (1.0%)
- 7 Benefits from employer or former employer ran out (2.6%)
- 8 Couldn't afford to pay the premiums (49.3%)
- 9 Insurance company refused coverage (1.4%)
- 10 Lost Medicaid or Medical Assistance eligibility (2.4%)
- 11 Other reason (20.1%)
  
- 77 DON'T KNOW/NOT SURE
- 99 REFUSED

ID2.2 **[IF Q3.1=2]** About how long has it been since you had health care coverage?  
(n=609)

- 1 Within the past 6 months (14.1%)
- 2 Within the past year (14.4%)
- 3 Within the past 2 years (11.5%)
- 4 Within the past 5 years (14.0%)
- 5 5 or more years ago (29.9%)
- 6 NEVER (16.2%)
  
- 7 DON'T KNOW/NOT SURE
- 9 REFUSED

ID2.3 **[IF Q3.1=1]** Previously you said that you had some kind of health care coverage. What type of health care coverage do you use to pay for most of your medical care?  
(n=4152)

- 1 Employer (39.1%)
- 2 Someone else's employer (19.2%)
- 3 A plan bought or someone else buys (12.6%)
- 4 Medicare (17.8%)
- 5 Medicaid or Medical Assistance (3.7%)
- 6 The military, CHAMPUS, Tricare, or the VA (4.1%)
- 7 The Indian Health Service (0.2%)
- 8 Some other source (3.4%)
- 88 NONE (0.1%)
  
- 77 DON'T KNOW/NOT SURE
- 99 REFUSED

ID2.4 Do you have any kind of insurance coverage that pays for some or all of your routine dental care, including dental insurance, prepaid plans such as HMOs, or government plans such as Medicaid?  
(n=4814)

- 1 Yes (57.0%)
- 2 No (43.0%)
  
- 7 DON'T KNOW/NOT SURE
- 9 REFUSED

**STATE ADDED 3: SEX EDUCATION**

ID3.1 At what grade level do you think children should begin to receive comprehensive sex education in school? Comprehensive sex education includes not only discussions about reproductive systems and associated biological functions, but also sexual behavior; outcomes of sexual behavior including STD/AIDS education and prevention; talking with parents about sex; birth control methods, availability, and usage; and may include discussion on personal sexuality choices.  
(n=4340)

- 55 Kindergarten (0.8%)
- 1 1st (1.3%)
- 2 2nd (0.7%)
- 3 3rd (2.5%)
- 4 4th (7.7%)
- 5 5th (17.9%)
- 6 6th (26.2%)
- 7 7th (16.6%)
- 8 8th (8.8%)
- 9 9th (5.2%)
- 10 10th (2.5%)
- 11 11th (0.6%)
- 12 12th (2.2%)
  
- 88 NEVER (7.3%)
- 77 DON'T KNOW/NOT SURE
- 99 REFUSED

**STATE ADDED 4: HOME WATER SUPPLY**

ID4.1 What is the main source of your home water supply?  
(n=4793)

- 1 A city, county or town water system (65.6%)
- 2 A small water system operated by a home association (3.4%)
- 3 A private well serving your home (28.7%)
- 4 Other source (2.3%)
  
- 7 DON'T KNOW/NOT SURE
- 9 REFUSED

ID4.2 How is the sewage or wastewater disposed of for your home?  
(n=4692)

- 1 A city, county or town sewer system (63.9%)
- 2 A small sewer district or association (1.8%)
- 3 A private sewer system or septic tank for your home (33.7%)
- 4 Other source (0.6%)
  
- 7 DON'T KNOW/NOT SURE
- 9 REFUSED

**STATE ADDED 5: HOME ENVIRONMENT**

ID5.1 During the past 12 months, on how many days were pesticides, sprays, or chemicals applied inside your home to kill bugs, mice, or other pests?  
(n=4748)

- Mean number of days: 1.1
  
- 888 NONE
- 777 DON'T KNOW/NOT SURE
- 999 REFUSED

**STATE ADDED 6: SUN BURNS**

ID6.1 The next question is about sun burns, including anytime that even a small part of your skin was red for more than 12 hours. Have you had a sun burn within the past 12 months?  
(n=4815)

- 1 Yes (48.4%)
- 2 No (51.6%)
  
- 7 DON'T KNOW/NOT SURE
- 9 REFUSED

ID6.2 **[IF ID6.1=1]** Including times when even a small part of your skin was red for more than 12 hours, how many sunburns have you had within the past 12 months?  
(n=1891)

- 1 One (27.7%)
- 2 Two (27.1%)
- 3 Three (17.5%)
- 4 Four (6.9%)
- 5 Five (4.9%)
- 6 Six or more (16.0%)
  
- 7 DON'T KNOW/NOT SURE
- 9 REFUSED

**STATE ADDED 7: FISH**

ID7.1 How often do you eat fish? Be sure to include canned tuna and canned salmon, imitation crab and fish eaten at restaurants. Do not include shellfish such as shrimp, oysters, clams or real crab?  
(n=4754)

Mean times eat fish per year: 54.4

- 555 NEVER
- 777 DON'T KNOW/NOT SURE
- 999 REFUSED

ID7.2 **[IF ID07Q01<->555, 777 OR 999]** How often do you eat fish that has been caught in Idaho waters? This includes lakes, reservoirs, rivers, creeks, ponds.  
(n=4153)

Mean times eat fish caught in Idaho per year: 8.0

- 555 NEVER
- 777 DON'T KNOW/NOT SURE
- 999 REFUSED

**STATE ADDED 8: NON-MEDICAL DRUG USE**

ID8.1 I am now going to ask about your non-medical use of drugs. "Non-medical use" means using drugs not prescribed by a doctor, or used to get high or for curiosity. I will be referring to cocaine (including crack), heroin or drugs like heroin (such as codeine or Demerol), marijuana (also referred to as pot or hashish), methamphetamine (also known as meth, crank or ice), hallucinogens, inhalants, prescription painkillers, stimulants and sedatives. Remember, all information on this survey is strictly confidential. Have you ever used any of the drugs I just mentioned?  
(n=4804)

- 1 Yes (23.2%)
- 2 No (76.8%)
  
- 7 DON'T KNOW/NOT SURE
- 9 REFUSED

ID8.2 **[IF ID8.1=1]** How long has it been since you last used any of these drugs for non-medical purposes? Would you say ...  
(n=916)

- 1 Within the last 30 days (0-30 DAYS) (8.4%)
- 2 More than 30 days but within the past 12 months (9.2%)
- 3 More than 12 months ago (82.4%)
  
- 7 DON'T KNOW/NOT SURE
- 9 REFUSED

ID8.3 **[IF ID8.1=1 AND ID8.2<3]** Used marijuana in past 12 months?  
(n=114)

- 1 Yes (91.3%)
- 2 No (8.7%)
  
- 7 DON'T KNOW/NOT SURE
- 9 REFUSED

ID8.4 **[IF ID8.1=1 AND ID8.2<3 AND ID8.3<->9]** Used meth in past 12 months?  
(n=115)

- 1 Yes (9.1%)
- 2 No (90.9%)
  
- 7 DON'T KNOW/NOT SURE
- 9 REFUSED

ID8.5 **[IF ID8.1=1 AND ID8.2<3 AND ID8.3<->9]** Used cocaine in past 12 months?  
(n=115)

- 1 Yes (8.0%)
- 2 No (92.0%)
  
- 7 DON'T KNOW/NOT SURE
- 9 REFUSED

ID8.6 **[IF ID8.1=1 AND ID8.2<3 AND ID8.3<->9]** Used heroin in past 12 months?  
(n=115)

- 1 Yes (0.3%)
- 2 No (99.7%)
  
- 7 DON'T KNOW/NOT SURE
- 9 REFUSED

ID8.7 **[IF ID8.1=1 AND ID8.2<3 AND ID8.3<->9]** Used prescription pain killers in past 12 months?  
(n=115)

- 1 Yes (28.9%)
- 2 No (71.1%)
  
- 7 DON'T KNOW/NOT SURE
- 9 REFUSED

ID8.8 **[IF ID8.1=1 AND ID8.2<3 AND ID8.3<->9]** Other than those I just mentioned, have you used any other drugs for non-medical purposes in the past 12 months?  
(n=114)

- 1 Yes (14.8%)
- 2 No (85.2%)
  
- 7 DON'T KNOW/NOT SURE
- 9 REFUSED

## STATE ADDED 9: INTIMATE PARTNER VIOLENCE

ID9.1 The next questions are about different types of violence in relationships with an intimate partner. By an intimate partner I mean any current or former spouse, boyfriend, or girlfriend. Someone you were dating, or romantically or sexually intimate with would also be considered an intimate partner. This information will help us to better understand the problem of violence in relationships. This is a sensitive topic. Some people may feel uncomfortable with these questions. At the end of this section, I will give you a phone number of an organization that can provide information and referral for these issues. Please keep in mind that if you are not in a safe place you can ask me to skip any question you do not want to answer.

Are you in a safe place to answer these questions?  
(n=4821)

- 1 Yes (98.0%)
- 2 No (2.0%)

ID9.2 **[IF ID9.0I=1]** Has an intimate partner EVER THREATENED you with physical violence? This includes threatening to hit, slap, push, kick, or physically hurt you in any way.  
(n=4691)

- 1 Yes (13.4%)
- 2 No (86.6%)
  
- 7 DON'T KNOW/NOT SURE
- 9 REFUSED

D9.3 **[IF ID9.0I=1]** Has an intimate partner EVER hit, slapped, pushed, kicked, or physically hurt you in any way?  
(n=4689)

- 1 Yes (15.7%)
- 2 No (84.4%)
  
- 7 DON'T KNOW/NOT SURE
- 9 REFUSED

ID9.4 **[IF ID9.0I=1]** Other than what you have already told me about, has an intimate partner EVER ATTEMPTED physical violence against you? This includes times when they tried to hit, slap, push, kick, or otherwise physically hurt you, but they were not able to?  
(n=4670)

- 1 Yes (11.5%)
- 2 No (88.5%)
  
- 7 DON'T KNOW/NOT SURE
- 9 REFUSED



# Appendix C

---

## References

Federal Communications Commission, Wireline Competition Bureau, Industry Analysis and Technology Division. *Statistics of Communications Common Carriers, 2005/2006 Edition*, Washington DC; Federal Communications Commission, 2008. [www.fcc.gov/wcb/iatd/socc.html](http://www.fcc.gov/wcb/iatd/socc.html) Accessed August 21, 2009.

Industry Analysis and Technology Division, Wireline Competition Bureau, *Telephone Subscribership in the United States, (Based on data from the Census Bureau's Current Population Survey)*, January 2007.

Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, *Prevalence and Trends Data: BRFSS 2008* <http://apps.nccd.cdc.gov/brfss/> Accessed July 22, 2009

National Cancer Institute, U.S. National Institutes of Health. *Common Cancer Types, Screening and Testing* [www.cancer.gov](http://www.cancer.gov) Accessed August 7, 2009.

American Cancer Society Guidelines for the Early Detection of Cancer  
[http://www.cancer.org/docroot/PED/ped\\_0.asp](http://www.cancer.org/docroot/PED/ped_0.asp) Accessed July 21, 2009