

P

R

A

T

S

**2010
ANNUAL
REPORT**

P

R

A

T

S

**2010
ANNUAL
REPORT**



Acknowledgments

This project was funded by grants through the State Systems Development Initiative (SSDI) Grant Program and the Title V MCH Services Block Grant Program (Maternal and Child Health Bureau, Health Resources and Services Administration, U.S. Department of Health and Human Services). Collaboration between Idaho's SSDI Program and Title V MCH Program has been essential to the success in establishing PRATS as an on-going data system.

The Idaho Department of Health and Welfare would like to thank the mothers of Idaho who have participated in the Pregnancy Risk Assessment Tracking System.

For more details on this project or any of the survey results, please contact the PRATS Program Director, Idaho Bureau of Vital Records and Health Statistics, at (208)332-7366.

Suggested Citation

Results From the 2010 Pregnancy Risk Assessment Tracking System (PRATS) Annual Report. Boise: Idaho Department of Health and Welfare, Division of Public Health, Bureau of Vital Records and Health Statistics, 2013.

Table of Contents

Title Page	i
Acknowledgments	ii
Table of Contents	iii
Overview	1
What is the Pregnancy Risk Assessment Tracking System (PRATS)?	1
Methodology	1
Reporting the Data	2
Data Limitations.....	3
References	3
Definition of Terms	3
Idaho Public Health Districts	4
Demographic Profile	5
Data Tables	
Understanding the Data Tables.....	7
Data Tables by Health District, 2010.....	8
Appendices	
Appendix A: Maternal and Infant Health Trends by Health District, 2004-2010	55
Appendix B: PRATS Questionnaire and Results, 2010	65
Appendix C: Technical Notes	83

Overview

What is the Pregnancy Risk Assessment Tracking System?

The Pregnancy Risk Assessment Tracking System (PRATS) is an annual survey of new mothers in Idaho, directed by the Bureau of Vital Records and Health Statistics within the Idaho Department of Health and Welfare. PRATS was first conducted in Idaho in 1999, and has been conducted annually since 2001. PRATS is modeled after the Centers for Disease Control and Prevention's (CDC) Pregnancy Risk Assessment Monitoring System (PRAMS).¹

PRATS is a population-based tracking system that identifies and examines maternal experiences and health practices surrounding pregnancy which may affect pregnancy outcomes and infant health. PRATS data are supplemental to information obtained from vital records and are used for planning, developing, and assessing perinatal health programs in Idaho. Results from PRATS have been used to determine progress towards goals, assess trends, and target at-risk populations relevant to maternal and child health. PRATS provides information about the intendedness of pregnancy, initiation of prenatal care, content of prenatal care, barriers to services, physical abuse among pregnant women, breastfeeding patterns, and many other important perinatal issues. This report summarizes major maternal health and behavior risk factors associated with maternal, fetal, and infant health outcomes. Results presented in this report are not inclusive of all data that may be provided by PRATS. Therefore, the 2010 PRATS questionnaire appears in its entirety at the end of this report in Appendix B (see page 88).

Methodology

Sample

The study population for PRATS included female residents of Idaho aged 18 or older (at the time of delivery) who had a live birth which occurred in-state (based on birth certificate data). The sampling frame included mothers who gave birth between January 1, 2010 and December 31, 2010. During the data collection period in 2011, babies were between the ages of 5 and 20 months.

Certain records were automatically excluded from the sampling frame, including records of mothers less than 18 years of age at the time of delivery, adopted infants, siblings of adopted infants within a multiple birth (twin, triplet, etc.), infants who had died, and siblings of infants who had died within a multiple birth. Also excluded from the sampling frame were Idaho resident mothers who delivered in another state or died after giving birth. If there was a multiple birth, only the firstborn infant was included in the sampling frame.

The sample design of PRATS was based on a disproportionate stratified systematic random sampling method designed to ensure representation of women from each of Idaho's seven public health districts.

Survey Methods

Between May and September 2011, 4,200 randomly selected mothers from across the state of Idaho were mailed an introductory letter requesting their participation in the PRATS survey. The introductory letter explained the purpose of the survey and provided a toll-free number to call for more information or to request a telephone interview. The mothers were also given the opportunity to decline participation in the study.

Approximately two weeks after the introductory letter was mailed, a full questionnaire packet was sent. Hispanic mothers were mailed both an English and Spanish version of the survey. In order to give women every opportunity to complete the questionnaire, up to two more questionnaire packets were mailed out during the course of a two-month period. Women were able to elect to complete the survey by telephone with a trained interviewer (English or Spanish). For women who did not respond, attempts were made to contact them by telephone. This survey strategy was developed by the CDC PRAMS project and proved to be very successful in achieving higher response rates.

Weighting the Data

Data in this report were weighted to adjust for the disproportionate stratified sampling design and response rate differences. For example, mothers in health districts with smaller populations are more likely to be chosen to respond to the survey than mothers in health districts with larger populations. Weighting procedures correct for this difference and ensure that sample data closely reflect the population from which the sample was obtained.

Analyzing the Data

Statistical testing and calculation of confidence limits were done using SUDAAN® (Survey Data Analysis) software. This software takes into account the complex sampling design methodology of PRATS. Data manipulation and variable creation were done using SAS® software. All missing (blank and “don’t know”) observations were excluded from analysis.

Reporting the Data

For purposes of this report, the terms “mothers” or “women” are used to refer to Idaho resident adult mothers (18 and older) who gave birth (live birth) in Idaho. PRATS data are only representative of Idaho resident women aged 18 and older whose pregnancies resulted in a live birth occurring in Idaho. Therefore, PRATS data are not generalizable to all pregnant women.

Prevalence estimates based on denominators with fewer than 30 respondents have been suppressed and are indicated in the data tables with an asterisk (*). PRATS has adopted this standard to maintain a high degree of reliability. Differences between estimates are reported throughout the report. Any difference determined to be

statistically significantly different through statistical testing has been designated using the terms “statistically” or “significantly.”

Data Limitations

The self-reported nature of PRATS data allows for potential information bias and error in estimation. Women may not have been able to accurately recall events surrounding their pregnancies because the timing of the survey was several months following delivery. In addition, women may be hesitant to report behaviors perceived to be risky or socially undesirable. The result of this bias would likely be an underestimation of the actual prevalence of these behaviors. Also, confidentiality concerns may have impacted how likely women were to honestly respond to survey questions. Because women were given the option of responding to the survey by mail or by phone, differences in interpretation may have existed between those who were read the question by a trained interviewer versus reading the question themselves.

References

1. *Pregnancy Risk Assessment Monitoring System (PRAMS)*, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Division of Reproductive Health, See website at: www.cdc.gov/PRAMS

Definition of Terms

AGE OF MOTHER

Age of mother at time of delivery is a calculated field from the birth certificate based on mother's date of birth and birth date of infant.

ETHNICITY OF MOTHER

Mother's ethnicity is based on the “mother of Hispanic origin” question on the birth certificate.


MARITAL STATUS OF MOTHER

Marital status indicates whether the mother was married at the time of conception, at the time of delivery, or at any time between conception and delivery. The marital status field must be completed on the Idaho Certificate of Live Birth.

Idaho Public Health Districts

Idaho is divided into seven public health districts, with each of the districts containing four to eight counties. The seven health districts are independent agencies governed by a local board appointed by the county commissioners within the district.

Number of Idaho Resident Mothers Aged 18 and Older Who Gave Birth In Idaho 2010



Overall:	21,144
District 1:	2,089
District 2:	1,017
District 3:	3,497
District 4:	5,394
District 5:	2,812
District 6:	2,520
District 7:	3,815

Note: Population counts were obtained from Idaho birth records and were used in weighting PRATS data (see page 2). Certain records were excluded such as mothers less than 18 years old at delivery, adopted infants, infant deaths, maternal deaths, non-resident mothers, and out-of-state births (see page 1).

Demographic Profile

The following table provides a demographic profile of Idaho adult mothers by public health district for 2010.

**Demographic Characteristics of Idaho Mothers By Health District
PRATS 2010**

	Idaho	Public Health District						
		1	2	3	4	5	6	7
AGE								
18-19	3.9%	6.6%	4.8%	4.4%	2.1%	4.8%	3.8%	3.5%
20-24	26.6	26.1	23.8	31.1	20.4	26.2	27.1	32.2
25-29	35.9	31.0	40.8	29.6	37.6	35.3	37.3	40.1
30-34	22.7	24.5	20.2	21.3	26.8	21.8	22.5	18.6
35+	10.9	11.8	10.5	13.5	13.1	12.0	9.3	5.5
ETHNICITY								
Non-Hispanic	87.1	96.2	96.8	75.1	90.3	75.0	89.2	93.6
Hispanic	12.9	3.8	3.2	24.9	9.7	25.0	10.8	6.4
MARITAL STATUS								
Not Married	24.8	32.8	27.7	30.1	22.1	30.9	23.0	15.4
Married	75.2	67.2	72.3	69.9	77.9	69.1	77.0	84.6
INCOME								
Less than \$15,000	24.5	23.0	25.7	31.3	18.5	24.2	27.4	25.6
\$15,000 - \$24,999	17.5	17.6	13.6	14.3	16.3	23.8	20.5	16.6
\$25,000 - \$34,999	14.8	17.3	15.4	15.2	12.7	16.0	15.1	14.6
\$35,000 - \$49,999	13.8	12.3	16.5	10.3	13.2	15.4	13.8	16.5
\$50,000+	29.5	29.9	28.8	28.8	39.5	20.6	23.2	26.7
EDUCATION								
K-11th Grade	9.7	6.5	3.6	16.3	6.8	19.1	8.1	5.3
High School or GED	22.6	24.6	28.9	25.3	19.3	21.9	29.7	18.0
Some College	39.4	43.6	34.8	37.6	38.2	40.7	37.0	42.1
College Graduate+	28.4	25.3	32.7	20.8	35.8	18.3	25.2	34.7

Note: The percentages are based on weighted PRATS data (see the *Weighting the Data* section on page 2 for further explanation).



PRATS

DATA TABLES

Understanding the Data Tables

The data tables on the following pages provide both state and health district data for various demographic subcategories. An understanding of the tables is important to interpreting the data correctly.

- A. This title refers to the risk factor.
- B. This title refers to the population surveyed. For all questions, this will be Idaho resident adult mothers (aged 18 and older) who had a live birth in Idaho in 2010. Certain groups may be excluded from analysis, such as birth control use being asked only of women who were not trying to get pregnant.
- C. The labels across the top of the table (i.e., Statewide and Public Health District) refer to the geographic region of residence.
- D. The labels on the left side of the table refer to the particular group the numbers represent. For example, the row labeled "20-24" will contain data for mothers in that age group for the geographic region indicated at the top of the table.
- E. The shaded columns contain the prevalence (percent) of the risk factor.
- F. The two numbers in the column labeled "95% CI" are the lower and upper limits of the confidence interval. This interval can be interpreted to mean that there is a 95% certainty that the true prevalence of the risk factor falls within the confidence interval.
- G. "n" refers to the number of people sampled who responded to the question within the demographic group and geographic region indicated. It does NOT represent the number of people who have the risk factor in the population. An asterisk indicates that fewer than 30 people in the group responded (see the *Reporting the Data* section on page 2).

Unintended Pregnancy A																
Idaho mothers who reported their pregnancy was unintended, 2010 B																
	Public Health District															
	Statewide C				District 1			District 2			District 3					
	%	95% CI		n	%	95% CI		n	%	95% CI		n	%	95% CI		n
TOTAL	33.2	31.2	35.3	2,357	34.2	29.3	39.4	321	31.9	27.8	36.3	340	33.0	27.6	38.9	272
AGE	E	F	G													
18-19	60.8	49.4	71.2	80	*	*	*	*	*	*	*	*	*	*	*	*
20-24 D	45.7	41.5	49.9	594	49.5	38.9	60.2	75	53.0	43.9	62.0	79	48.4	37.8	59.3	79
25-29	28.8	25.6	32.2	872	24.5	17.4	33.3	105	19.6	14.6	25.7	140	24.3	16.1	35.0	84
30-34	23.9	20.3	27.9	550	25.5	17.6	35.5	84	26.7	18.9	36.2	70	20.7	12.6	32.2	62
35+	27.0	21.6	33.0	261	32.9	20.9	47.7	41	21.3	12.4	34.0	38	30.1	17.7	46.2	37

Unintended Pregnancy

Idaho mothers who reported their pregnancy was unintended, 2010

	Statewide			Public Health District								
				District 1			District 2			District 3		
	%	95% CI	n	%	95% CI	n	%	95% CI	n	%	95% CI	n
TOTAL	33.2	31.2 35.3	2,357	34.2	29.3 39.4	321	31.9	27.8 36.3	340	33.0	27.6 38.9	272
AGE												
18-19	60.8	49.4 71.2	80	*	* *	*	*	* *	*	*	* *	*
20-24	45.7	41.5 49.9	594	49.5	38.9 60.2	75	53.0	43.9 62.0	79	48.4	37.8 59.3	79
25-29	28.8	25.6 32.2	872	24.5	17.4 33.3	105	19.6	14.6 25.7	140	24.3	16.1 35.0	84
30-34	23.9	20.3 27.9	550	25.5	17.6 35.5	84	26.7	18.9 36.2	70	20.7	12.6 32.2	62
35+	27.0	21.6 33.0	261	32.9	20.9 47.7	41	21.3	12.4 34.0	38	30.1	17.7 46.2	37
ETHNICITY												
Non-Hispanic	32.0	29.9 34.2	2,107	34.2	29.2 39.6	307	31.2	27.0 35.6	329	29.5	23.7 36.2	208
Hispanic	41.8	35.7 48.3	247	*	* *	*	*	* *	*	43.3	31.9 55.6	64
MARITAL STATUS												
Not Married	60.8	55.9 65.5	407	57.1	46.3 67.4	70	53.9	44.0 63.4	65	55.2	42.8 66.9	58
Married	24.1	22.2 26.2	1,950	23.1	18.7 28.3	251	23.6	19.8 28.0	275	23.4	18.4 29.3	214
INCOME												
Less than \$15,000	51.4	46.8 56.0	495	52.1	40.1 63.8	59	49.0	39.6 58.5	74	49.0	37.9 60.3	72
\$15,000 - \$24,999	41.0	36.0 46.2	388	49.3	36.6 62.1	52	41.6	29.8 54.4	43	45.5	31.0 60.9	39
\$25,000 - \$34,999	36.8	31.5 42.3	352	35.8	24.8 48.5	56	31.3	21.9 42.7	52	39.8	26.2 55.3	41
\$35,000 - \$49,999	23.1	18.6 28.4	350	15.5	8.0 27.9	43	18.1	11.1 27.9	59	*	* *	*
\$50,000+	16.2	13.5 19.3	705	18.5	12.3 26.8	105	22.0	16.2 29.2	105	11.2	6.0 19.9	82
EDUCATION												
K-11th Grade	45.9	38.6 53.4	184	*	* *	*	*	* *	*	55.5	40.0 69.9	38
High School or GED	42.0	37.5 46.6	519	43.1	32.8 54.2	74	40.4	32.2 49.3	89	36.2	25.6 48.3	66
Some College	35.9	32.7 39.2	934	40.8	33.0 49.0	135	38.3	31.2 46.0	117	31.4	23.2 40.9	104
College Graduate+	18.3	15.3 21.6	714	12.5	7.3 20.4	92	15.6	11.0 21.7	123	13.8	7.2 24.9	61

*Figure not reliable by PRATS standards (n < 30)

Unintended Pregnancy

Idaho mothers who reported their pregnancy was unintended, 2010

	Public Health District															
	District 4				District 5				District 6				District 7			
	%	95% CI		n	%	95% CI		n	%	95% CI		n	%	95% CI		n
TOTAL	34.6	29.9	39.7	363	38.5	33.5	43.9	335	32.6	28.0	37.6	354	27.9	23.6	32.7	372
AGE																
18-19	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
20-24	50.2	38.9	61.5	70	52.3	41.8	62.6	83	43.1	33.6	53.2	91	33.5	25.5	42.7	117
25-29	35.8	28.2	44.2	137	30.1	22.5	38.9	121	30.6	23.6	38.6	134	25.0	18.8	32.3	151
30-34	23.7	16.4	33.1	101	31.6	22.3	42.7	76	15.6	9.6	24.4	85	25.2	16.8	36.0	72
35+	25.0	15.0	38.7	48	31.1	19.4	45.8	42	32.9	19.5	49.8	34	*	*	*	*
ETHNICITY																
Non-Hispanic	34.4	29.5	39.7	330	38.3	32.6	44.3	261	31.4	26.7	36.6	320	26.2	21.8	31.0	352
Hispanic	36.4	21.9	53.8	33	39.4	29.1	50.7	74	42.3	27.4	58.7	34	*	*	*	*
MARITAL STATUS																
Not Married	69.0	56.5	79.2	58	58.5	47.0	69.1	65	60.4	47.8	71.8	53	65.8	50.4	78.4	38
Married	24.9	20.5	29.9	305	29.6	24.8	35.0	270	24.3	20.0	29.0	301	21.0	17.1	25.4	334
INCOME																
Less than \$15,000	57.6	45.1	69.3	59	58.6	46.8	69.4	66	53.2	42.8	63.4	81	41.5	31.5	52.2	84
\$15,000 - \$24,999	44.2	31.7	57.4	54	40.1	29.6	51.6	72	37.4	27.1	48.9	69	32.0	21.7	44.3	59
\$25,000 - \$34,999	46.3	32.8	60.5	45	36.2	24.3	50.0	52	29.8	19.7	42.3	54	29.5	18.8	43.2	52
\$35,000 - \$49,999	29.6	18.1	44.4	47	30.1	20.3	42.2	58	19.6	11.4	31.7	51	15.9	9.0	26.4	63
\$50,000+	17.4	12.2	24.3	146	20.0	13.0	29.6	79	12.2	7.0	20.6	87	15.4	9.7	23.5	101
EDUCATION																
K-11th Grade	*	*	*	*	45.9	33.4	58.9	55	*	*	*	*	*	*	*	*
High School or GED	47.2	35.7	59.0	66	40.8	29.7	53.0	66	43.9	34.6	53.7	96	40.3	28.9	52.9	62
Some College	42.8	34.8	51.1	137	42.5	34.8	50.6	144	33.1	26.0	41.0	138	24.3	18.3	31.5	159
College Graduate+	20.2	14.3	27.8	137	19.3	11.9	29.9	70	16.9	10.7	25.5	96	21.0	15.0	28.5	135

*Figure not reliable by PRATS standards (n < 30)

No Birth Control Use

Idaho mothers not using birth control when they became pregnant (among those not trying to get pregnant)¹, 2010

	Statewide			Public Health District												
				District 1			District 2			District 3						
	%	95% CI		n	%	95% CI		n	%	95% CI		n	%	95% CI		n
TOTAL	54.9	51.5	58.3	911	62.1	54.0	69.6	134	56.5	49.3	63.4	131	52.9	43.7	62.0	109
AGE																
18-19	74.0	62.0	83.3	54	*	*	*	*	*	*	*	*	*	*	*	*
20-24	54.8	48.6	60.8	280	60.5	44.8	74.3	36	58.6	46.1	70.1	43	56.9	41.3	71.3	38
25-29	51.2	45.1	57.2	296	67.1	52.5	79.0	42	54.8	41.9	67.1	40	*	*	*	*
30-34	52.9	45.0	60.6	173	*	*	*	*	*	*	*	*	*	*	*	*
35+	58.1	48.2	67.4	108	*	*	*	*	*	*	*	*	*	*	*	*
ETHNICITY																
Non-Hispanic	56.0	52.3	59.6	790	61.2	53.0	68.9	129	57.1	49.8	64.2	124	56.5	45.5	66.9	78
Hispanic	49.1	40.0	58.2	120	*	*	*	*	*	*	*	*	44.3	28.2	61.7	31
MARITAL STATUS																
Not Married	59.4	53.5	65.0	294	62.8	50.0	74.0	51	60.0	48.0	70.9	45	60.0	44.9	73.4	40
Married	51.5	47.4	55.5	617	61.5	51.5	70.6	83	53.5	44.9	61.9	86	46.4	35.4	57.7	69
INCOME																
Less than \$15,000	59.5	53.4	65.4	281	73.5	58.6	84.4	37	64.4	52.0	75.1	44	62.5	47.4	75.4	42
\$15,000 - \$24,999	47.7	40.3	55.1	192	*	*	*	*	*	*	*	*	*	*	*	*
\$25,000 - \$34,999	50.4	41.8	59.0	141	*	*	*	*	*	*	*	*	*	*	*	*
\$35,000 - \$49,999	52.4	42.5	62.1	112	*	*	*	*	*	*	*	*	*	*	*	*
\$50,000+	57.4	49.1	65.3	156	*	*	*	*	*	*	*	*	*	*	*	*
EDUCATION																
K-11th Grade	53.7	43.2	63.8	94	*	*	*	*	*	*	*	*	*	*	*	*
High School or GED	61.8	55.4	67.9	256	70.2	54.9	82.1	40	62.9	50.9	73.5	47	*	*	*	*
Some College	50.4	45.2	55.5	396	59.9	48.5	70.3	68	45.6	34.5	57.2	50	39.5	26.9	53.7	47
College Graduate+	56.4	48.2	64.2	161	*	*	*	*	*	*	*	*	*	*	*	*

*Figure not reliable by PRATS standards (n < 30)

1. Only asked of mothers who were not trying to get pregnant at the time of conception.

No Birth Control Use

Idaho mothers not using birth control when they became pregnant (among those not trying to get pregnant)¹, 2010

	Public Health District															
	District 4			District 5			District 6			District 7						
	%	95% CI		n	%	95% CI		n	%	95% CI		n	%	95% CI		n
TOTAL	51.4	43.2	59.5	138	54.1	46.1	62.0	142	53.8	45.8	61.7	137	59.0	50.2	67.2	120
AGE																
18-19	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
20-24	48.9	34.2	63.9	40	56.1	41.5	69.7	43	43.3	29.6	58.1	41	62.9	47.2	76.2	39
25-29	43.4	30.8	57.0	51	41.0	27.6	55.9	42	62.2	48.9	73.9	50	59.4	45.0	72.4	44
30-34	*	*	*	*	50.2	33.7	66.7	31	*	*	*	*	*	*	*	*
35+	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
ETHNICITY																
Non-Hispanic	51.4	42.7	60.0	122	56.0	46.7	64.8	108	51.2	42.7	59.7	120	62.5	53.4	70.8	109
Hispanic	*	*	*	*	48.7	32.9	64.8	34	*	*	*	*	*	*	*	*
MARITAL STATUS																
Not Married	55.1	41.6	67.9	49	59.1	45.1	71.8	44	60.0	45.5	73.0	40	*	*	*	*
Married	48.3	38.5	58.3	89	50.0	40.8	59.2	98	49.5	40.4	58.6	97	56.8	47.2	66.0	95
INCOME																
Less than \$15,000	46.9	31.8	62.5	37	59.1	43.8	72.8	39	57.9	44.3	70.4	48	64.7	48.3	78.3	34
\$15,000 - \$24,999	*	*	*	*	52.1	36.7	67.0	38	48.8	33.1	64.7	33	*	*	*	*
\$25,000 - \$34,999	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
\$35,000 - \$49,999	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
\$50,000+	59.7	42.7	74.7	32	*	*	*	*	*	*	*	*	*	*	*	*
EDUCATION																
K-11th Grade	*	*	*	*	62.0	44.3	77.0	30	*	*	*	*	*	*	*	*
High School or GED	49.9	33.9	65.9	34	*	*	*	*	59.9	46.8	71.7	53	*	*	*	*
Some College	46.7	35.0	58.7	64	54.7	43.0	66.0	67	48.4	35.8	61.2	52	59.5	45.5	72.0	48
College Graduate+	66.7	48.7	80.8	30	*	*	*	*	*	*	*	*	61.8	46.1	75.4	37

*Figure not reliable by PRATS standards (n < 30)

1. Only asked of mothers who were not trying to get pregnant at the time of conception.

No Health Insurance

Idaho mothers who did not have private health insurance prior to pregnancy¹, 2010

	Statewide			Public Health District								
				District 1			District 2			District 3		
	%	95% CI	n	%	95% CI	n	%	95% CI	n	%	95% CI	n
TOTAL	37.4	35.4 39.5	2,374	40.3	35.2 45.6	322	34.7	30.5 39.1	344	45.2	39.5 51.2	272
AGE												
18-19	75.8	64.8 84.2	80	*	* *	*	*	* *	*	*	* *	*
20-24	51.9	47.7 56.0	597	61.2	50.5 71.0	75	50.7	41.6 59.9	78	69.7	59.2 78.5	79
25-29	33.0	29.7 36.4	876	35.0	26.7 44.2	105	29.3	23.2 36.1	142	29.3	20.3 40.2	83
30-34	26.2	22.5 30.3	556	24.4	16.5 34.7	85	29.8	21.5 39.6	73	27.1	17.6 39.4	62
35+	26.4	21.1 32.5	265	21.5	11.6 36.4	41	13.9	6.9 26.0	38	48.1	33.1 63.4	38
ETHNICITY												
Non-Hispanic	32.6	30.5 34.7	2,122	40.1	34.9 45.5	308	33.9	29.7 38.4	333	34.8	28.6 41.5	209
Hispanic	70.4	64.4 75.7	249	*	* *	*	*	* *	*	77.3	66.1 85.6	63
MARITAL STATUS												
Not Married	70.6	65.9 74.9	411	69.0	58.4 78.0	71	65.7	55.9 74.3	67	74.1	62.0 83.5	58
Married	26.5	24.5 28.5	1,963	26.3	21.6 31.6	251	22.7	19.0 27.0	277	32.7	27.0 39.0	214
INCOME												
Less than \$15,000	76.1	72.1 79.7	497	79.0	68.0 87.0	60	69.9	60.8 77.7	76	86.5	76.8 92.6	71
\$15,000 - \$24,999	56.7	51.6 61.7	391	67.5	54.7 78.2	52	60.7	48.3 71.8	43	56.3	41.1 70.5	39
\$25,000 - \$34,999	37.9	32.7 43.3	358	37.4	26.2 50.1	56	30.8	21.5 42.0	53	45.1	31.0 60.1	41
\$35,000 - \$49,999	14.4	10.9 18.8	352	10.5	4.2 23.7	43	14.8	8.6 24.5	59	6.5	1.7 21.8	30
\$50,000+	4.0	2.8 5.8	708	8.7	4.6 15.8	105	3.7	1.7 8.0	106	8.9	4.4 17.2	82
EDUCATION												
K-11th Grade	85.3	79.4 89.7	185	*	* *	*	*	* *	*	86.0	70.7 93.9	37
High School or GED	53.5	49.0 57.9	524	68.1	57.7 77.0	74	57.0	48.4 65.1	90	53.6	41.8 65.1	66
Some College	35.3	32.1 38.6	940	41.4	33.6 49.6	135	33.9	27.1 41.4	119	38.4	29.5 48.1	104
College Graduate+	10.8	8.6 13.6	719	4.3	1.7 10.1	92	12.9	8.7 18.8	123	12.7	6.6 22.9	62

*Figure not reliable by PRATS standards (n < 30)

1. Does not include Medicaid.

No Health Insurance

Idaho mothers who did not have private health insurance prior to pregnancy¹, 2010

	Public Health District															
	District 4				District 5				District 6				District 7			
	%	95% CI		n	%	95% CI		n	%	95% CI		n	%	95% CI		n
TOTAL	29.8	25.3	34.7	367	47.9	42.7	53.1	340	39.3	34.5	44.3	356	31.2	26.7	36.1	373
AGE																
18-19	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
20-24	47.5	36.3	58.9	71	56.3	45.9	66.2	84	49.7	40.0	59.3	92	34.7	26.7	43.7	118
25-29	30.1	23.0	38.4	137	47.7	39.2	56.3	123	37.3	29.8	45.4	135	27.1	20.6	34.8	151
30-34	21.6	14.6	30.8	102	41.7	31.4	52.9	76	24.9	17.1	34.8	86	22.5	14.6	33.1	72
35+	9.6	3.7	22.4	50	24.4	13.6	39.8	44	43.9	28.9	60.1	33	*	*	*	*
ETHNICITY																
Non-Hispanic	28.1	23.5	33.3	333	37.4	31.7	43.4	263	35.4	30.5	40.6	323	27.8	23.4	32.7	353
Hispanic	45.4	29.9	61.8	34	79.3	69.6	86.5	77	72.7	56.1	84.8	33	*	*	*	*
MARITAL STATUS																
Not Married	62.7	50.2	73.7	59	75.8	64.8	84.1	66	71.2	58.6	81.2	52	76.3	61.3	86.8	38
Married	20.5	16.4	25.2	308	35.4	30.3	40.9	274	29.9	25.4	34.9	304	23.0	19.0	27.6	335
INCOME																
Less than \$15,000	73.9	61.7	83.3	59	80.6	69.6	88.3	66	81.5	72.3	88.2	80	59.9	49.7	69.3	85
\$15,000 - \$24,999	56.7	43.8	68.8	55	65.8	54.9	75.3	74	51.6	40.4	62.6	69	44.3	32.6	56.8	59
\$25,000 - \$34,999	33.0	21.2	47.3	46	42.7	30.7	55.7	55	37.5	26.3	50.2	55	35.6	24.1	49.0	52
\$35,000 - \$49,999	14.8	7.0	28.7	47	24.9	15.7	37.1	58	11.5	5.6	22.3	52	14.3	7.9	24.6	63
\$50,000+	2.0	0.7	5.8	148	6.3	2.8	13.5	79	1.1	0.2	6.7	87	1.0	0.2	6.1	101
EDUCATION																
K-11th Grade	*	*	*	*	90.8	82.3	95.4	56	*	*	*	*	*	*	*	*
High School or GED	43.7	32.6	55.5	68	64.2	52.5	74.3	68	48.2	38.7	57.8	96	51.9	39.9	63.6	62
Some College	34.6	27.2	42.9	138	35.1	27.8	43.1	145	38.8	31.4	46.8	139	28.6	22.3	35.9	160
College Graduate+	8.4	4.7	14.8	138	12.2	6.4	22.0	71	12.8	7.7	20.6	98	14.0	9.2	20.8	135

*Figure not reliable by PRATS standards (n < 30)

1. Does not include Medicaid.

Medicaid Utilization

Idaho mothers who reported Medicaid as a payment source for prenatal care and/or delivery, 2010

	Statewide			Public Health District								
				District 1			District 2			District 3		
	%	95% CI	n	%	95% CI	n	%	95% CI	n	%	95% CI	n
TOTAL	43.9	41.8 46.0	2,363	43.9	38.7 49.2	319	43.1	38.8 47.6	343	49.1	43.3 55.0	271
AGE												
18-19	90.1	81.0 95.1	80	*	* *	*	*	* *	*	*	* *	*
20-24	63.8	59.8 67.7	596	72.7	62.7 80.8	74	58.2	49.1 66.9	78	73.1	62.9 81.3	79
25-29	39.2	35.8 42.7	869	34.5	26.2 43.9	103	41.9	35.2 48.9	141	35.0	25.4 46.1	83
30-34	28.5	24.6 32.7	555	20.7	13.4 30.4	85	23.9	16.3 33.5	73	29.6	19.5 42.1	62
35+	26.2	20.9 32.2	263	22.2	12.4 36.3	41	25.2	15.3 38.6	38	43.5	28.8 59.4	37
ETHNICITY												
Non-Hispanic	39.8	37.6 42.0	2,114	43.5	38.2 48.9	305	41.9	37.5 46.5	332	39.5	33.1 46.4	208
Hispanic	72.3	66.5 77.5	246	*	* *	*	*	* *	*	78.7	68.0 86.5	63
MARITAL STATUS												
Not Married	80.9	76.6 84.5	409	77.1	66.9 85.0	70	80.6	71.7 87.2	67	87.7	76.9 93.9	57
Married	31.7	29.7 33.8	1,954	27.7	22.9 33.1	249	28.6	24.5 33.2	276	32.7	27.0 39.0	214
INCOME												
Less than \$15,000	91.0	88.0 93.3	498	92.3	83.4 96.7	60	92.1	86.0 95.7	77	91.0	82.4 95.6	71
\$15,000 - \$24,999	71.3	66.4 75.7	387	69.4	56.6 79.8	51	68.4	56.3 78.5	43	66.3	51.0 78.9	38
\$25,000 - \$34,999	41.3	36.1 46.8	358	44.4	32.5 57.0	56	40.5	30.1 51.8	53	46.5	32.1 61.4	41
\$35,000 - \$49,999	14.1	10.6 18.5	349	13.0	5.8 26.5	42	19.0	11.5 29.8	57	8.5	2.2 27.6	30
\$50,000+	2.3	1.3 4.0	704	2.5	0.7 8.9	104	2.4	0.8 7.3	106	9.1	4.2 18.5	82
EDUCATION												
K-11th Grade	88.8	83.7 92.4	183	*	* *	*	*	* *	*	90.4	77.1 96.3	37
High School or GED	60.5	56.0 64.7	525	67.0	56.6 76.0	74	68.5	60.5 75.6	91	65.9	54.3 75.9	66
Some College	45.0	41.7 48.4	934	47.4	39.4 55.5	134	43.5	36.1 51.1	117	40.6	31.6 50.4	103
College Graduate+	13.6	11.2 16.5	715	6.5	3.1 13.0	90	15.3	10.7 21.4	123	9.5	4.5 19.1	62

*Figure not reliable by PRATS standards (n < 30)

Medicaid Utilization

Idaho mothers who reported Medicaid as a payment source for prenatal care and/or delivery, 2010

	Public Health District															
	District 4				District 5				District 6				District 7			
	%	95% CI		n	%	95% CI		n	%	95% CI		n	%	95% CI		n
TOTAL	32.1	27.5	37.2	363	54.5	49.4	59.6	339	48.4	43.5	53.4	356	45.1	40.2	50.0	372
AGE																
18-19	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
20-24	48.7	37.4	60.0	71	64.6	54.5	73.5	84	67.5	58.2	75.6	92	63.8	55.2	71.6	118
25-29	32.1	24.8	40.5	134	50.8	42.3	59.4	122	45.7	37.9	53.8	135	40.8	33.4	48.7	151
30-34	24.7	17.2	34.1	101	52.5	41.7	63.1	76	24.1	16.5	33.8	86	24.7	16.2	35.6	72
35+	10.6	4.6	22.6	50	30.7	18.9	45.8	44	45.9	30.7	62.0	33	*	*	*	*
ETHNICITY																
Non-Hispanic	30.9	26.1	36.1	332	45.2	39.4	51.2	262	46.5	41.3	51.7	323	42.1	37.2	47.2	352
Hispanic	45.0	28.8	62.2	31	82.2	72.9	88.7	77	65.4	48.9	78.9	33	*	*	*	*
MARITAL STATUS																
Not Married	66.1	53.7	76.7	59	89.4	80.2	94.6	66	84.6	73.2	91.7	52	86.8	73.0	94.2	38
Married	22.4	18.2	27.2	304	38.8	33.6	44.4	273	37.8	32.9	43.0	304	37.4	32.6	42.5	334
INCOME																
Less than \$15,000	89.5	78.7	95.1	59	93.9	85.1	97.6	66	94.1	88.1	97.2	80	87.5	79.1	92.8	85
\$15,000 - \$24,999	56.1	42.9	68.5	53	84.8	76.3	90.6	74	75.7	64.6	84.2	69	79.4	68.2	87.4	59
\$25,000 - \$34,999	35.0	23.0	49.4	46	41.1	29.2	54.1	55	44.5	32.7	57.0	55	40.3	28.2	53.7	52
\$35,000 - \$49,999	13.8	6.5	27.2	47	26.3	16.7	39.0	58	5.8	2.0	15.3	52	12.7	6.7	22.8	63
\$50,000+	0.0	0.0	0.0	147	1.3	0.2	7.6	78	0.0	0.0	0.0	87	2.0	0.5	7.1	100
EDUCATION																
K-11th Grade	*	*	*	*	89.6	80.0	94.9	56	*	*	*	*	*	*	*	*
High School or GED	48.3	36.9	59.9	68	68.1	57.1	77.4	68	63.7	54.4	72.2	96	53.3	41.3	64.9	62
Some College	36.8	29.2	45.1	137	50.4	42.5	58.2	144	48.2	40.4	56.0	139	52.2	44.8	59.6	160
College Graduate+	8.5	4.7	14.9	137	10.8	5.4	20.5	71	18.5	12.2	27.2	98	24.3	18.0	32.0	134

*Figure not reliable by PRATS standards (n < 30)

Overweight/Obesity

Idaho mothers who were overweight or obese prior to pregnancy¹, 2010

	Statewide			Public Health District												
				District 1			District 2			District 3						
	%	95% CI		n	%	95% CI		n	%	95% CI		n	%	95% CI		n
TOTAL	44.4	42.3	46.6	2,313	47.1	41.9	52.3	320	45.4	41.1	49.8	341	52.3	46.3	58.3	256
AGE																
18-19	29.6	20.3	40.9	77	*	*	*	*	*	*	*	*	*	*	*	*
20-24	44.2	40.0	48.5	585	47.5	37.0	58.3	75	46.5	37.5	55.7	79	57.3	46.2	67.8	76
25-29	40.9	37.5	44.4	858	44.8	36.1	53.8	105	40.0	33.5	46.9	140	44.0	33.7	54.8	79
30-34	49.1	44.7	53.5	537	48.4	38.5	58.4	84	54.6	45.0	63.9	71	58.4	45.7	70.1	57
35+	52.4	46.0	58.8	256	46.6	32.9	60.8	41	58.9	45.7	70.9	38	63.7	46.9	77.8	34
ETHNICITY																
Non-Hispanic	43.0	40.8	45.2	2,100	46.1	40.8	51.4	306	46.7	42.3	51.2	330	49.9	43.2	56.6	204
Hispanic	56.5	49.6	63.2	210	*	*	*	*	*	*	*	*	61.1	47.5	73.2	52
MARITAL STATUS																
Not Married	51.1	46.0	56.2	388	54.3	43.5	64.7	70	42.4	33.1	52.3	66	51.9	39.2	64.3	54
Married	42.3	40.1	44.6	1,925	43.6	38.0	49.3	250	46.6	41.8	51.4	275	52.5	45.8	59.0	202
INCOME																
Less than \$15,000	47.5	42.8	52.2	478	53.3	41.2	64.9	59	49.6	40.4	58.8	77	48.2	36.3	60.2	63
\$15,000 - \$24,999	49.1	43.8	54.3	375	49.0	36.4	61.8	52	46.7	34.7	59.1	43	58.4	42.0	73.2	35
\$25,000 - \$34,999	49.9	44.4	55.4	350	49.3	37.2	61.5	56	46.4	35.6	57.5	53	64.2	49.0	77.0	41
\$35,000 - \$49,999	44.0	38.6	49.5	350	37.9	25.4	52.2	42	38.4	28.7	49.1	59	57.5	40.3	73.1	30
\$50,000+	38.8	35.1	42.7	699	43.6	34.9	52.6	105	44.2	36.6	52.2	103	48.2	37.7	58.8	79
EDUCATION																
K-11th Grade	51.4	43.3	59.4	155	*	*	*	*	*	*	*	*	*	*	*	*
High School or GED	48.9	44.4	53.5	513	49.6	38.8	60.4	73	42.9	34.7	51.6	89	55.9	43.7	67.4	64
Some College	47.6	44.2	51.0	927	51.8	43.7	59.7	135	55.5	48.0	62.7	119	56.7	47.0	65.8	101
College Graduate+	34.2	30.7	38.0	712	34.8	26.3	44.5	91	35.3	28.7	42.6	121	40.2	29.0	52.6	60

*Figure not reliable by PRATS standards (n < 30)

1. Mothers with a Body Mass Index (BMI) of 25.0 or greater.

Overweight/Obesity

Idaho mothers who were overweight or obese prior to pregnancy¹, 2010

	Public Health District															
	District 4				District 5				District 6				District 7			
	%	95% CI		n	%	95% CI		n	%	95% CI		n	%	95% CI		n
TOTAL	40.9	36.0	45.9	362	50.6	45.3	55.9	324	43.0	38.1	48.0	346	37.4	32.7	42.3	364
AGE																
18-19	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
20-24	40.1	29.3	52.0	67	43.2	33.1	53.9	82	41.9	32.6	51.9	90	36.0	27.9	45.0	116
25-29	39.6	31.8	47.9	137	49.0	40.4	57.8	118	39.0	31.4	47.2	131	35.3	28.3	43.1	148
30-34	43.5	34.4	53.1	101	58.9	47.5	69.4	71	47.1	37.3	57.2	83	43.5	32.8	54.8	70
35+	42.2	29.6	56.0	50	59.3	44.4	72.8	41	59.8	43.6	74.2	32	*	*	*	*
ETHNICITY																
Non-Hispanic	41.1	36.0	46.4	329	44.2	38.4	50.1	262	42.7	37.7	47.9	320	37.2	32.5	42.3	349
Hispanic	39.0	24.0	56.5	33	74.5	62.8	83.5	62	*	*	*	*	*	*	*	*
MARITAL STATUS																
Not Married	51.8	39.3	64.1	56	52.5	40.8	63.9	61	44.9	32.6	57.9	49	53.1	36.9	68.7	32
Married	37.9	32.8	43.3	306	49.8	44.2	55.5	263	42.4	37.3	47.7	297	34.9	30.2	40.0	332
INCOME																
Less than \$15,000	42.6	30.5	55.5	56	53.8	41.7	65.4	63	43.4	33.4	53.9	80	46.9	36.5	57.6	80
\$15,000 - \$24,999	44.2	31.8	57.4	54	54.4	42.7	65.6	67	47.8	36.6	59.2	66	45.3	33.5	57.7	58
\$25,000 - \$34,999	52.7	38.6	66.3	45	53.0	39.9	65.7	53	44.7	32.5	57.7	50	34.9	23.4	48.5	52
\$35,000 - \$49,999	50.4	36.8	64.1	47	54.0	41.8	65.8	57	42.3	30.6	55.0	52	27.0	17.9	38.6	63
\$50,000+	34.1	27.0	41.9	148	41.9	32.0	52.4	78	38.4	29.4	48.3	87	33.7	25.5	43.2	99
EDUCATION																
K-11th Grade	*	*	*	*	57.9	43.6	71.0	46	*	*	*	*	*	*	*	*
High School or GED	45.9	34.5	57.8	66	52.5	40.7	64.1	66	47.6	38.2	57.1	96	44.6	32.8	57.1	59
Some College	44.6	36.7	52.9	137	49.9	41.9	57.8	141	40.2	32.7	48.2	135	42.6	35.4	50.2	159
College Graduate+	33.9	26.7	42.0	138	43.8	33.3	54.9	71	35.9	27.6	45.2	98	26.4	19.9	34.2	133

*Figure not reliable by PRATS standards (n < 30)

1. Mothers with a Body Mass Index (BMI) of 25.0 or greater.

First Trimester Prenatal Care

Idaho mothers who began prenatal care during the first trimester, 2010

	Statewide			Public Health District								
				District 1			District 2			District 3		
	%	95% CI	n	%	95% CI	n	%	95% CI	n	%	95% CI	n
TOTAL	90.9	89.5 92.0	2,352	89.6	85.7 92.5	319	87.9	84.6 90.5	341	90.2	85.8 93.3	271
AGE												
18-19	79.1	68.2 87.0	79	*	* *	*	*	* *	*	*	* *	*
20-24	90.0	87.1 92.3	588	86.4	77.0 92.3	73	86.5	79.1 91.6	78	83.9	73.5 90.7	78
25-29	92.1	90.0 93.8	868	92.0	84.7 96.0	104	89.7	84.4 93.4	141	96.0	88.5 98.7	83
30-34	91.9	89.3 93.9	553	93.7	86.4 97.3	85	89.5	82.4 93.9	72	92.4	83.4 96.7	62
35+	90.9	86.4 94.0	264	91.5	77.8 97.1	41	87.6	76.7 93.8	38	91.4	76.7 97.1	38
ETHNICITY												
Non-Hispanic	91.7	90.3 92.8	2,104	89.4	85.4 92.4	305	88.4	85.1 91.0	330	92.0	87.4 95.0	208
Hispanic	85.3	80.0 89.3	245	*	* *	*	*	* *	*	84.5	72.6 91.9	63
MARITAL STATUS												
Not Married	86.2	82.5 89.3	401	81.4	71.6 88.4	70	84.4	75.7 90.3	64	82.5	70.9 90.1	57
Married	92.4	91.1 93.5	1,951	93.6	90.1 95.9	249	89.2	85.8 91.8	277	93.5	89.5 96.0	214
INCOME												
Less than \$15,000	84.7	81.0 87.8	485	83.8	73.0 90.8	60	78.5	69.5 85.4	74	82.6	72.0 89.7	72
\$15,000 - \$24,999	91.0	87.6 93.5	389	86.9	74.7 93.8	51	93.0	83.3 97.3	43	92.3	75.3 97.9	38
\$25,000 - \$34,999	86.9	82.7 90.1	357	82.0	69.8 90.0	55	86.9	77.4 92.7	53	89.8	76.1 96.1	41
\$35,000 - \$49,999	93.9	90.8 96.0	351	97.8	87.7 99.6	43	93.4	86.2 97.0	58	93.5	78.2 98.3	30
\$50,000+	96.6	94.9 97.8	705	99.1	94.6 99.9	104	89.9	84.2 93.6	106	96.5	90.1 98.8	82
EDUCATION												
K-11th Grade	83.9	77.4 88.8	180	*	* *	*	*	* *	*	83.8	67.6 92.8	38
High School or GED	89.4	86.5 91.8	520	83.8	73.7 90.5	73	83.9	76.4 89.4	90	89.1	79.0 94.7	65
Some College	90.4	88.2 92.2	930	89.2	82.7 93.4	134	88.2	82.3 92.4	117	91.9	84.7 95.8	103
College Graduate+	95.1	93.2 96.5	716	97.9	92.6 99.4	91	90.4	85.3 93.9	123	95.3	86.8 98.4	62

*Figure not reliable by PRATS standards (n < 30)

First Trimester Prenatal Care

Idaho mothers who began prenatal care during the first trimester, 2010

	Public Health District															
	District 4				District 5				District 6				District 7			
	%	95% CI		n	%	95% CI		n	%	95% CI		n	%	95% CI		n
TOTAL	92.8	89.7	95.0	361	89.5	85.8	92.4	335	90.8	87.4	93.4	354	91.2	87.9	93.7	371
AGE																
18-19	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
20-24	90.4	81.4	95.3	69	91.4	83.2	95.8	82	93.0	85.7	96.7	91	94.8	89.1	97.6	117
25-29	94.2	89.0	97.0	135	86.5	79.2	91.4	120	90.5	84.1	94.5	134	91.9	86.2	95.4	151
30-34	94.2	88.0	97.3	100	89.9	80.6	95.0	76	90.2	82.2	94.8	86	89.2	80.3	94.3	72
35+	91.4	79.5	96.7	50	93.7	83.1	97.8	44	85.8	71.3	93.6	33	*	*	*	*
ETHNICITY																
Non-Hispanic	92.9	89.6	95.2	328	91.1	86.9	94.0	260	91.2	87.7	93.7	321	92.5	89.3	94.8	352
Hispanic	92.0	78.4	97.3	33	84.9	75.4	91.2	75	87.4	69.7	95.5	33	*	*	*	*
MARITAL STATUS																
Not Married	92.9	82.9	97.2	56	87.7	78.1	93.5	65	86.5	75.4	93.1	52	83.8	69.2	92.2	37
Married	92.8	89.4	95.1	305	90.4	86.5	93.2	270	92.1	88.7	94.5	302	92.5	89.4	94.8	334
INCOME																
Less than \$15,000	84.0	72.1	91.4	55	88.2	77.7	94.2	62	83.5	74.0	89.9	78	88.3	79.9	93.5	84
\$15,000 - \$24,999	95.2	86.5	98.4	55	85.3	75.5	91.6	74	92.3	84.5	96.3	69	91.1	80.8	96.2	59
\$25,000 - \$34,999	85.5	73.1	92.8	46	87.1	76.8	93.2	55	91.8	80.8	96.7	55	85.2	72.6	92.6	52
\$35,000 - \$49,999	98.0	87.9	99.7	47	88.5	77.2	94.6	58	96.2	87.1	98.9	52	90.5	81.0	95.5	63
\$50,000+	96.3	91.7	98.4	147	97.5	91.1	99.3	78	95.4	89.2	98.2	87	98.0	92.9	99.5	101
EDUCATION																
K-11th Grade	*	*	*	*	84.1	72.4	91.5	54	*	*	*	*	*	*	*	*
High School or GED	96.0	88.6	98.7	67	85.4	75.1	91.9	67	93.7	87.0	97.0	96	85.6	75.0	92.2	62
Some College	86.9	80.3	91.5	136	93.8	88.8	96.7	143	85.3	78.6	90.2	137	95.2	91.0	97.5	160
College Graduate+	97.2	92.9	98.9	137	90.6	81.0	95.6	71	98.0	93.0	99.4	98	92.5	86.5	96.0	134

*Figure not reliable by PRATS standards (n < 30)

Idaho mothers who participated in the WIC Program during pregnancy¹, 2010

	Statewide			Public Health District												
				District 1			District 2			District 3						
	%	95% CI		n	%	95% CI		n	%	95% CI		n	%	95% CI		n
TOTAL	39.9	37.8	42.0	2,366	42.9	37.8	48.2	321	42.2	37.8	46.6	345	42.9	37.2	48.9	269
AGE																
18-19	77.4	65.8	85.9	79	*	*	*	*	*	*	*	*	*	*	*	*
20-24	58.0	53.8	62.1	594	64.8	54.2	74.1	75	58.7	49.6	67.2	79	57.7	46.7	68.0	77
25-29	36.0	32.7	39.5	875	35.5	27.3	44.6	105	37.0	30.6	44.0	142	35.3	25.7	46.3	84
30-34	24.8	21.2	28.8	553	23.3	15.6	33.2	84	28.3	20.3	37.8	73	22.6	14.0	34.3	61
35+	26.6	21.3	32.6	265	24.4	14.2	38.7	41	27.7	17.4	41.1	38	41.9	27.3	58.0	37
ETHNICITY																
Non-Hispanic	35.5	33.3	37.7	2,116	43.1	37.8	48.5	307	41.4	37.0	45.9	334	30.8	24.8	37.6	207
Hispanic	69.9	63.9	75.4	247	*	*	*	*	*	*	*	*	80.2	68.9	88.1	62
MARITAL STATUS																
Not Married	67.9	63.0	72.4	409	70.0	59.3	78.9	70	76.1	66.9	83.4	67	73.7	61.4	83.2	57
Married	30.6	28.6	32.8	1,957	29.9	24.9	35.3	251	29.1	25.0	33.7	278	29.7	24.2	36.0	212
INCOME																
Less than \$15,000	74.7	70.4	78.5	497	82.9	71.8	90.2	60	81.6	73.7	87.5	77	77.4	66.3	85.6	69
\$15,000 - \$24,999	61.6	56.5	66.5	390	68.7	56.0	79.0	52	65.3	53.0	75.9	43	61.0	45.6	74.5	39
\$25,000 - \$34,999	48.5	43.1	53.9	357	43.7	31.9	56.3	55	41.2	30.8	52.4	53	47.8	33.3	62.6	41
\$35,000 - \$49,999	19.3	15.4	24.0	351	23.7	13.8	37.7	43	26.4	17.9	37.2	59	18.3	8.0	36.5	30
\$50,000+	2.5	1.5	4.0	706	2.5	0.7	8.9	105	4.2	1.9	9.2	106	5.4	2.1	13.3	82
EDUCATION																
K-11th Grade	79.2	72.6	84.6	184	*	*	*	*	*	*	*	*	87.6	71.9	95.1	36
High School or GED	57.4	52.9	61.8	523	62.1	51.4	71.8	74	67.6	59.5	74.8	91	59.6	47.6	70.6	65
Some College	38.0	34.7	41.3	935	46.0	38.1	54.2	134	40.0	32.9	47.6	119	32.0	23.6	41.8	103
College Graduate+	14.9	12.4	17.8	718	10.6	6.1	17.9	92	16.9	12.1	23.2	123	5.7	1.9	16.0	62

*Figure not reliable by PRATS standards (n < 30)

1. The WIC Program is the supplemental nutrition program for women, infants, and children.

Idaho mothers who participated in the WIC Program during pregnancy¹, 2010

	Public Health District															
	District 4				District 5				District 6				District 7			
	%	95% CI		n	%	95% CI		n	%	95% CI		n	%	95% CI		n
TOTAL	29.4	25.0	34.3	364	46.8	41.6	52.1	338	45.0	40.1	50.0	357	41.0	36.2	46.0	372
AGE																
18-19	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
20-24	44.2	33.1	55.9	69	66.7	56.6	75.4	84	59.9	50.3	68.9	92	61.0	52.3	69.0	118
25-29	31.2	24.0	39.3	137	39.0	30.8	47.9	122	45.7	37.9	53.8	135	35.1	28.1	42.9	150
30-34	24.2	16.9	33.5	101	34.8	24.9	46.1	76	25.2	17.5	34.9	86	19.3	11.8	29.9	72
35+	10.6	4.6	22.6	50	38.9	25.8	53.8	44	33.7	20.4	50.2	34	*	*	*	*
ETHNICITY																
Non-Hispanic	27.8	23.3	32.9	330	38.9	33.2	44.9	263	41.2	36.1	46.4	323	37.8	33.0	42.8	352
Hispanic	44.1	28.7	60.7	34	71.3	60.0	80.4	75	77.2	62.1	87.5	34	*	*	*	*
MARITAL STATUS																
Not Married	50.9	38.7	62.9	59	72.3	61.1	81.3	65	73.6	61.3	83.1	53	73.7	58.5	84.8	38
Married	23.3	19.0	28.2	305	35.5	30.4	41.0	273	36.5	31.7	41.7	304	35.0	30.3	40.0	334
INCOME																
Less than \$15,000	64.8	52.1	75.7	59	73.1	61.9	82.0	66	76.6	66.8	84.2	81	75.6	66.0	83.3	85
\$15,000 - \$24,999	59.9	46.9	71.7	55	62.8	51.4	72.9	73	64.1	52.8	74.0	69	56.0	43.7	67.7	59
\$25,000 - \$34,999	40.2	27.5	54.4	46	63.1	50.4	74.2	55	52.0	39.7	64.1	55	50.0	37.2	62.9	52
\$35,000 - \$49,999	15.2	7.5	28.2	46	23.4	14.8	35.1	58	17.3	9.7	29.0	52	19.1	11.5	30.0	63
\$50,000+	0.0	0.0	0.0	147	1.3	0.2	7.5	79	4.2	1.4	11.8	87	4.0	1.6	9.7	100
EDUCATION																
K-11th Grade	*	*	*	*	71.6	58.4	81.9	55	*	*	*	*	*	*	*	*
High School or GED	43.0	31.9	54.9	67	64.8	53.5	74.7	68	60.8	51.3	69.6	96	58.1	46.0	69.2	62
Some College	34.2	26.8	42.4	137	41.9	34.2	50.0	144	40.4	32.9	48.3	139	38.4	31.4	45.9	159
College Graduate+	9.2	5.4	15.4	137	11.0	5.8	19.8	71	22.8	16.0	31.6	98	27.1	20.5	34.9	135

*Figure not reliable by PRATS standards (n < 30)

1. The WIC Program is the supplemental nutrition program for women, infants, and children.

Multivitamin Use

Idaho mothers who regularly took a multivitamin during the third trimester¹, 2010

	Statewide			Public Health District								
				District 1			District 2			District 3		
	%	95% CI	n	%	95% CI	n	%	95% CI	n	%	95% CI	n
TOTAL	83.1	81.5 84.7	2,367	83.8	79.6 87.3	321	85.2	81.7 88.1	345	80.8	75.7 85.0	269
AGE												
18-19	77.7	66.9 85.7	80	*	* *	*	*	* *	*	*	* *	*
20-24	81.0	77.5 84.1	597	83.1	74.3 89.4	75	83.0	74.5 89.1	79	76.4	66.0 84.4	79
25-29	83.4	80.7 85.8	874	86.3	78.3 91.6	105	82.7	76.7 87.4	142	85.3	76.1 91.3	83
30-34	85.8	82.6 88.6	552	85.8	77.3 91.5	84	87.7	80.0 92.7	73	82.7	71.0 90.3	61
35+	83.9	78.8 87.9	264	81.8	68.7 90.2	41	95.1	85.7 98.4	38	83.3	67.9 92.2	36
ETHNICITY												
Non-Hispanic	83.3	81.6 84.9	2,116	83.6	79.3 87.2	307	85.9	82.5 88.8	334	81.3	75.5 86.0	207
Hispanic	81.9	76.6 86.2	248	*	* *	*	*	* *	*	79.2	67.5 87.4	62
MARITAL STATUS												
Not Married	83.3	79.2 86.7	410	85.7	76.4 91.8	70	77.6	68.5 84.7	67	79.0	67.0 87.4	57
Married	83.1	81.3 84.7	1,957	82.9	78.1 86.8	251	88.1	84.7 90.9	278	81.6	76.1 86.1	212
INCOME												
Less than \$15,000	79.1	75.1 82.6	499	79.0	68.0 87.0	60	76.3	67.2 83.4	77	78.6	67.9 86.4	71
\$15,000 - \$24,999	85.1	81.2 88.3	390	74.9	62.4 84.2	52	89.9	79.0 95.5	43	82.2	67.4 91.1	39
\$25,000 - \$34,999	77.6	72.6 81.8	356	82.6	71.3 90.0	55	85.1	75.4 91.4	53	72.4	57.0 83.8	40
\$35,000 - \$49,999	83.7	79.4 87.2	352	80.7	67.0 89.6	43	86.1	77.0 92.0	59	86.9	70.7 94.8	30
\$50,000+	88.0	85.2 90.3	704	94.5	89.0 97.4	105	89.3	83.3 93.3	106	85.6	76.1 91.7	81
EDUCATION												
K-11th Grade	77.7	71.2 83.1	185	*	* *	*	*	* *	*	79.9	63.8 90.0	36
High School or GED	80.9	77.2 84.2	524	73.6	63.4 81.7	74	83.8	76.4 89.1	91	77.3	65.9 85.8	66
Some College	83.1	80.4 85.5	935	86.3	79.9 90.9	134	85.8	79.5 90.3	119	82.5	74.1 88.6	103
College Graduate+	86.9	84.1 89.3	717	93.6	87.3 96.9	92	88.7	83.0 92.7	123	83.9	73.1 90.9	61

*Figure not reliable by PRATS standards (n < 30)

1. Multivitamin use four or more times per week during the last three months of pregnancy.

Multivitamin Use

Idaho mothers who regularly took a multivitamin during the third trimester¹, 2010

	Public Health District															
	District 4				District 5				District 6				District 7			
	%	95% CI		n	%	95% CI		n	%	95% CI		n	%	95% CI		n
TOTAL	87.1	83.3	90.1	365	79.9	75.5	83.8	338	84.2	80.3	87.4	357	80.5	76.3	84.0	372
AGE																
18-19	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
20-24	83.8	73.2	90.7	70	72.2	61.7	80.7	84	85.6	77.5	91.1	92	84.0	76.3	89.5	118
25-29	86.2	79.6	91.0	137	79.4	71.8	85.3	122	84.6	78.1	89.4	135	79.2	72.2	84.8	150
30-34	90.5	83.5	94.7	101	87.4	77.9	93.2	75	85.4	77.1	91.0	86	78.3	68.0	86.0	72
35+	90.4	79.3	95.8	50	83.1	70.5	91.0	44	73.6	57.6	85.2	34	*	*	*	*
ETHNICITY																
Non-Hispanic	86.9	82.9	90.1	331	79.2	74.0	83.6	262	84.9	80.9	88.2	323	80.4	76.0	84.1	352
Hispanic	88.3	73.0	95.5	34	82.3	72.4	89.1	76	77.9	62.2	88.3	34	*	*	*	*
MARITAL STATUS																
Not Married	86.4	75.6	92.9	59	80.3	69.8	87.8	66	90.6	80.3	95.8	53	81.6	67.0	90.6	38
Married	87.3	83.2	90.5	306	79.8	74.9	83.9	272	82.2	77.9	85.9	304	80.2	75.9	84.0	334
INCOME																
Less than \$15,000	81.0	69.1	89.1	59	74.5	62.9	83.4	66	82.8	73.8	89.2	81	79.1	69.5	86.3	85
\$15,000 - \$24,999	90.4	80.5	95.5	55	88.4	78.7	94.0	73	81.0	71.0	88.1	69	84.8	73.8	91.7	59
\$25,000 - \$34,999	78.4	64.6	87.8	46	71.1	58.7	81.0	55	91.3	81.6	96.1	55	72.0	59.0	82.1	52
\$35,000 - \$49,999	91.1	78.9	96.6	47	79.9	68.6	87.9	58	73.1	60.6	82.7	52	82.5	71.8	89.8	63
\$50,000+	90.0	84.3	93.8	147	84.8	75.8	90.9	78	88.6	80.8	93.5	87	83.1	75.0	89.0	100
EDUCATION																
K-11th Grade	*	*	*	*	69.4	56.7	79.8	56	*	*	*	*	*	*	*	*
High School or GED	87.4	77.1	93.5	67	79.9	69.1	87.7	68	82.0	73.8	88.0	96	79.7	68.0	87.9	62
Some College	82.7	75.8	88.0	137	82.2	75.3	87.4	144	84.5	78.0	89.3	139	81.5	75.1	86.6	159
College Graduate+	90.1	83.9	94.1	138	86.1	76.7	92.1	70	86.9	79.5	92.0	98	81.1	74.0	86.6	135

*Figure not reliable by PRATS standards (n < 30)

1. Multivitamin use four or more times per week during the last three months of pregnancy.

HIV Testing

Idaho mothers who received an HIV test during pregnancy, 2010

	Statewide			Public Health District								
				District 1			District 2			District 3		
	%	95% CI	n	%	95% CI	n	%	95% CI	n	%	95% CI	n
TOTAL	45.5	43.4 47.6	2,332	51.2	45.9 56.4	316	44.4	40.0 48.8	340	45.9	40.1 51.8	270
AGE												
18-19	62.1	50.9 72.2	80	*	* *	*	*	* *	*	*	* *	*
20-24	50.8	46.6 55.0	588	65.9	55.1 75.3	75	45.8	36.7 55.1	77	50.1	39.3 60.9	78
25-29	42.8	39.3 46.3	864	42.8	34.1 51.9	105	45.0	38.2 52.0	140	40.8	31.0 51.4	84
30-34	41.5	37.2 46.0	538	43.9	34.0 54.3	80	45.5	36.3 55.0	73	39.0	27.8 51.5	60
35+	43.7	37.5 50.1	262	45.3	31.6 59.9	40	36.0	24.4 49.6	37	42.8	28.4 58.5	38
ETHNICITY												
Non-Hispanic	44.0	41.8 46.3	2,086	50.3	45.0 55.7	302	43.8	39.4 48.3	329	43.0	36.5 49.7	209
Hispanic	55.6	49.2 61.9	243	*	* *	*	*	* *	*	55.1	42.7 66.8	61
MARITAL STATUS												
Not Married	61.8	56.9 66.5	406	64.3	53.4 73.9	70	56.7	46.9 66.0	67	63.2	50.5 74.2	57
Married	40.1	37.9 42.4	1,926	44.7	39.1 50.5	246	39.6	35.0 44.4	273	38.5	32.4 45.0	213
INCOME												
Less than \$15,000	54.5	49.8 59.0	494	60.5	48.3 71.5	60	45.8	36.7 55.2	77	60.4	48.9 70.9	71
\$15,000 - \$24,999	47.2	42.1 52.4	385	59.8	46.9 71.4	52	33.7	22.9 46.5	42	30.0	17.8 46.0	39
\$25,000 - \$34,999	44.7	39.2 50.2	347	48.1	35.8 60.6	54	33.0	23.1 44.6	51	66.1	51.1 78.4	40
\$35,000 - \$49,999	42.9	37.5 48.5	346	45.3	32.0 59.3	43	52.4	41.9 62.7	58	37.9	22.9 55.7	30
\$50,000+	39.1	35.4 43.0	695	44.8	36.0 53.9	101	49.7	41.9 57.5	105	32.2	23.3 42.7	82
EDUCATION												
K-11th Grade	53.1	45.6 60.4	183	*	* *	*	*	* *	*	41.3	26.8 57.5	37
High School or GED	51.3	46.7 55.8	520	61.5	50.5 71.4	73	50.4	41.9 59.0	90	52.5	40.6 64.1	65
Some College	46.1	42.8 49.5	915	53.4	45.3 61.2	134	48.6	41.1 56.2	116	46.7	37.5 56.2	103
College Graduate+	37.4	33.7 41.2	708	36.3	27.6 46.0	88	33.9	27.4 41.1	122	38.9	28.0 51.1	62

*Figure not reliable by PRATS standards (n < 30)

HIV Testing

Idaho mothers who received an HIV test during pregnancy, 2010

	Public Health District															
	District 4				District 5				District 6				District 7			
	%	95% CI		n	%	95% CI		n	%	95% CI		n	%	95% CI		n
TOTAL	47.6	42.6	52.6	361	53.0	47.8	58.3	329	40.2	35.4	45.1	351	37.4	32.7	42.4	365
AGE																
18-19	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
20-24	55.4	43.8	66.4	69	65.7	55.0	75.0	82	36.6	27.6	46.6	90	40.5	32.1	49.4	117
25-29	48.1	39.9	56.3	135	47.4	38.8	56.1	121	36.5	29.1	44.6	133	37.3	30.0	45.3	146
30-34	45.2	35.9	54.8	100	51.6	40.3	62.7	70	39.6	30.3	49.8	84	27.5	18.8	38.3	71
35+	37.5	25.4	51.3	50	49.8	35.9	63.8	43	50.8	35.3	66.2	34	*	*	*	*
ETHNICITY																
Non-Hispanic	47.1	41.8	52.4	327	53.4	47.5	59.3	255	38.9	34.0	44.2	317	34.9	30.2	40.0	347
Hispanic	52.0	35.8	67.7	34	51.8	40.7	62.8	74	50.0	34.4	65.6	34	*	*	*	*
MARITAL STATUS																
Not Married	68.4	55.8	78.8	57	59.4	47.8	70.0	64	45.3	33.4	57.8	53	65.8	50.4	78.4	38
Married	41.8	36.5	47.2	304	50.2	44.6	55.8	265	38.6	33.6	43.8	298	32.1	27.5	37.1	327
INCOME																
Less than \$15,000	58.3	45.5	70.1	57	60.5	48.4	71.5	65	45.9	35.8	56.4	81	45.0	34.8	55.6	83
\$15,000 - \$24,999	55.7	42.6	68.0	54	64.1	52.7	74.1	72	36.0	25.9	47.5	68	36.0	25.0	48.8	58
\$25,000 - \$34,999	43.3	30.2	57.4	46	52.0	39.1	64.7	53	28.0	18.2	40.5	54	32.0	20.7	45.9	49
\$35,000 - \$49,999	53.5	39.5	67.0	46	35.0	24.1	47.8	55	41.2	29.4	54.0	51	36.5	26.1	48.4	63
\$50,000+	38.3	31.0	46.2	147	46.8	36.6	57.4	76	42.0	32.7	51.9	85	35.1	26.8	44.5	99
EDUCATION																
K-11th Grade	*	*	*	*	51.6	38.6	64.3	54	*	*	*	*	*	*	*	*
High School or GED	50.3	38.7	61.9	67	52.8	41.2	64.1	68	44.0	34.8	53.7	96	50.3	38.3	62.3	61
Some College	51.8	43.5	60.0	134	57.2	49.1	64.9	139	36.2	29.0	44.2	134	31.6	24.9	39.0	155
College Graduate+	38.4	30.8	46.6	137	45.7	34.9	57.0	68	39.0	30.4	48.4	97	32.4	25.3	40.4	134

*Figure not reliable by PRATS standards (n < 30)

No Dental Care

Idaho mothers who did not receive routine dental care during pregnancy, 2010

	Statewide			Public Health District								
				District 1			District 2			District 3		
	%	95% CI	n	%	95% CI	n	%	95% CI	n	%	95% CI	n
TOTAL	48.9	46.8 51.0	2,367	49.5	44.3 54.7	321	49.6	45.3 54.0	345	53.1	47.3 58.9	272
AGE												
18-19	69.9	59.0 79.0	80	*	* *	*	*	* *	*	*	* *	*
20-24	57.8	53.7 61.9	597	55.0	44.1 65.4	75	59.4	50.2 68.0	79	59.7	48.7 69.8	79
25-29	45.4	42.0 48.9	874	47.3	38.4 56.3	105	52.0	45.2 58.7	142	37.3	27.8 47.9	84
30-34	42.8	38.6 47.1	550	49.7	39.8 59.6	84	35.4	26.8 45.2	73	57.3	45.0 68.7	61
35+	43.8	37.6 50.1	266	42.0	28.8 56.5	41	45.1	32.6 58.2	38	62.5	46.9 75.9	38
ETHNICITY												
Non-Hispanic	46.8	44.6 49.0	2,116	49.0	43.8 54.4	307	48.7	44.3 53.2	334	50.7	44.1 57.3	209
Hispanic	63.6	57.4 69.5	248	*	* *	*	*	* *	*	60.6	48.3 71.7	63
MARITAL STATUS												
Not Married	59.6	54.6 64.4	410	52.9	42.1 63.3	70	55.2	45.5 64.6	67	60.3	47.8 71.6	58
Married	45.4	43.2 47.7	1,957	47.8	42.2 53.5	251	47.5	42.7 52.3	278	50.0	43.6 56.4	214
INCOME												
Less than \$15,000	62.9	58.4 67.2	499	48.4	36.7 60.3	60	54.4	45.0 63.4	77	63.0	51.5 73.3	72
\$15,000 - \$24,999	57.6	52.4 62.6	390	64.1	50.9 75.5	52	69.7	56.8 80.0	43	53.5	38.3 68.1	39
\$25,000 - \$34,999	55.1	49.7 60.5	356	60.3	47.8 71.6	55	56.4	45.2 66.9	53	58.4	43.3 72.1	41
\$35,000 - \$49,999	48.8	43.4 54.3	351	50.3	36.6 63.9	43	56.1	45.6 66.2	59	42.5	26.9 59.8	30
\$50,000+	27.7	24.4 31.2	705	34.3	26.3 43.1	105	28.8	22.2 36.4	106	41.4	31.4 52.0	82
EDUCATION												
K-11th Grade	71.7	64.7 77.8	185	*	* *	*	*	* *	*	57.0	41.1 71.6	37
High School or GED	58.7	54.2 63.1	524	58.2	47.2 68.4	74	59.4	50.8 67.4	91	60.2	48.2 71.0	66
Some College	50.2	46.8 53.5	935	53.3	45.2 61.2	134	52.5	45.0 59.9	119	54.8	45.4 63.9	104
College Graduate+	32.1	28.7 35.8	717	31.6	23.5 41.0	92	39.7	32.9 47.0	123	42.1	30.9 54.2	62

*Figure not reliable by PRATS standards (n < 30)

No Dental Care

Idaho mothers who did not receive routine dental care during pregnancy, 2010

	Public Health District															
	District 4				District 5				District 6				District 7			
	%	95% CI		n	%	95% CI		n	%	95% CI		n	%	95% CI		n
TOTAL	40.0	35.1	45.0	363	55.7	50.5	60.7	339	48.3	43.4	53.2	357	52.7	47.8	57.6	370
AGE																
18-19	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
20-24	54.9	43.5	65.9	70	55.5	45.1	65.5	84	57.3	47.5	66.6	92	61.4	52.6	69.6	118
25-29	39.5	31.8	47.8	136	54.0	45.4	62.4	122	48.4	40.4	56.3	135	48.9	41.3	56.6	150
30-34	28.9	21.0	38.2	100	59.8	48.9	69.7	76	36.2	27.2	46.2	86	43.5	33.0	54.7	70
35+	30.7	19.7	44.5	50	43.7	30.4	58.0	44	41.9	27.4	57.9	34	*	*	*	*
ETHNICITY																
Non-Hispanic	38.3	33.2	43.5	329	51.1	45.3	57.0	263	47.6	42.4	52.8	323	50.7	45.7	55.7	351
Hispanic	55.8	39.4	70.9	34	69.3	58.6	78.4	76	54.0	38.1	69.2	34	*	*	*	*
MARITAL STATUS																
Not Married	54.2	41.9	66.0	59	72.7	61.6	81.6	66	62.3	49.6	73.4	53	56.8	41.4	70.9	37
Married	35.9	30.8	41.2	304	48.0	42.5	53.6	273	44.1	39.0	49.3	304	52.0	46.8	57.0	333
INCOME																
Less than \$15,000	70.4	57.9	80.4	59	75.4	64.0	84.0	66	53.8	43.3	63.9	81	62.3	51.8	71.7	84
\$15,000 - \$24,999	50.5	37.7	63.2	55	55.1	43.9	65.9	73	65.4	54.1	75.2	69	60.4	47.8	71.8	59
\$25,000 - \$34,999	50.5	36.7	64.2	46	56.8	44.0	68.8	55	49.8	37.6	61.9	55	56.4	43.1	68.9	51
\$35,000 - \$49,999	45.5	32.1	59.5	46	48.1	36.2	60.2	58	51.9	39.5	64.2	52	52.4	40.7	63.8	63
\$50,000+	16.8	11.6	23.6	146	33.8	24.8	44.2	79	20.5	13.8	29.5	87	32.8	24.7	42.1	100
EDUCATION																
K-11th Grade	*	*	*	*	78.3	66.2	86.9	56	*	*	*	*	*	*	*	*
High School or GED	49.7	38.1	61.2	67	71.9	60.7	80.9	68	54.6	45.0	63.9	96	63.4	51.2	74.1	62
Some College	40.4	32.6	48.8	136	49.0	41.2	56.9	144	52.7	44.9	60.4	139	55.8	48.3	63.1	159
College Graduate+	24.2	17.9	31.8	137	27.3	18.6	38.3	71	27.9	20.3	36.9	98	40.5	32.8	48.6	134

*Figure not reliable by PRATS standards (n < 30)

Exercise

Idaho mothers who exercised during the third trimester¹, 2010

	Statewide			Public Health District												
				District 1			District 2			District 3						
	%	95% CI		n	%	95% CI		n	%	95% CI		n	%	95% CI		n
TOTAL	57.3	55.2	59.4	2,353	59.3	54.1	64.3	318	59.7	55.3	64.0	342	52.9	47.0	58.7	269
AGE																
18-19	69.5	57.9	79.1	79	*	*	*	*	*	*	*	*	*	*	*	*
20-24	61.1	56.9	65.1	592	55.7	44.9	66.1	75	59.8	50.4	68.6	77	58.7	47.6	68.9	78
25-29	54.4	50.9	57.9	870	63.1	54.0	71.4	104	59.8	52.9	66.3	142	42.1	32.2	52.7	83
30-34	56.9	52.6	61.2	552	53.7	43.7	63.5	85	62.5	53.0	71.1	73	56.6	44.3	68.2	60
35+	54.3	47.9	60.6	260	61.8	46.9	74.8	39	52.3	39.1	65.2	37	54.8	39.3	69.4	38
ETHNICITY																
Non-Hispanic	57.1	54.8	59.3	2,106	59.6	54.2	64.7	304	59.8	55.3	64.1	331	50.6	43.9	57.2	208
Hispanic	58.9	52.5	65.1	244	*	*	*	*	*	*	*	*	60.3	47.9	71.6	61
MARITAL STATUS																
Not Married	58.1	53.1	62.9	408	56.5	45.6	66.8	69	55.2	45.5	64.6	67	56.9	44.5	68.5	58
Married	57.1	54.8	59.3	1,945	60.6	54.9	66.1	249	61.5	56.7	66.0	275	51.2	44.7	57.6	211
INCOME																
Less than \$15,000	62.0	57.5	66.4	496	51.3	39.4	63.0	60	57.4	48.0	66.3	76	57.2	45.7	68.0	71
\$15,000 - \$24,999	61.0	55.8	65.9	386	68.7	54.9	79.8	50	45.1	33.2	57.6	43	54.0	38.8	68.4	39
\$25,000 - \$34,999	54.4	48.9	59.8	353	60.0	47.3	71.5	55	66.5	55.1	76.2	53	58.4	43.0	72.3	40
\$35,000 - \$49,999	52.3	46.8	57.8	348	60.5	46.0	73.3	42	57.7	46.8	67.9	57	*	*	*	*
\$50,000+	55.3	51.4	59.1	705	61.9	53.0	70.0	105	66.2	58.5	73.2	106	44.2	34.1	54.7	82
EDUCATION																
K-11th Grade	62.3	54.7	69.2	182	*	*	*	*	*	*	*	*	67.1	50.7	80.2	36
High School or GED	57.4	52.9	61.8	521	61.9	50.8	71.9	72	58.3	49.7	66.4	91	62.0	49.9	72.7	66
Some College	55.4	52.1	58.7	930	53.1	45.0	61.1	133	58.9	51.3	66.1	116	42.6	33.6	52.1	103
College Graduate+	58.3	54.4	62.1	714	67.4	57.9	75.6	92	64.3	57.0	70.9	123	50.8	38.9	62.7	61

*Figure not reliable by PRATS standards (n < 30)

1. Among mothers who exercised one or more days per week during pregnancy.

Exercise

Idaho mothers who exercised during the third trimester¹, 2010

	Public Health District															
	District 4				District 5				District 6				District 7			
	%	95% CI		n	%	95% CI		n	%	95% CI		n	%	95% CI		n
TOTAL	58.0	52.9	62.8	364	59.7	54.5	64.7	337	55.0	50.0	59.9	353	58.5	53.6	63.3	370
AGE																
18-19	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
20-24	64.6	52.8	74.8	69	55.0	44.5	65.0	84	58.5	48.6	67.7	91	67.9	59.2	75.4	118
25-29	53.7	45.5	61.7	137	66.9	58.3	74.5	122	48.7	40.7	56.7	133	53.7	46.0	61.3	149
30-34	63.3	53.8	71.9	101	49.3	38.4	60.1	75	57.9	47.9	67.3	86	50.9	40.0	61.8	72
35+	50.0	36.8	63.2	50	65.9	51.3	77.9	43	43.9	28.9	60.1	33	*	*	*	*
ETHNICITY																
Non-Hispanic	58.5	53.2	63.6	330	59.6	53.7	65.2	262	53.2	48.0	58.4	321	58.7	53.7	63.6	350
Hispanic	53.2	37.0	68.8	34	60.2	49.0	70.4	75	70.8	54.2	83.2	32	*	*	*	*
MARITAL STATUS																
Not Married	54.2	41.9	66.0	59	63.6	52.2	73.7	66	60.8	47.9	72.3	51	60.5	45.2	74.0	38
Married	59.0	53.6	64.2	305	57.9	52.3	63.3	271	53.3	48.1	58.5	302	58.1	53.0	63.1	332
INCOME																
Less than \$15,000	69.8	57.0	80.1	59	63.5	51.4	74.0	66	58.3	47.7	68.2	80	67.7	57.3	76.6	84
\$15,000 - \$24,999	63.2	50.0	74.7	55	68.8	57.8	77.9	73	52.1	40.8	63.2	67	61.4	48.8	72.6	59
\$25,000 - \$34,999	46.3	32.8	60.5	45	56.3	43.4	68.5	54	48.5	36.4	60.7	55	55.7	42.4	68.2	51
\$35,000 - \$49,999	44.6	31.4	58.5	47	45.0	33.3	57.2	58	55.8	43.1	67.7	52	58.7	46.9	69.6	63
\$50,000+	58.3	50.5	65.8	147	62.6	52.1	72.1	79	55.0	45.2	64.5	86	48.7	39.5	58.0	100
EDUCATION																
K-11th Grade	*	*	*	*	62.8	49.3	74.6	54	*	*	*	*	*	*	*	*
High School or GED	47.0	35.7	58.7	67	65.1	53.6	75.1	68	54.1	44.4	63.4	95	60.0	47.9	71.0	62
Some College	61.5	53.2	69.1	137	57.7	49.7	65.3	144	51.0	43.2	58.8	138	59.5	52.0	66.6	159
College Graduate+	58.4	50.2	66.1	137	54.6	43.6	65.2	71	60.0	50.6	68.7	97	57.8	49.6	65.5	133

*Figure not reliable by PRATS standards (n < 30)

1. Among mothers who exercised one or more days per week during pregnancy.

Stressful Life Events

Idaho mothers who experienced three or more stressful life events in the 12 months prior to delivery, 2010

	Statewide			Public Health District												
				District 1			District 2			District 3						
	%	95% CI		n	%	95% CI		n	%	95% CI		n	%	95% CI		n
TOTAL	27.4	25.5	29.4	2,370	27.9	23.4	33.0	321	24.9	21.1	29.0	344	31.5	26.2	37.3	271
AGE																
18-19	59.1	47.8	69.4	80	*	*	*	*	*	*	*	*	*	*	*	*
20-24	35.2	31.2	39.3	596	30.2	21.0	41.3	75	37.5	29.0	46.9	78	40.7	30.6	51.7	79
25-29	24.2	21.3	27.4	874	28.8	21.2	37.8	105	19.3	14.3	25.6	142	22.4	14.4	33.1	83
30-34	19.4	16.1	23.2	555	22.1	14.7	31.6	85	21.3	14.1	30.8	73	22.9	14.0	35.2	61
35+	24.1	19.0	30.1	265	20.1	10.4	35.3	40	13.9	6.9	26.0	38	37.0	23.3	53.2	38
ETHNICITY																
Non-Hispanic	26.5	24.6	28.6	2,119	28.4	23.7	33.6	307	24.9	21.1	29.2	333	27.5	21.9	34.0	209
Hispanic	33.3	27.5	39.6	248	*	*	*	*	*	*	*	*	43.7	32.0	56.1	62
MARITAL STATUS																
Not Married	48.9	44.0	53.9	410	42.9	32.7	53.7	70	46.3	36.8	56.0	67	46.6	34.6	58.9	58
Married	20.3	18.5	22.2	1,960	20.7	16.5	25.7	251	16.6	13.3	20.5	277	24.9	19.7	30.9	213
INCOME																
Less than \$15,000	44.5	39.9	49.1	500	44.3	32.9	56.3	60	43.2	34.2	52.6	77	52.5	41.1	63.6	72
\$15,000 - \$24,999	38.8	33.9	44.0	389	41.2	29.1	54.5	51	43.5	31.7	56.2	43	23.5	13.3	38.1	39
\$25,000 - \$34,999	29.8	24.9	35.2	356	38.5	27.1	51.3	56	22.1	14.0	33.1	53	37.1	23.7	52.8	40
\$35,000 - \$49,999	16.9	13.1	21.5	351	8.9	3.6	20.1	43	17.4	10.7	27.0	58	15.0	5.9	33.3	30
\$50,000+	9.4	7.3	12.0	708	9.8	5.6	16.7	105	6.5	3.6	11.4	106	14.3	8.2	23.7	82
EDUCATION																
K-11th Grade	43.7	36.5	51.2	185	*	*	*	*	*	*	*	*	49.6	34.2	65.0	37
High School or GED	39.0	34.7	43.6	524	24.2	16.0	34.8	74	36.1	28.2	44.8	91	47.9	36.4	59.7	66
Some College	27.3	24.4	30.5	936	38.1	30.6	46.4	134	29.6	23.0	37.1	118	23.1	15.9	32.4	103
College Graduate+	12.6	10.2	15.5	719	10.3	5.7	18.0	92	7.3	4.2	12.6	123	12.0	6.0	22.8	62

*Figure not reliable by PRATS standards (n < 30)

Stressful Life Events

Idaho mothers who experienced three or more stressful life events in the 12 months prior to delivery, 2010

	Public Health District															
	District 4				District 5				District 6				District 7			
	%	95% CI		n	%	95% CI		n	%	95% CI		n	%	95% CI		n
TOTAL	24.9	20.7	29.6	366	32.2	27.4	37.5	338	28.4	24.1	33.2	357	23.4	19.3	28.0	373
AGE																
18-19	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
20-24	31.2	21.4	43.0	70	44.6	34.5	55.2	84	39.1	30.0	49.1	92	27.6	20.2	36.6	118
25-29	24.1	17.7	32.0	137	24.6	17.5	33.3	121	27.7	20.9	35.6	135	22.6	16.6	30.0	151
30-34	20.3	13.6	29.1	102	24.3	15.6	35.9	76	15.1	9.1	24.1	86	10.8	5.7	19.7	72
35+	16.3	8.5	29.0	50	32.8	20.7	47.9	44	22.9	11.8	39.6	34	*	*	*	*
ETHNICITY																
Non-Hispanic	25.6	21.2	30.6	332	30.6	25.2	36.5	262	29.1	24.5	34.2	323	22.6	18.5	27.3	353
Hispanic	18.2	8.5	34.6	34	37.2	27.1	48.6	76	23.1	12.0	40.0	34	*	*	*	*
MARITAL STATUS																
Not Married	42.4	30.9	54.8	59	55.4	44.0	66.3	65	50.9	38.7	63.1	53	63.2	47.8	76.3	38
Married	19.9	15.9	24.5	307	22.0	17.7	26.9	273	21.7	17.7	26.3	304	16.1	12.7	20.2	335
INCOME																
Less than \$15,000	35.2	24.2	48.0	59	53.7	41.9	65.0	66	46.1	35.9	56.6	81	37.7	28.1	48.4	85
\$15,000 - \$24,999	48.0	35.4	60.8	55	44.9	34.0	56.2	73	41.2	30.7	52.7	69	27.6	17.7	40.2	59
\$25,000 - \$34,999	35.0	23.0	49.4	46	22.3	13.0	35.5	54	19.3	11.4	30.6	55	25.9	15.7	39.6	52
\$35,000 - \$49,999	15.8	7.9	29.3	47	21.5	13.0	33.5	58	23.1	14.2	35.3	52	15.9	9.0	26.4	63
\$50,000+	9.6	5.9	15.4	148	6.3	2.8	13.5	79	6.8	3.3	13.7	87	7.9	4.1	14.5	101
EDUCATION																
K-11th Grade	*	*	*	*	48.6	35.9	61.4	55	*	*	*	*	*	*	*	*
High School or GED	37.1	26.5	49.0	67	39.3	28.4	51.4	68	41.8	32.7	51.6	96	39.8	28.4	52.4	62
Some College	29.0	22.1	37.1	138	30.0	23.2	37.8	144	27.8	21.2	35.4	139	20.0	14.4	26.9	160
College Graduate+	13.1	8.4	19.8	138	12.2	6.4	22.0	71	15.5	9.7	23.9	98	13.1	8.6	19.5	135

*Figure not reliable by PRATS standards (n < 30)

Physical Abuse

Idaho mothers who experienced physical abuse during pregnancy, 2010

	Statewide			Public Health District								
				District 1			District 2			District 3		
	%	95% CI	n	%	95% CI	n	%	95% CI	n	%	95% CI	n
TOTAL	4.2	3.4 5.2	2,364	6.9	4.6 10.4	320	4.4	2.8 6.9	343	4.5	2.6 7.7	273
AGE												
18-19	14.9	8.6 24.5	80	*	* *	*	*	* *	*	*	* *	*
20-24	6.5	4.6 9.1	596	8.4	4.0 16.9	75	2.2	0.7 6.6	79	10.0	5.4 18.0	79
25-29	3.7	2.5 5.4	870	2.6	0.9 7.2	104	6.4	3.5 11.4	141	2.9	0.7 10.5	84
30-34	1.5	0.8 2.9	554	4.2	1.4 11.5	84	4.1	1.3 11.8	73	0.0	0.0 0.0	62
35+	2.4	1.1 5.3	264	4.6	1.3 15.0	41	0.0	0.0 0.0	37	0.0	0.0 0.0	38
ETHNICITY												
Non-Hispanic	4.0	3.2 5.1	2,110	6.9	4.5 10.5	306	4.6	2.9 7.1	332	4.2	2.2 7.7	209
Hispanic	5.4	3.1 9.1	251	*	* *	*	*	* *	*	5.5	1.8 15.3	64
MARITAL STATUS												
Not Married	8.5	6.1 11.7	409	13.0	7.3 22.3	69	12.1	7.0 20.1	66	5.2	1.8 14.3	58
Married	2.8	2.2 3.7	1,955	4.0	2.3 6.9	251	1.4	0.7 3.2	277	4.2	2.3 7.7	215
INCOME												
Less than \$15,000	9.0	6.7 12.1	496	12.9	6.6 23.8	59	6.8	3.1 14.2	75	9.3	4.7 17.3	72
\$15,000 - \$24,999	6.1	4.0 9.1	390	11.2	5.0 23.2	52	6.2	2.0 17.4	43	2.4	0.4 14.0	39
\$25,000 - \$34,999	3.1	1.5 6.2	355	6.5	2.6 15.1	55	2.8	0.6 12.5	53	5.8	1.5 19.9	41
\$35,000 - \$49,999	2.1	0.9 4.9	350	0.0	0.0 0.0	43	0.0	0.0 0.0	59	5.2	0.8 27.3	30
\$50,000+	0.8	0.4 1.7	706	1.8	0.5 6.3	105	3.7	1.7 8.0	106	0.0	0.0 0.0	82
EDUCATION												
K-11th Grade	6.6	3.8 11.4	187	*	* *	*	*	* *	*	5.3	1.3 18.5	38
High School or GED	6.2	4.2 9.1	524	9.7	4.7 19.2	74	2.9	0.9 8.5	91	10.7	5.2 20.6	66
Some College	3.9	2.8 5.5	930	7.5	4.1 13.4	133	10.4	6.3 16.6	117	0.9	0.1 5.6	104
College Graduate+	2.3	1.3 3.8	717	2.1	0.6 7.3	92	0.0	0.0 0.0	123	3.2	0.8 11.3	62

*Figure not reliable by PRATS standards (n < 30)

Physical Abuse

Idaho mothers who experienced physical abuse during pregnancy, 2010

	Public Health District															
	District 4				District 5				District 6				District 7			
	%	95% CI		n	%	95% CI		n	%	95% CI		n	%	95% CI		n
TOTAL	4.9	3.1	7.8	364	4.7	2.9	7.5	339	3.3	1.8	5.8	354	1.8	0.8	3.9	371
AGE																
18-19	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
20-24	8.7	3.8	18.8	70	9.5	4.9	17.6	83	0.9	0.2	5.6	92	2.8	0.9	8.2	118
25-29	5.1	2.3	10.7	135	2.8	0.9	8.0	123	4.4	1.9	9.8	134	2.3	0.8	6.7	149
30-34	1.9	0.5	6.9	102	1.2	0.2	7.0	76	1.1	0.2	6.7	85	0.0	0.0	0.0	72
35+	1.9	0.3	11.8	50	6.0	1.6	20.5	44	4.8	0.8	24.2	33	*	*	*	*
ETHNICITY																
Non-Hispanic	5.2	3.2	8.4	330	3.9	2.1	7.3	262	3.2	1.7	5.9	320	1.3	0.5	3.2	351
Hispanic	2.6	0.4	15.5	34	6.9	3.2	14.3	77	4.0	0.7	21.0	34	*	*	*	*
MARITAL STATUS																
Not Married	11.9	5.9	22.4	59	6.1	2.4	14.3	66	7.6	3.1	17.4	53	5.3	1.4	17.7	38
Married	3.0	1.6	5.5	305	4.0	2.3	6.9	273	2.0	1.0	4.1	301	1.2	0.5	3.0	333
INCOME																
Less than \$15,000	14.3	7.4	25.8	58	12.2	6.3	22.4	66	5.2	2.1	12.6	81	2.7	0.7	9.7	85
\$15,000 - \$24,999	9.6	4.1	20.8	55	5.5	2.4	12.0	73	6.1	2.4	14.6	69	1.6	0.2	9.5	59
\$25,000 - \$34,999	3.1	0.5	18.0	45	0.0	0.0	0.0	55	0.0	0.0	0.0	54	2.9	0.5	16.4	52
\$35,000 - \$49,999	2.0	0.3	12.1	47	1.7	0.3	9.8	58	0.0	0.0	0.0	51	3.2	0.9	11.3	62
\$50,000+	0.7	0.1	4.3	148	1.3	0.2	7.5	79	1.2	0.2	6.8	86	0.0	0.0	0.0	100
EDUCATION																
K-11th Grade	*	*	*	*	10.2	4.7	20.7	56	*	*	*	*	*	*	*	*
High School or GED	9.9	4.6	20.1	67	2.1	0.3	12.3	68	1.7	0.5	6.0	96	2.3	0.4	13.2	62
Some College	5.0	2.3	10.5	136	4.9	2.4	9.7	144	4.5	1.9	9.9	137	0.6	0.1	3.8	159
College Graduate+	2.1	0.7	6.2	138	1.4	0.2	8.2	71	2.8	0.7	9.8	97	2.6	0.9	7.7	134

*Figure not reliable by PRATS standards (n < 30)

Smoking

Idaho mothers who smoked during the third trimester¹, 2010

	Statewide			Public Health District								
				District 1			District 2			District 3		
	%	95% CI	n	%	95% CI	n	%	95% CI	n	%	95% CI	n
TOTAL	8.0	6.9 9.3	2,362	10.7	7.7 14.7	317	9.8	7.3 13.0	341	10.2	7.0 14.6	272
AGE												
18-19	20.0	12.5 30.4	80	*	* *	*	*	* *	*	*	* *	*
20-24	10.7	8.2 13.9	594	8.1	3.6 17.1	75	14.6	8.9 22.9	77	17.3	10.4 27.4	79
25-29	8.1	6.3 10.4	871	12.3	7.2 20.3	102	8.3	5.0 13.4	141	6.4	2.5 15.3	84
30-34	4.0	2.5 6.1	554	9.2	4.6 17.6	85	6.8	3.0 14.4	72	4.0	1.0 14.4	61
35+	5.2	2.9 9.2	263	6.6	1.8 21.6	39	7.4	3.0 17.4	38	2.4	0.4 14.3	38
ETHNICITY												
Non-Hispanic	9.0	7.7 10.4	2,110	11.1	8.0 15.3	303	10.2	7.6 13.5	330	13.5	9.3 19.2	209
Hispanic	1.6	0.6 4.1	249	*	* *	*	*	* *	*	0.0	0.0 0.0	63
MARITAL STATUS												
Not Married	18.7	15.1 22.9	408	20.0	12.7 30.0	70	21.2	14.3 30.3	66	20.7	12.4 32.5	58
Married	4.5	3.6 5.5	1,954	6.1	3.8 9.5	247	5.5	3.6 8.1	275	5.6	3.3 9.4	214
INCOME												
Less than \$15,000	18.9	15.5 22.9	499	22.7	14.3 34.2	60	24.8	17.5 33.9	77	25.8	17.1 37.1	72
\$15,000 - \$24,999	9.3	6.5 13.1	387	13.1	6.3 25.3	51	11.5	5.2 23.3	42	3.8	0.6 20.9	39
\$25,000 - \$34,999	5.4	3.4 8.5	354	10.4	4.6 21.8	55	7.3	3.3 15.2	51	8.0	2.6 21.7	41
\$35,000 - \$49,999	4.9	2.9 8.2	350	10.1	3.6 25.4	42	1.6	0.3 7.7	58	3.3	0.5 18.7	30
\$50,000+	0.9	0.5 1.9	706	0.9	0.2 5.5	103	2.4	0.8 7.3	106	0.0	0.0 0.0	82
EDUCATION												
K-11th Grade	17.4	12.4 23.8	185	*	* *	*	*	* *	*	21.5	11.2 37.2	37
High School or GED	15.2	12.2 18.9	521	19.2	11.8 29.6	73	18.1	12.2 26.0	90	13.5	7.0 24.5	66
Some College	6.9	5.3 9.1	934	9.2	5.3 15.4	133	7.5	4.2 12.8	117	9.1	4.7 17.0	104
College Graduate+	0.7	0.2 1.9	716	0.0	0.0 0.0	90	0.8	0.2 3.9	122	0.0	0.0 0.0	62

*Figure not reliable by PRATS standards (n < 30)

1. Smoking one or more cigarettes during the last three months of pregnancy.

Smoking

Idaho mothers who smoked during the third trimester¹, 2010

	Public Health District															
	District 4				District 5				District 6				District 7			
	%	95% CI		n	%	95% CI		n	%	95% CI		n	%	95% CI		n
TOTAL	8.3	5.8	11.8	366	6.3	4.0	9.8	337	8.7	6.1	12.2	356	4.5	2.7	7.3	373
AGE																
18-19	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
20-24	12.5	6.5	22.7	70	8.4	3.9	17.2	83	6.9	3.3	14.2	92	7.1	3.3	14.4	118
25-29	9.4	5.3	15.9	137	9.0	4.7	16.5	121	8.9	5.2	15.1	135	4.8	2.4	9.5	151
30-34	4.2	1.6	10.6	102	1.2	0.2	7.0	76	5.3	2.1	12.8	86	0.0	0.0	0.0	72
35+	6.7	2.2	18.5	50	6.0	1.6	20.5	44	7.6	2.0	24.4	33	*	*	*	*
ETHNICITY																
Non-Hispanic	9.2	6.4	13.0	332	6.8	4.2	11.1	260	9.2	6.5	13.0	323	4.8	2.9	7.8	353
Hispanic	0.0	0.0	0.0	34	4.8	1.6	13.3	77	4.1	0.7	21.5	33	*	*	*	*
MARITAL STATUS																
Not Married	20.3	12.2	32.0	59	14.1	7.8	24.1	64	18.9	10.9	30.6	53	15.8	7.6	30.1	38
Married	4.9	3.0	7.8	307	2.9	1.5	5.5	273	5.6	3.6	8.6	303	2.4	1.2	4.6	335
INCOME																
Less than \$15,000	20.4	11.8	32.8	59	16.5	9.3	27.7	65	18.7	11.6	28.8	81	8.0	3.7	16.4	85
\$15,000 - \$24,999	16.8	9.1	29.0	55	6.3	2.5	15.4	73	7.5	3.3	16.2	68	5.1	1.4	17.0	59
\$25,000 - \$34,999	2.1	0.3	12.5	46	3.1	0.5	16.8	54	8.8	3.9	18.4	55	2.9	0.5	16.4	52
\$35,000 - \$49,999	6.9	2.3	19.0	47	0.0	0.0	0.0	58	3.9	1.1	12.9	52	6.4	2.5	15.1	63
\$50,000+	0.7	0.1	4.3	148	1.3	0.2	7.5	79	2.3	0.6	7.9	87	1.0	0.2	6.1	101
EDUCATION																
K-11th Grade	*	*	*	*	7.6	2.9	18.1	56	*	*	*	*	*	*	*	*
High School or GED	11.9	5.9	22.5	67	11.1	5.2	22.0	67	20.2	13.4	29.3	96	16.4	8.7	28.7	62
Some College	10.5	6.3	17.1	138	6.0	3.0	12.0	143	3.4	1.5	7.5	139	1.8	0.6	5.2	160
College Graduate+	1.4	0.4	5.3	138	0.0	0.0	0.0	71	0.0	0.0	0.0	98	0.7	0.1	4.6	135

*Figure not reliable by PRATS standards (n < 30)

1. Smoking one or more cigarettes during the last three months of pregnancy.

Alcohol Use

Idaho mothers who drank alcohol during the third trimester¹, 2010

	Statewide			Public Health District								
				District 1			District 2			District 3		
	%	95% CI	n	%	95% CI	n	%	95% CI	n	%	95% CI	n
TOTAL	3.8	3.0 4.8	2,359	2.9	1.6 5.1	319	5.3	3.7 7.6	342	3.6	1.9 6.6	269
AGE												
18-19	1.3	0.2 7.8	80	*	* *	*	*	* *	*	*	* *	*
20-24	3.2	1.9 5.3	596	1.8	0.3 10.0	75	6.2	3.0 12.3	78	4.4	1.5 12.5	79
25-29	4.6	3.2 6.4	872	4.4	2.0 9.4	104	6.2	3.6 10.3	141	1.8	0.3 10.8	83
30-34	3.6	2.2 5.8	548	3.3	1.2 9.0	84	3.9	1.6 9.5	72	6.3	2.5 15.1	60
35+	4.1	2.1 7.7	263	2.4	0.4 13.2	40	5.0	1.6 14.3	38	2.5	0.4 14.6	37
ETHNICITY												
Non-Hispanic	3.7	2.9 4.7	2,110	2.8	1.5 5.0	305	5.5	3.8 7.9	331	4.0	2.0 7.8	207
Hispanic	4.6	2.5 8.3	246	*	* *	*	*	* *	*	2.1	0.3 12.8	62
MARITAL STATUS												
Not Married	6.0	4.0 8.9	409	1.4	0.2 8.2	70	3.0	1.0 8.8	67	5.2	1.8 14.3	58
Married	3.1	2.3 4.0	1,950	3.6	2.0 6.5	249	6.2	4.2 9.0	275	2.8	1.3 6.0	211
INCOME												
Less than \$15,000	3.3	1.9 5.6	499	0.0	0.0 0.0	60	2.7	0.9 8.2	77	3.4	0.9 12.1	72
\$15,000 - \$24,999	2.6	1.4 5.0	385	3.2	0.9 10.8	51	3.1	0.6 13.9	43	3.8	0.6 21.3	38
\$25,000 - \$34,999	4.8	2.8 8.0	356	5.9	2.0 16.0	56	6.9	3.1 14.6	53	2.3	0.3 13.5	40
\$35,000 - \$49,999	2.7	1.3 5.5	350	0.0	0.0 0.0	42	6.6	3.0 13.9	58	0.0	0.0 0.0	30
\$50,000+	4.9	3.4 7.1	703	4.6	2.1 10.0	104	7.5	4.3 12.8	104	4.7	1.9 11.6	81
EDUCATION												
K-11th Grade	3.7	1.8 7.5	183	*	* *	*	*	* *	*	0.0	0.0 0.0	37
High School or GED	3.9	2.3 6.3	522	1.9	0.3 10.6	74	1.8	0.6 5.5	91	4.2	1.1 14.6	65
Some College	3.4	2.2 5.0	933	3.7	1.8 7.7	134	5.0	2.6 9.4	118	4.1	1.6 10.3	102
College Graduate+	4.4	3.0 6.4	715	3.3	1.2 8.9	90	8.1	5.0 13.0	121	4.8	1.6 13.2	62

*Figure not reliable by PRATS standards (n < 30)

1. Consuming one or more alcoholic beverages during the last three months of pregnancy.

Alcohol Use

Idaho mothers who drank alcohol during the third trimester¹, 2010

	Public Health District															
	District 4				District 5				District 6				District 7			
	%	95% CI		n	%	95% CI		n	%	95% CI		n	%	95% CI		n
TOTAL	5.9	3.9	8.9	366	4.7	2.8	7.8	335	2.3	1.1	4.7	356	1.4	0.6	3.3	372
AGE																
18-19	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
20-24	4.4	1.4	12.5	70	4.6	1.5	12.7	84	0.0	0.0	0.0	92	2.0	0.5	7.6	118
25-29	6.4	3.3	12.1	137	9.1	4.9	16.3	121	4.9	2.1	10.7	135	0.6	0.1	4.0	151
30-34	6.1	2.8	12.8	102	0.0	0.0	0.0	73	0.0	0.0	0.0	86	1.4	0.2	8.4	71
35+	7.7	3.0	18.3	50	2.1	0.3	12.2	44	0.0	0.0	0.0	33	*	*	*	*
ETHNICITY																
Non-Hispanic	5.9	3.8	9.0	332	4.6	2.5	8.2	260	1.5	0.6	3.9	323	1.3	0.5	3.2	352
Hispanic	6.5	1.7	22.1	34	4.9	1.6	13.5	75	8.2	2.3	25.4	33	*	*	*	*
MARITAL STATUS																
Not Married	8.5	3.7	18.3	59	9.4	4.5	18.6	64	7.6	3.1	17.4	53	2.6	0.4	15.2	38
Married	5.2	3.3	8.2	307	2.6	1.3	5.1	271	0.7	0.2	2.4	303	1.2	0.5	3.0	334
INCOME																
Less than \$15,000	3.5	0.9	12.8	59	9.2	4.1	19.6	65	2.6	0.7	9.4	81	1.0	0.2	6.3	85
\$15,000 - \$24,999	0.0	0.0	0.0	55	4.2	1.2	14.2	71	6.7	2.4	17.4	68	0.0	0.0	0.0	59
\$25,000 - \$34,999	9.3	3.6	21.9	46	1.6	0.3	9.7	54	1.8	0.3	10.1	55	4.7	1.2	16.4	52
\$35,000 - \$49,999	4.9	1.3	17.3	47	5.0	1.7	13.6	58	0.0	0.0	0.0	52	1.6	0.3	9.6	63
\$50,000+	8.6	5.0	14.4	148	2.5	0.7	8.8	79	0.0	0.0	0.0	87	1.0	0.2	6.1	100
EDUCATION																
K-11th Grade	*	*	*	*	8.9	3.5	20.7	54	*	*	*	*	*	*	*	*
High School or GED	3.3	0.9	12.0	67	5.6	1.9	15.3	67	5.2	2.1	12.5	96	3.7	1.0	13.1	62
Some College	5.9	2.9	11.7	138	3.1	1.2	7.7	143	0.7	0.1	4.1	139	0.6	0.1	3.8	159
College Graduate+	7.8	4.4	13.3	138	2.8	0.8	9.6	71	1.7	0.3	9.8	98	0.7	0.1	4.6	135

*Figure not reliable by PRATS standards (n < 30)

1. Consuming one or more alcoholic beverages during the last three months of pregnancy.

Breastfeeding Initiation

Idaho mothers who ever breastfed, 2010

	Statewide			Public Health District								
				District 1			District 2			District 3		
	%	95% CI	n	%	95% CI	n	%	95% CI	n	%	95% CI	n
TOTAL	91.5	90.2 92.6	2,370	91.0	87.4 93.6	321	91.9	89.0 94.1	344	91.4	87.4 94.2	273
AGE												
18-19	88.6	79.2 94.0	80	*	* *	*	*	* *	*	*	* *	*
20-24	91.2	88.4 93.4	596	91.9	82.9 96.4	75	88.0	80.0 93.1	79	89.8	81.0 94.8	79
25-29	92.1	90.0 93.8	872	91.5	84.9 95.3	104	92.7	87.8 95.7	141	91.0	82.5 95.6	84
30-34	92.1	89.5 94.1	556	91.6	83.9 95.8	85	92.8	85.8 96.5	73	97.0	89.1 99.2	62
35+	89.9	85.3 93.1	266	93.2	82.2 97.6	41	92.6	82.7 97.0	38	88.0	74.7 94.8	38
ETHNICITY												
Non-Hispanic	91.9	90.6 93.1	2,116	91.1	87.4 93.7	307	92.5	89.6 94.6	333	92.0	87.5 95.0	209
Hispanic	88.5	83.8 91.9	251	*	* *	*	*	* *	*	89.3	79.3 94.8	64
MARITAL STATUS												
Not Married	88.2	84.7 91.0	410	87.1	78.1 92.8	70	84.9	76.4 90.6	66	89.7	79.4 95.1	58
Married	92.5	91.2 93.7	1,960	92.8	89.3 95.3	251	94.6	92.0 96.4	278	92.1	87.9 94.9	215
INCOME												
Less than \$15,000	89.7	86.4 92.2	498	89.2	78.7 94.8	59	86.8	78.6 92.1	76	92.7	83.8 96.9	72
\$15,000 - \$24,999	91.0	87.7 93.5	389	82.6	70.5 90.4	52	93.8	82.6 98.0	43	93.0	80.9 97.6	39
\$25,000 - \$34,999	89.3	85.3 92.3	357	88.2	76.9 94.4	56	93.8	85.1 97.6	53	81.9	66.9 91.0	41
\$35,000 - \$49,999	91.5	87.7 94.2	351	97.8	87.7 99.6	43	93.5	86.4 97.1	59	93.5	78.2 98.3	30
\$50,000+	94.3	92.2 95.8	708	95.4	90.1 98.0	105	93.0	87.6 96.2	106	92.3	84.1 96.4	82
EDUCATION												
K-11th Grade	87.4	81.7 91.6	187	*	* *	*	*	* *	*	90.7	77.8 96.4	38
High School or GED	87.0	83.6 89.8	523	85.1	75.1 91.5	74	88.8	82.1 93.2	91	91.2	80.8 96.2	66
Some College	91.0	88.8 92.7	936	91.3	85.4 95.0	134	93.6	88.2 96.7	118	88.4	80.9 93.3	104
College Graduate+	97.0	95.4 98.0	718	94.7	88.6 97.6	92	93.2	88.3 96.1	123	96.8	88.7 99.2	62

*Figure not reliable by PRATS standards (n < 30)

Breastfeeding Initiation

Idaho mothers who ever breastfed, 2010

	Public Health District															
	District 4				District 5				District 6				District 7			
	%	95% CI		n	%	95% CI		n	%	95% CI		n	%	95% CI		n
TOTAL	92.0	88.8	94.4	365	89.6	86.0	92.4	339	90.8	87.3	93.4	357	92.7	89.7	94.9	371
AGE																
18-19	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
20-24	87.5	77.3	93.6	70	89.6	81.3	94.4	83	92.1	84.8	96.1	92	96.4	91.0	98.6	118
25-29	95.3	90.0	97.8	136	91.0	84.3	95.0	123	87.4	80.5	92.1	135	92.2	86.7	95.6	149
30-34	92.0	85.0	95.9	102	87.3	78.9	92.6	76	93.3	86.5	96.8	86	90.5	82.0	95.3	72
35+	88.5	77.0	94.6	50	93.7	83.1	97.8	44	95.4	76.4	99.2	34	*	*	*	*
ETHNICITY																
Non-Hispanic	92.3	88.9	94.7	331	91.8	87.8	94.5	262	90.6	87.0	93.4	323	92.5	89.3	94.7	351
Hispanic	89.7	73.0	96.5	34	83.3	73.9	89.8	77	92.0	75.2	97.7	34	*	*	*	*
MARITAL STATUS																
Not Married	88.1	77.6	94.1	59	89.4	80.2	94.6	66	83.0	71.5	90.5	53	92.1	79.2	97.3	38
Married	93.1	89.8	95.4	306	89.7	85.8	92.7	273	93.1	89.9	95.3	304	92.8	89.7	95.0	333
INCOME																
Less than \$15,000	87.4	76.1	93.8	59	83.8	72.8	90.9	66	86.9	77.9	92.6	81	95.7	87.9	98.6	85
\$15,000 - \$24,999	93.6	82.4	97.9	55	92.3	85.2	96.2	73	88.2	78.0	94.1	69	91.0	80.5	96.1	58
\$25,000 - \$34,999	89.5	77.5	95.4	45	88.7	78.8	94.3	55	96.5	88.1	99.0	55	91.0	80.8	96.1	52
\$35,000 - \$49,999	87.1	74.5	94.0	47	95.0	86.4	98.3	58	90.4	80.0	95.7	52	90.3	80.7	95.4	62
\$50,000+	95.4	90.8	97.7	148	93.7	86.5	97.2	79	95.4	89.2	98.2	87	93.1	86.7	96.6	101
EDUCATION																
K-11th Grade	*	*	*	*	86.2	74.3	93.1	56	*	*	*	*	*	*	*	*
High School or GED	86.8	76.1	93.2	67	86.5	76.9	92.5	68	84.2	75.8	90.0	96	86.3	76.0	92.6	61
Some College	91.0	85.3	94.7	137	89.5	83.8	93.3	144	93.8	89.1	96.6	139	91.5	86.1	94.9	160
College Graduate+	97.2	92.9	98.9	138	97.3	90.4	99.3	71	100.0	100.0	100.0	98	97.1	92.8	98.8	134

*Figure not reliable by PRATS standards (n < 30)

Breastfeeding Duration

Idaho mothers who breastfed for at least six months¹, 2010

	Statewide			Public Health District												
				District 1			District 2			District 3						
	%	95% CI		n	%	95% CI		n	%	95% CI		n	%	95% CI		n
TOTAL	53.0	50.8	55.2	2,208	56.3	50.8	61.7	293	57.6	53.0	62.0	325	49.3	43.3	55.3	255
AGE																
18-19	28.6	19.5	39.9	77	*	*	*	*	*	*	*	*	*	*	*	*
20-24	44.3	40.0	48.7	549	45.1	34.4	56.4	69	44.4	35.2	54.0	72	39.9	29.5	51.4	71
25-29	55.9	52.2	59.5	807	59.0	49.4	68.0	95	61.3	54.2	67.9	136	53.5	42.6	64.1	77
30-34	60.3	55.8	64.6	525	69.0	58.3	78.1	77	71.8	61.9	79.9	68	61.0	48.6	72.1	60
35+	57.9	51.3	64.2	250	70.1	54.7	81.9	37	56.3	42.7	69.0	36	52.2	36.7	67.3	37
ETHNICITY																
Non-Hispanic	54.2	51.8	56.5	1,976	56.4	50.8	61.9	279	58.2	53.6	62.7	318	54.4	47.5	61.2	194
Hispanic	45.0	38.6	51.6	229	*	*	*	*	*	*	*	*	34.3	23.9	46.4	61
MARITAL STATUS																
Not Married	34.7	30.0	39.9	373	39.1	28.8	50.5	64	32.8	24.0	43.0	61	34.7	23.2	48.3	49
Married	58.9	56.5	61.2	1,835	64.6	58.7	70.1	229	66.7	61.9	71.1	264	54.9	48.3	61.3	206
INCOME																
Less than \$15,000	41.0	36.3	45.8	452	43.0	31.2	55.6	54	32.7	24.4	42.4	70	33.2	22.9	45.4	64
\$15,000 - \$24,999	49.4	44.1	54.7	365	41.4	29.4	54.4	51	55.1	42.3	67.2	42	61.1	44.4	75.5	35
\$25,000 - \$34,999	52.4	46.7	58.0	327	57.7	44.0	70.4	47	65.4	53.5	75.7	49	47.4	32.8	62.5	39
\$35,000 - \$49,999	54.4	48.7	60.0	329	59.9	44.6	73.4	37	68.5	57.2	77.9	54	*	*	*	*
\$50,000+	64.6	60.7	68.3	675	71.7	62.8	79.2	98	69.5	61.7	76.3	103	59.5	48.7	69.5	79
EDUCATION																
K-11th Grade	42.6	35.0	50.5	164	*	*	*	*	*	*	*	*	39.0	24.2	56.2	32
High School or GED	40.0	35.6	44.7	487	49.4	38.1	60.8	66	39.0	30.8	47.9	84	42.9	31.6	55.0	63
Some College	54.2	50.7	57.6	870	52.1	43.6	60.5	122	56.0	48.3	63.5	112	55.4	45.7	64.7	99
College Graduate+	65.2	61.2	68.9	682	72.3	62.7	80.1	87	76.9	70.0	82.7	118	55.9	43.6	67.6	59

*Figure not reliable by PRATS standards (n < 30)

1. Babies who were not at least six months old at the time of the survey were excluded from analysis.

Breastfeeding Duration

Idaho mothers who breastfed for at least six months¹, 2010

	Public Health District															
	District 4				District 5				District 6				District 7			
	%	95% CI		n	%	95% CI		n	%	95% CI		n	%	95% CI		n
TOTAL	55.8	50.6	60.9	343	46.2	40.9	51.6	314	53.0	47.8	58.1	327	54.3	49.3	59.3	351
AGE																
18-19	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
20-24	44.1	32.8	56.0	65	33.5	24.2	44.4	77	45.7	35.9	55.9	84	53.1	44.0	61.9	111
25-29	63.0	54.3	71.0	124	52.9	43.9	61.8	112	48.7	40.4	57.0	123	51.7	43.7	59.6	140
30-34	56.3	46.7	65.5	100	49.0	37.9	60.2	71	67.1	56.8	76.0	79	63.0	51.6	73.0	70
35+	57.2	43.1	70.1	47	53.4	38.8	67.4	41	62.5	45.3	77.0	31	*	*	*	*
ETHNICITY																
Non-Hispanic	55.4	49.9	60.8	312	48.1	42.1	54.2	246	54.2	48.8	59.6	296	53.4	48.1	58.5	331
Hispanic	59.2	41.6	74.7	31	40.1	29.4	51.9	68	42.6	27.2	59.6	31	*	*	*	*
MARITAL STATUS																
Not Married	35.1	24.2	47.8	57	34.5	24.0	46.7	58	29.8	19.2	43.1	47	35.1	22.2	50.7	37
Married	61.9	56.3	67.2	286	51.2	45.4	56.9	256	59.6	54.2	64.8	280	58.0	52.7	63.1	314
INCOME																
Less than \$15,000	45.5	33.0	58.6	54	33.5	22.8	46.1	58	44.4	33.9	55.5	72	48.4	37.9	59.0	80
\$15,000 - \$24,999	53.6	40.1	66.5	51	39.2	28.5	50.9	68	49.3	37.9	60.8	64	49.1	36.6	61.8	54
\$25,000 - \$34,999	52.8	38.4	66.8	43	36.4	24.8	49.9	50	60.1	46.7	72.1	49	57.5	44.0	69.9	50
\$35,000 - \$49,999	35.2	22.9	49.7	44	63.2	50.5	74.4	55	56.0	43.1	68.1	50	63.3	51.2	74.0	60
\$50,000+	69.5	61.6	76.4	141	64.4	53.7	73.8	75	59.2	49.1	68.7	82	57.0	47.5	66.0	97
EDUCATION																
K-11th Grade	*	*	*	*	36.3	23.9	50.7	45	*	*	*	*	*	*	*	*
High School or GED	43.6	32.1	55.9	62	34.9	24.7	46.8	67	37.0	28.0	46.9	87	32.8	22.4	45.3	58
Some College	57.0	48.2	65.3	125	47.7	39.6	56.0	133	57.6	49.3	65.5	126	53.0	45.3	60.5	153
College Graduate+	61.9	53.6	69.6	133	65.1	53.6	75.0	69	71.7	62.2	79.6	92	66.0	57.7	73.4	124

*Figure not reliable by PRATS standards (n < 30)

1. Babies who were not at least six months old at the time of the survey were excluded from analysis.

Self-Reported Postpartum Depression

Idaho mothers who reported being moderately or severely depressed during the three months after pregnancy¹, 2010

	Statewide			Public Health District								
				District 1			District 2			District 3		
	%	95% CI	n	%	95% CI	n	%	95% CI	n	%	95% CI	n
TOTAL	16.8	15.3 18.5	2,361	12.7	9.5 16.8	320	17.1	13.9 20.8	342	16.2	12.2 21.2	272
AGE												
18-19	22.1	13.9 33.1	80	*	* *	*	*	* *	*	*	* *	*
20-24	19.2	16.1 22.8	595	14.7	8.4 24.5	75	21.2	14.7 29.6	79	23.4	15.2 34.1	79
25-29	15.8	13.4 18.6	867	13.0	8.1 20.2	103	11.5	7.8 16.7	140	11.3	6.1 20.1	83
30-34	13.5	10.8 16.8	553	6.0	2.4 14.2	85	20.0	13.2 29.2	72	8.5	3.7 18.7	62
35+	19.3	14.6 25.1	266	13.0	5.8 26.7	41	15.3	7.6 28.4	38	23.6	12.5 39.9	38
ETHNICITY												
Non-Hispanic	16.7	15.0 18.5	2,108	13.0	9.7 17.3	306	17.0	13.8 20.7	332	15.9	11.4 21.7	208
Hispanic	17.8	13.4 23.3	250	*	* *	*	*	* *	*	17.0	9.5 28.5	64
MARITAL STATUS												
Not Married	23.2	19.3 27.7	410	20.0	12.7 30.0	70	25.8	18.1 35.2	66	27.6	18.0 39.9	58
Married	14.7	13.2 16.4	1,951	9.2	6.4 13.1	250	13.8	10.8 17.4	276	11.2	7.7 16.0	214
INCOME												
Less than \$15,000	22.8	19.1 26.9	497	17.8	10.4 28.8	59	26.5	19.0 35.8	75	22.6	14.3 33.8	72
\$15,000 - \$24,999	19.2	15.3 23.7	388	23.5	13.8 37.0	52	15.2	8.0 26.9	43	7.0	2.4 19.1	39
\$25,000 - \$34,999	17.1	13.4 21.6	356	12.2	6.2 22.7	56	18.0	10.7 28.5	52	10.2	3.9 23.9	41
\$35,000 - \$49,999	16.5	12.5 21.3	347	0.0	0.0 0.0	42	9.7	5.1 17.6	59	18.3	8.0 36.5	30
\$50,000+	11.1	8.9 13.7	707	6.4	3.3 12.2	105	13.5	8.9 19.8	106	13.1	7.3 22.4	82
EDUCATION												
K-11th Grade	22.5	16.8 29.3	187	*	* *	*	*	* *	*	13.8	5.9 28.7	38
High School or GED	19.6	16.1 23.6	522	13.0	7.1 22.6	74	18.6	12.8 26.2	91	22.9	14.2 34.8	66
Some College	16.1	13.8 18.7	930	16.1	10.7 23.3	133	23.4	17.5 30.5	118	13.3	7.8 21.7	103
College Graduate+	13.6	11.1 16.6	716	6.4	3.1 12.7	92	8.6	5.2 13.8	121	13.6	7.1 24.6	62

*Figure not reliable by PRATS standards (n < 30)

1. During the three months after delivery.

Self-Reported Postpartum Depression

Idaho mothers who reported being moderately or severely depressed during the three months after pregnancy¹, 2010

	Public Health District															
	District 4				District 5				District 6				District 7			
	%	95% CI		n	%	95% CI		n	%	95% CI		n	%	95% CI		n
TOTAL	17.5	14.0	21.7	364	19.5	15.7	24.1	336	17.6	14.1	21.8	356	16.2	12.9	20.2	371
AGE																
18-19	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
20-24	13.7	7.4	24.1	70	24.2	16.1	34.8	83	15.6	9.5	24.6	91	21.0	14.8	29.0	118
25-29	17.2	11.8	24.4	136	18.5	12.8	26.1	121	17.9	12.3	25.2	135	16.5	11.4	23.1	149
30-34	17.2	11.2	25.4	101	18.1	10.8	28.8	75	20.7	13.7	30.2	86	5.4	2.2	13.0	72
35+	21.1	12.0	34.6	50	19.0	10.5	31.9	44	19.0	9.8	33.8	34	*	*	*	*
ETHNICITY																
Non-Hispanic	18.0	14.3	22.5	330	19.0	14.7	24.2	259	16.9	13.3	21.2	322	15.9	12.5	20.0	351
Hispanic	13.0	5.1	29.4	34	21.2	13.4	31.8	77	23.1	12.0	40.0	34	*	*	*	*
MARITAL STATUS																
Not Married	20.3	12.2	32.0	59	24.2	15.9	35.2	66	24.5	15.4	36.8	53	21.1	11.3	35.9	38
Married	16.7	13.1	21.2	305	17.4	13.6	22.1	270	15.5	12.1	19.7	303	15.3	12.0	19.4	333
INCOME																
Less than \$15,000	24.7	15.5	36.9	59	24.6	16.0	36.0	66	22.8	15.2	32.9	81	21.1	13.7	31.0	85
\$15,000 - \$24,999	27.2	17.2	40.1	55	23.2	14.9	34.3	73	15.5	8.5	26.4	68	14.5	7.9	25.0	58
\$25,000 - \$34,999	17.9	9.4	31.5	45	17.3	9.4	29.7	55	18.8	10.8	30.6	55	24.5	15.0	37.2	52
\$35,000 - \$49,999	24.2	14.1	38.3	46	15.4	8.1	27.2	56	19.2	11.2	31.1	52	14.5	8.0	25.0	62
\$50,000+	9.3	5.7	14.8	148	16.5	10.1	25.7	78	14.8	9.1	23.1	87	9.8	5.5	16.9	101
EDUCATION																
K-11th Grade	*	*	*	*	21.1	12.1	34.2	56	*	*	*	*	*	*	*	*
High School or GED	24.5	15.7	36.1	67	14.5	7.8	25.2	67	15.8	10.0	24.2	96	22.0	13.4	33.9	61
Some College	14.0	9.2	20.8	136	22.4	16.5	29.7	142	18.9	13.5	25.8	138	13.2	9.0	18.9	160
College Graduate+	15.9	10.8	22.8	138	17.7	10.6	28.0	71	14.5	8.9	22.8	98	12.5	8.1	18.8	134

*Figure not reliable by PRATS standards (n < 30)

1. During the three months after delivery.

No Well-Baby Care

Idaho mothers who never took their baby for routine well-baby care, 2010

	Statewide			Public Health District								
				District 1			District 2			District 3		
	%	95% CI	n	%	95% CI	n	%	95% CI	n	%	95% CI	n
TOTAL	2.1	1.6 2.8	2,262	2.2	1.2 4.2	308	1.4	0.7 2.8	327	1.4	0.6 3.6	251
AGE												
18-19	3.2	0.8 11.5	77	*	* *	*	*	* *	*	*	* *	*
20-24	1.5	0.8 2.9	556	0.0	0.0 0.0	70	2.4	0.8 7.2	71	1.2	0.2 7.6	68
25-29	2.7	1.7 4.2	845	5.4	2.6 10.8	101	2.0	0.8 4.8	139	0.0	0.0 0.0	78
30-34	1.8	1.0 3.4	536	1.1	0.2 6.4	84	0.0	0.0 0.0	68	3.2	0.9 11.5	59
35+	1.9	0.8 4.4	248	2.4	0.4 13.5	38	0.0	0.0 0.0	36	2.6	0.4 15.1	36
ETHNICITY												
Non-Hispanic	1.8	1.3 2.5	2,036	2.3	1.2 4.4	294	1.4	0.7 2.9	316	0.0	0.0 0.0	196
Hispanic	4.6	2.6 8.0	223	*	* *	*	*	* *	*	6.2	2.4 15.0	55
MARITAL STATUS												
Not Married	1.9	0.9 4.2	381	0.0	0.0 0.0	68	0.0	0.0 0.0	65	0.0	0.0 0.0	50
Married	2.2	1.6 3.0	1,881	3.3	1.8 6.2	240	1.9	0.9 3.9	262	2.0	0.8 5.0	201
INCOME												
Less than \$15,000	1.0	0.4 2.4	460	0.0	0.0 0.0	57	0.0	0.0 0.0	70	1.3	0.2 7.8	62
\$15,000 - \$24,999	2.5	1.2 5.2	376	1.6	0.3 9.3	49	2.0	0.4 9.3	42	2.4	0.4 14.0	39
\$25,000 - \$34,999	3.4	1.9 6.0	339	5.1	1.8 13.7	52	7.3	3.3 15.4	50	2.4	0.4 14.1	38
\$35,000 - \$49,999	2.7	1.3 5.7	342	0.0	0.0 0.0	41	0.0	0.0 0.0	58	*	* *	*
\$50,000+	1.4	0.8 2.5	691	3.7	1.5 8.9	103	0.0	0.0 0.0	101	1.2	0.2 7.6	79
EDUCATION												
K-11th Grade	6.2	3.4 11.1	162	*	* *	*	*	* *	*	9.8	3.8 22.9	32
High School or GED	2.7	1.5 4.7	486	0.0	0.0 0.0	67	2.0	0.6 5.9	83	0.0	0.0 0.0	59
Some College	1.8	1.1 3.0	906	1.9	0.7 5.3	132	1.6	0.5 4.7	115	0.0	0.0 0.0	97
College Graduate+	0.9	0.5 1.8	702	4.4	1.8 10.4	89	0.8	0.2 4.0	118	0.0	0.0 0.0	60

*Figure not reliable by PRATS standards (n < 30)

No Well-Baby Care

Idaho mothers who never took their baby for routine well-baby care, 2010

	Public Health District															
	District 4				District 5				District 6				District 7			
	%	95% CI		n	%	95% CI		n	%	95% CI		n	%	95% CI		n
TOTAL	2.1	1.0	4.3	354	2.2	1.1	4.1	319	2.0	0.9	4.2	341	2.9	1.6	5.3	362
AGE																
18-19	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
20-24	2.6	0.7	9.6	67	1.0	0.2	6.2	79	2.7	0.7	9.5	88	0.8	0.1	5.1	113
25-29	3.5	1.3	8.8	131	3.0	1.2	7.3	118	2.7	0.9	7.6	129	2.3	0.8	6.7	149
30-34	1.0	0.1	6.1	100	1.2	0.2	7.4	71	1.2	0.2	6.8	84	4.2	1.4	11.6	70
35+	0.0	0.0	0.0	49	4.8	1.3	16.1	40	0.0	0.0	0.0	30	*	*	*	*
ETHNICITY																
Non-Hispanic	2.3	1.1	4.8	320	1.7	0.8	3.9	255	0.9	0.3	2.5	311	2.7	1.4	4.9	344
Hispanic	0.0	0.0	0.0	34	3.7	1.3	10.2	64	11.7	4.1	29.2	30	*	*	*	*
MARITAL STATUS																
Not Married	3.6	1.0	12.9	55	0.0	0.0	0.0	59	4.1	1.1	13.7	49	5.7	1.5	19.1	35
Married	1.7	0.7	3.8	299	3.1	1.6	5.8	260	1.4	0.6	3.4	292	2.5	1.3	4.7	327
INCOME																
Less than \$15,000	1.5	0.2	9.4	56	2.4	0.7	8.5	59	1.1	0.2	6.3	74	0.0	0.0	0.0	82
\$15,000 - \$24,999	5.8	1.9	16.3	53	0.0	0.0	0.0	68	2.3	0.4	12.7	67	1.6	0.3	9.7	58
\$25,000 - \$34,999	2.2	0.3	13.0	44	3.3	0.9	11.6	53	1.9	0.3	10.6	52	4.9	1.3	16.9	50
\$35,000 - \$49,999	5.2	1.3	18.1	45	3.4	0.9	11.7	57	2.0	0.3	11.2	51	3.2	0.9	11.3	62
\$50,000+	0.0	0.0	0.0	147	2.5	0.7	8.9	78	1.2	0.2	6.9	85	3.0	1.0	8.6	98
EDUCATION																
K-11th Grade	*	*	*	*	1.6	0.3	9.6	46	*	*	*	*	*	*	*	*
High School or GED	3.4	0.9	12.3	65	4.9	1.9	11.7	64	0.9	0.1	5.2	93	7.3	2.8	17.6	55
Some College	3.1	1.2	7.9	132	0.7	0.1	4.0	139	1.4	0.4	5.1	132	2.4	1.0	6.0	159
College Graduate+	0.0	0.0	0.0	135	2.8	0.8	9.7	70	1.0	0.2	6.0	97	0.7	0.1	4.6	133

*Figure not reliable by PRATS standards (n < 30)

Hearing Screening

Idaho mothers who reported their baby's hearing was tested, 2010

	Statewide			Public Health District												
				District 1			District 2			District 3						
	%	95% CI		n	%	95% CI		n	%	95% CI		n	%	95% CI		n
TOTAL	95.1	94.1	95.9	2,366	93.7	90.8	95.8	319	91.8	89.1	93.9	343	91.6	87.6	94.4	273
AGE																
18-19	88.3	78.7	93.9	79	*	*	*	*	*	*	*	*	*	*	*	*
20-24	94.8	92.5	96.4	596	96.2	89.3	98.7	75	92.8	86.6	96.3	79	88.5	79.0	94.1	79
25-29	96.0	94.5	97.2	870	92.3	85.8	95.9	103	95.1	91.2	97.3	141	96.7	90.7	98.9	84
30-34	96.2	94.3	97.4	555	93.3	86.7	96.8	84	87.1	79.7	92.1	73	95.4	87.2	98.5	62
35+	92.6	88.4	95.4	266	95.5	85.0	98.7	41	86.2	74.0	93.2	38	85.1	69.3	93.5	38
ETHNICITY																
Non-Hispanic	95.7	94.7	96.5	2,112	93.4	90.3	95.6	305	91.9	89.2	94.0	332	94.0	89.9	96.6	209
Hispanic	90.7	86.1	93.9	251	*	*	*	*	*	*	*	*	84.3	72.9	91.5	64
MARITAL STATUS																
Not Married	93.8	90.9	95.8	407	95.7	88.3	98.5	69	93.9	87.0	97.2	65	86.2	75.3	92.8	58
Married	95.5	94.5	96.3	1,959	92.8	89.2	95.3	250	91.0	87.9	93.4	278	94.0	90.1	96.4	215
INCOME																
Less than \$15,000	94.6	92.1	96.4	495	92.0	82.7	96.5	58	92.9	86.8	96.3	75	91.0	81.6	95.8	72
\$15,000 - \$24,999	94.6	91.8	96.4	389	94.2	84.2	98.0	52	95.0	85.2	98.4	43	88.3	75.2	94.9	39
\$25,000 - \$34,999	93.6	90.3	95.8	357	98.4	91.0	99.7	56	85.1	75.4	91.4	53	92.0	78.3	97.4	41
\$35,000 - \$49,999	94.7	91.8	96.6	350	86.4	74.1	93.4	42	82.9	73.5	89.5	59	96.7	81.3	99.5	30
\$50,000+	97.3	95.8	98.3	708	95.4	90.1	98.0	105	97.2	93.2	98.9	106	94.6	86.7	97.9	82
EDUCATION																
K-11th Grade	88.7	83.0	92.6	186	*	*	*	*	*	*	*	*	84.2	69.3	92.7	38
High School or GED	94.4	92.0	96.2	521	88.6	80.0	93.8	74	87.9	81.4	92.3	90	91.2	80.8	96.2	66
Some College	96.4	94.9	97.5	936	96.4	92.0	98.4	134	97.0	93.4	98.6	118	92.8	85.8	96.5	104
College Graduate+	95.8	94.1	97.0	717	92.5	85.8	96.2	91	90.0	84.6	93.6	123	95.3	86.8	98.4	62

*Figure not reliable by PRATS standards (n < 30)

Hearing Screening

Idaho mothers who reported their baby's hearing was tested, 2010

	Public Health District															
	District 4				District 5				District 6				District 7			
	%	95% CI		n	%	95% CI		n	%	95% CI		n	%	95% CI		n
TOTAL	97.1	94.9	98.4	365	94.1	91.0	96.2	339	97.4	95.1	98.7	356	96.0	93.7	97.6	371
AGE																
18-19	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
20-24	97.5	90.9	99.3	70	94.3	87.2	97.6	83	99.1	94.4	99.9	92	95.6	90.1	98.1	118
25-29	96.3	91.5	98.4	136	94.4	88.9	97.3	123	98.1	93.2	99.5	134	96.8	92.9	98.6	149
30-34	99.1	94.0	99.9	102	97.9	87.7	99.7	76	93.9	85.5	97.6	86	95.9	88.7	98.6	72
35+	96.2	86.4	99.0	50	93.7	83.1	97.8	44	97.3	85.0	99.6	34	*	*	*	*
ETHNICITY																
Non-Hispanic	97.3	95.0	98.6	331	95.7	92.5	97.6	262	97.6	95.2	98.8	322	95.8	93.2	97.4	351
Hispanic	94.8	82.0	98.6	34	89.4	80.3	94.5	77	96.0	79.0	99.3	34	*	*	*	*
MARITAL STATUS																
Not Married	98.3	89.6	99.8	59	92.4	83.8	96.6	66	94.2	84.7	98.0	52	97.4	84.8	99.6	38
Married	96.7	94.2	98.2	306	94.9	91.8	96.8	273	98.4	96.3	99.3	304	95.8	93.2	97.4	333
INCOME																
Less than \$15,000	95.8	88.0	98.6	59	96.9	88.9	99.2	66	98.0	93.1	99.5	80	95.3	88.4	98.2	85
\$15,000 - \$24,999	96.8	88.5	99.2	55	92.6	83.7	96.8	73	93.0	83.5	97.2	69	100.0	100.0	100.0	58
\$25,000 - \$34,999	91.6	80.1	96.7	45	93.5	84.7	97.4	55	97.0	83.8	99.5	55	94.6	85.3	98.2	52
\$35,000 - \$49,999	97.1	82.9	99.6	47	95.2	83.3	98.8	58	98.1	89.0	99.7	52	95.2	86.7	98.3	62
\$50,000+	99.3	95.7	99.9	148	96.2	89.7	98.7	79	100.0	100.0	100.0	87	96.1	90.4	98.4	101
EDUCATION																
K-11th Grade	*	*	*	*	86.0	74.7	92.8	56	*	*	*	*	*	*	*	*
High School or GED	96.7	88.0	99.2	67	96.6	90.4	98.8	68	98.3	93.9	99.5	95	96.3	86.7	99.0	61
Some College	97.3	93.3	99.0	137	97.6	92.9	99.2	144	96.8	92.0	98.7	139	97.0	93.3	98.7	160
College Graduate+	98.6	94.8	99.6	138	91.9	82.5	96.5	71	100.0	100.0	100.0	98	94.1	89.1	96.9	134

*Figure not reliable by PRATS standards (n < 30)

1. Among mothers who were working or attending school.

Immunization Information from PNC Provider

Idaho mothers who reported receiving immunization information from PNC providers, 2010

	Statewide			Public Health District								
				District 1			District 2			District 3		
	%	95% CI	n	%	95% CI	n	%	95% CI	n	%	95% CI	n
TOTAL	70.7	68.6 72.6	2,190	78.4	73.7 82.4	302	76.7	72.7 80.2	327	70.5	64.7 75.6	252
AGE												
18-19	67.2	55.9 76.8	78	*	* *	*	*	* *	*	*	* *	*
20-24	74.4	70.5 78.0	562	87.3	77.9 93.1	72	78.5	69.6 85.3	74	73.0	62.2 81.6	77
25-29	69.5	66.0 72.7	817	80.4	72.0 86.7	99	77.8	71.7 82.9	138	65.4	54.7 74.7	80
30-34	67.4	63.0 71.5	498	70.3	60.1 78.8	79	72.5	63.0 80.3	67	68.5	55.4 79.1	56
35+	73.5	67.2 79.0	235	74.5	59.8 85.1	37	72.8	59.5 83.0	35	*	* *	*
ETHNICITY												
Non-Hispanic	70.1	68.0 72.2	1,958	78.1	73.3 82.2	290	77.2	73.2 80.8	316	68.2	61.4 74.2	192
Hispanic	74.3	68.0 79.6	229	*	* *	*	*	* *	*	77.2	65.2 86.0	60
MARITAL STATUS												
Not Married	78.4	73.8 82.3	382	83.8	74.1 90.4	68	83.9	75.0 90.0	62	78.2	65.9 86.9	55
Married	68.1	65.8 70.3	1,808	75.6	70.2 80.4	234	74.0	69.4 78.0	265	67.0	60.4 73.0	197
INCOME												
Less than \$15,000	75.5	71.3 79.3	469	81.7	70.8 89.1	58	82.5	74.1 88.6	75	81.2	70.4 88.7	68
\$15,000 - \$24,999	70.3	65.2 75.0	362	80.5	68.0 88.9	50	78.3	66.1 87.0	42	69.7	52.7 82.6	34
\$25,000 - \$34,999	70.2	64.7 75.1	330	76.2	63.4 85.6	50	77.5	66.4 85.7	48	72.9	58.2 83.8	40
\$35,000 - \$49,999	72.1	66.7 76.9	329	81.5	68.3 90.0	41	66.6	56.1 75.7	57	70.6	53.1 83.6	30
\$50,000+	66.1	62.2 69.9	643	72.7	63.7 80.2	97	74.9	67.4 81.1	101	57.6	46.3 68.2	72
EDUCATION												
K-11th Grade	73.3	66.1 79.5	176	*	* *	*	*	* *	*	79.9	63.8 90.0	36
High School or GED	73.5	69.1 77.5	478	80.8	70.2 88.2	69	86.7	80.0 91.4	85	72.0	59.9 81.6	60
Some College	70.7	67.5 73.8	873	84.5	77.8 89.5	129	77.1	70.1 82.9	113	70.9	61.3 79.0	95
College Graduate+	67.3	63.5 70.9	658	67.6	57.6 76.1	84	69.4	62.1 75.7	117	59.1	46.7 70.5	59

*Figure not reliable by PRATS standards (n < 30)

Immunization Information from PNC Provider

Idaho mothers who reported receiving immunization information from PNC providers, 2010

	Public Health District															
	District 4			District 5			District 6			District 7						
	%	95% CI		n	%	95% CI		n	%	95% CI		n	%	95% CI		n
TOTAL	72.2	67.3	76.7	330	71.6	66.5	76.2	308	66.6	61.6	71.2	327	64.8	59.8	69.5	344
AGE																
18-19	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
20-24	71.6	59.7	81.0	62	76.1	66.0	83.9	80	77.3	68.3	84.3	89	68.7	59.8	76.4	108
25-29	70.1	61.7	77.4	126	72.4	63.9	79.5	111	61.7	53.3	69.4	122	67.2	59.5	74.1	141
30-34	73.8	64.1	81.6	90	73.4	61.7	82.6	65	56.7	46.0	66.8	75	53.4	41.9	64.6	66
35+	73.4	59.2	84.0	45	67.3	52.1	79.7	39	79.3	63.6	89.4	31	*	*	*	*
ETHNICITY																
Non-Hispanic	73.4	68.2	77.9	298	70.0	64.1	75.4	238	65.6	60.3	70.5	298	63.8	58.6	68.6	326
Hispanic	62.0	44.7	76.7	32	76.3	65.3	84.6	70	*	*	*	*	*	*	*	*
MARITAL STATUS																
Not Married	77.4	64.7	86.4	53	74.6	62.9	83.6	59	77.6	65.0	86.6	49	77.8	62.4	88.1	36
Married	70.8	65.3	75.7	277	70.3	64.7	75.3	249	63.3	57.9	68.4	278	62.3	57.1	67.3	308
INCOME																
Less than \$15,000	77.9	64.6	87.2	51	72.5	60.4	81.9	60	73.9	63.9	82.0	78	64.9	54.2	74.2	79
\$15,000 - \$24,999	65.4	52.0	76.8	51	78.8	67.6	86.9	70	59.1	47.0	70.2	60	68.6	55.9	79.0	55
\$25,000 - \$34,999	67.1	52.2	79.2	43	69.9	56.7	80.5	50	74.7	62.1	84.2	50	62.8	49.2	74.6	49
\$35,000 - \$49,999	77.3	63.1	87.2	45	77.3	64.9	86.3	55	58.7	45.2	71.0	46	67.3	54.6	77.8	55
\$50,000+	72.1	64.1	78.9	132	62.0	50.5	72.4	65	64.5	54.4	73.5	81	58.6	48.9	67.6	95
EDUCATION																
K-11th Grade	*	*	*	*	70.1	56.5	80.8	53	*	*	*	*	*	*	*	*
High School or GED	74.8	62.6	84.1	60	77.9	65.7	86.6	58	68.7	59.0	76.9	91	63.7	50.7	75.0	55
Some College	67.5	59.1	74.9	125	71.1	63.3	77.9	135	65.4	57.3	72.7	127	67.9	60.4	74.5	149
College Graduate+	76.7	68.7	83.1	123	67.0	55.3	77.0	62	64.2	54.5	72.9	88	59.5	51.1	67.4	125

*Figure not reliable by PRATS standards (n < 30)

Baby Sleep Position

Idaho mothers who lay their baby on his or her back to sleep, 2010

	Statewide			Public Health District								
				District 1			District 2			District 3		
	%	95% CI	n	%	95% CI	n	%	95% CI	n	%	95% CI	n
TOTAL	80.8	79.0 82.4	2,333	80.4	75.9 84.3	317	83.2	79.6 86.3	340	74.2	68.6 79.1	266
AGE												
18-19	72.1	60.2 81.5	77	*	* *	*	*	* *	*	*	* *	*
20-24	80.4	76.9 83.6	583	79.6	69.4 87.0	73	78.5	69.8 85.3	77	70.5	59.4 79.6	76
25-29	83.5	80.7 85.9	865	84.0	76.0 89.7	103	86.8	81.4 90.7	142	82.1	72.2 89.0	81
30-34	81.8	78.1 85.0	545	79.8	70.2 86.8	84	84.7	76.5 90.4	71	79.6	67.7 87.9	61
35+	73.8	67.6 79.2	263	73.3	58.9 84.0	41	76.2	63.4 85.6	38	63.0	46.8 76.7	38
ETHNICITY												
Non-Hispanic	83.4	81.6 84.9	2,088	80.4	75.7 84.3	303	82.6	78.9 85.7	329	78.8	72.7 83.8	204
Hispanic	63.1	56.5 69.2	242	*	* *	*	*	* *	*	60.3	47.8 71.7	62
MARITAL STATUS												
Not Married	76.1	71.6 80.1	401	78.3	68.0 85.9	69	82.8	74.0 89.1	64	63.8	51.3 74.7	58
Married	82.3	80.5 84.0	1,932	81.5	76.6 85.5	248	83.3	79.4 86.6	276	78.9	73.0 83.7	208
INCOME												
Less than \$15,000	77.9	73.7 81.5	484	85.9	75.0 92.5	59	80.9	72.2 87.3	74	72.5	61.1 81.6	70
\$15,000 - \$24,999	81.4	76.9 85.2	383	78.9	66.3 87.6	50	87.2	76.8 93.3	43	77.3	61.2 88.1	37
\$25,000 - \$34,999	76.9	72.1 81.1	355	67.4	54.6 78.0	56	78.9	68.3 86.7	53	71.0	55.3 82.9	40
\$35,000 - \$49,999	84.8	80.5 88.2	349	82.3	69.6 90.4	43	82.9	73.5 89.5	59	86.9	70.7 94.8	30
\$50,000+	84.3	81.2 87.0	698	85.3	77.6 90.6	103	86.4	80.0 91.0	105	76.3	65.8 84.3	80
EDUCATION												
K-11th Grade	60.2	52.5 67.4	173	*	* *	*	*	* *	*	58.6	42.3 73.3	35
High School or GED	81.5	77.7 84.8	511	80.2	70.5 87.3	72	86.0	78.9 90.9	90	76.8	64.8 85.6	65
Some College	82.6	79.9 85.0	931	81.6	74.3 87.2	134	81.4	74.6 86.6	118	78.2	68.9 85.4	101
College Graduate+	84.4	81.4 87.0	712	83.1	74.6 89.2	91	83.9	77.9 88.5	122	76.3	64.7 84.9	62

*Figure not reliable by PRATS standards (n < 30)

Baby Sleep Position

Idaho mothers who lay their baby on his or her back to sleep, 2010

	Public Health District															
	District 4				District 5				District 6				District 7			
	%	95% CI		n	%	95% CI		n	%	95% CI		n	%	95% CI		n
TOTAL	84.9	80.9	88.2	362	76.9	72.2	81.1	330	83.3	79.1	86.8	348	81.7	77.5	85.2	370
AGE																
18-19	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
20-24	89.0	79.7	94.3	68	78.6	69.0	85.9	82	81.8	72.5	88.5	91	82.6	74.9	88.4	116
25-29	85.8	79.2	90.6	136	76.6	68.5	83.2	121	86.4	79.4	91.3	132	82.8	76.2	87.8	150
30-34	86.7	78.6	92.0	101	79.7	68.9	87.4	72	78.3	68.8	85.5	84	79.4	68.7	87.1	72
35+	75.0	61.4	85.0	50	72.6	57.7	83.8	43	85.4	70.6	93.4	32	*	*	*	*
ETHNICITY																
Non-Hispanic	87.6	83.8	90.7	328	82.0	77.1	86.1	257	84.8	80.6	88.3	317	82.6	78.4	86.1	350
Hispanic	59.8	43.0	74.6	34	61.3	49.9	71.6	73	68.9	51.3	82.4	31	*	*	*	*
MARITAL STATUS																
Not Married	86.2	75.2	92.8	58	76.6	65.5	84.9	64	74.0	61.3	83.7	50	73.7	58.5	84.8	38
Married	84.5	80.2	88.1	304	77.1	72.0	81.5	266	85.9	81.8	89.2	298	83.1	79.0	86.6	332
INCOME																
Less than \$15,000	88.3	77.6	94.3	57	70.2	58.1	80.0	63	81.4	71.5	88.4	77	71.6	61.1	80.2	84
\$15,000 - \$24,999	75.2	62.5	84.6	55	82.5	72.4	89.5	72	84.0	73.4	90.9	68	90.4	80.7	95.5	58
\$25,000 - \$34,999	83.5	70.7	91.4	46	77.2	65.0	86.1	54	79.5	67.5	87.9	55	78.0	65.7	86.8	51
\$35,000 - \$49,999	90.9	78.5	96.5	46	79.6	68.1	87.6	57	86.3	75.0	93.0	51	81.0	70.0	88.6	63
\$50,000+	86.7	80.3	91.2	147	80.7	71.1	87.7	77	86.3	77.7	92.0	85	86.6	78.7	91.9	101
EDUCATION																
K-11th Grade	*	*	*	*	52.5	39.4	65.3	53	*	*	*	*	*	*	*	*
High School or GED	84.5	74.4	91.1	66	74.1	62.7	83.0	64	86.1	77.2	91.8	94	83.8	72.7	90.9	60
Some College	85.1	78.1	90.1	137	84.0	77.4	88.9	144	82.4	75.5	87.6	137	82.8	76.6	87.7	160
College Graduate+	86.6	80.1	91.1	138	88.7	79.7	94.0	69	87.7	80.2	92.6	96	83.0	75.9	88.3	134

*Figure not reliable by PRATS standards (n < 30)

Day Care

Idaho mothers who took their baby to day care¹, 2010

	Statewide			Public Health District								
				District 1			District 2			District 3		
	%	95% CI	n	%	95% CI	n	%	95% CI	n	%	95% CI	n
TOTAL	36.4	33.5 39.4	1,133	36.2	29.7 43.4	169	53.6	47.5 59.6	180	33.8	26.3 42.3	127
AGE												
18-19	20.7	11.4 34.5	43	*	* *	*	*	* *	*	*	* *	*
20-24	23.3	18.6 28.8	300	19.5	10.1 34.4	40	39.8	27.5 53.5	37	25.5	14.8 40.2	43
25-29	40.4	35.4 45.5	413	40.3	29.2 52.4	61	53.6	44.0 63.0	71	32.7	20.7 47.6	41
30-34	46.3	39.8 52.8	250	47.2	33.5 61.4	41	67.4	52.9 79.2	35	*	* *	*
35+	44.7	35.6 54.1	127	*	* *	*	*	* *	*	*	* *	*
ETHNICITY												
Non-Hispanic	37.2	34.1 40.3	1,024	36.6	29.9 43.8	165	54.3	48.1 60.3	178	34.9	26.5 44.4	102
Hispanic	31.2	22.9 40.9	108	*	* *	*	*	* *	*	*	* *	*
MARITAL STATUS												
Not Married	34.3	28.5 40.5	254	34.0	22.8 47.4	47	42.1	30.1 55.1	38	*	* *	*
Married	37.4	34.2 40.8	879	37.7	30.2 45.9	122	58.5	51.7 64.9	142	36.7	28.1 46.3	98
INCOME												
Less than \$15,000	24.1	18.9 30.3	236	30.1	17.3 47.0	33	35.7	23.3 50.3	33	25.5	14.0 42.0	35
\$15,000 - \$24,999	31.1	24.3 38.8	177	*	* *	*	*	* *	*	*	* *	*
\$25,000 - \$34,999	31.4	24.1 39.8	146	*	* *	*	*	* *	*	*	* *	*
\$35,000 - \$49,999	40.5	32.9 48.5	173	*	* *	*	63.5	48.6 76.1	32	*	* *	*
\$50,000+	50.2	44.8 55.7	369	42.9	31.9 54.7	60	70.2	60.8 78.1	72	48.9	34.4 63.5	41
EDUCATION												
K-11th Grade	27.6	17.6 40.6	59	*	* *	*	*	* *	*	*	* *	*
High School or GED	27.5	22.0 33.8	245	20.7	10.0 38.1	30	39.8	28.6 52.1	45	24.9	13.0 42.3	31
Some College	32.9	28.5 37.7	462	35.7	26.5 46.0	84	49.0	39.0 59.2	64	34.2	22.6 48.1	49
College Graduate+	50.0	44.6 55.5	367	47.0	34.7 59.7	51	68.7	58.8 77.1	69	43.0	28.5 58.8	36

*Figure not reliable by PRATS standards (n < 30)

1. Among mothers who were working or attending school.

Day Care

Idaho mothers who took their baby to day care¹, 2010

	Public Health District															
	District 4				District 5				District 6				District 7			
	%	95% CI		n	%	95% CI		n	%	95% CI		n	%	95% CI		n
TOTAL	39.6	33.2	46.4	200	39.3	32.3	46.7	162	31.4	25.0	38.6	164	27.7	20.8	35.9	131
AGE																
18-19	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
20-24	27.2	15.5	43.2	38	25.6	15.8	38.6	47	20.6	11.7	33.6	49	14.9	7.0	28.7	46
25-29	48.1	36.7	59.7	68	38.0	26.4	51.2	54	37.0	26.5	48.9	64	32.3	21.5	45.5	54
30-34	40.6	29.1	53.3	59	63.2	46.3	77.4	33	*	*	*	*	*	*	*	*
35+	40.9	25.7	58.2	31	*	*	*	*	*	*	*	*	*	*	*	*
ETHNICITY																
Non-Hispanic	39.7	33.0	46.9	181	40.9	33.1	49.2	132	32.4	25.5	40.1	144	28.0	20.9	36.5	122
Hispanic	*	*	*	*	32.6	18.6	50.5	30	*	*	*	*	*	*	*	*
MARITAL STATUS																
Not Married	40.4	27.9	54.4	47	25.0	14.1	40.4	36	33.3	20.8	48.8	36	*	*	*	*
Married	39.2	32.0	46.9	153	46.8	38.8	55.0	126	30.5	23.6	38.3	128	24.6	17.7	33.0	110
INCOME																
Less than \$15,000	18.9	8.4	37.2	30	*	*	*	*	26.0	15.0	41.0	40	24.6	13.2	41.1	37
\$15,000 - \$24,999	*	*	*	*	37.4	24.1	53.0	39	31.2	17.6	49.0	30	*	*	*	*
\$25,000 - \$34,999	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
\$35,000 - \$49,999	40.5	25.1	58.1	31	*	*	*	*	*	*	*	*	*	*	*	*
\$50,000+	50.6	40.2	60.9	83	61.0	46.4	73.8	41	42.1	28.6	56.9	39	43.5	28.4	59.8	33
EDUCATION																
K-11th Grade	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
High School or GED	25.0	13.7	41.2	36	33.3	19.2	51.4	30	25.5	15.5	38.8	51	*	*	*	*
Some College	42.6	32.0	54.0	74	30.2	21.3	41.0	75	27.4	17.9	39.4	58	11.6	5.4	23.0	58
College Graduate+	47.9	37.4	58.5	79	69.0	54.4	80.6	43	44.2	31.2	58.1	43	46.5	33.2	60.3	46

*Figure not reliable by PRATS standards (n < 30)

1. Among mothers who were working or attending school.



PRATS

APPENDICES

Appendix A: Maternal and Infant Health Trends

The tables on the following pages present maternal and infant health data for Idaho and each health district from 2004 to 2010. The variable names and descriptions are provided below.

Variable	Description
Unintended Pregnancy	Mothers who reported their pregnancy was unintended
No Birth Control Use	Mothers not using birth control when they became pregnant (among those not trying to get pregnant)
No Health Insurance	Mothers who did not have health insurance prior to pregnancy (excluding Medicaid)
Medicaid Utilization	Mothers who reported Medicaid as a payment source for prenatal care and/or delivery
Overweight/Obesity	Mothers who were overweight or obese prior to pregnancy (BMI greater than or equal to 25)
First Trimester Prenatal Care	Mothers who began prenatal care during the first trimester
WIC	Mothers who participated in the WIC Program during pregnancy
Multivitamin Use	Mothers who regularly took a multivitamin during the third trimester
HIV Testing	Mothers who received an HIV test during pregnancy
No Dental Care	Mothers who did not receive routine dental care during pregnancy
Exercise	Mothers who exercised during the third trimester
Stressful Life Events	Mothers who experienced three or more stressful life events in the 12 months prior to delivery
Physical Abuse	Mothers who experienced physical abuse during pregnancy
Smoking	Mothers who smoked during the third trimester
Alcohol Use	Mothers who drank alcohol during the third trimester
Breastfeeding Initiation	Mothers who ever breastfed
Breastfeeding Duration (6 mos+)	Mothers who breastfed for at least six months
Postpartum Depression	Mothers who reported being moderately or severely depressed during the three months after pregnancy
No Well-Baby Care	Mothers who never took their baby for routine well-baby care
Hearing Screening	Mothers who reported their baby's hearing was tested
Immunization Information from PNC Provider	Mothers who reported receiving information about their baby's immunizations from a prenatal care provider
Baby Sleep Position (On Back)	Mothers who lay their baby on his or her back to sleep
Day Care	Mothers who took their baby to day care (among those working or attending school)

Appendix A: Maternal and Infant Health Trends

By Health District, PRATS 2004-2010

		Year								
		2004			2005			2006		
Idaho		%	95% CI	n	%	95% CI	n	%	95% CI	n
	Unintended Pregnancy	34.6	32.7 36.6	2,667	34.8	32.8 36.8	2,609	34.4	32.4 36.5	2,327
	No Birth Control Use	58.1	54.9 61.1	1,102	60.1	56.9 63.2	1,080	59.7	56.4 63.0	970
	No Health Insurance	39.7	37.7 41.6	2,696	38.0	36.0 40.0	2,634	38.7	36.7 40.8	2,355
	Medicaid Utilization	44.9	42.9 46.8	2,651	41.9	40.0 43.9	2,595	39.0	37.0 41.1	2,326
	Overweight/Obesity	40.7	38.8 42.8	2,590	39.3	37.3 41.4	2,513	37.5	35.4 39.6	2,261
	First Trimester Prenatal Care	88.4	87.0 89.7	2,614	86.9	85.4 88.3	2,565	86.1	84.5 87.5	2,298
	WIC	39.6	37.7 41.6	2,695	38.3	36.3 40.3	2,620	36.9	34.9 39.0	2,341
	Multivitamin Use	81.1	79.5 82.6	2,686	82.7	81.1 84.2	2,617	80.2	78.5 81.8	2,336
	HIV Testing	54.7	52.7 56.7	2,680	53.3	51.2 55.3	2,609	52.6	50.5 54.8	2,326
	No Dental Care	60.8	58.8 62.7	2,675	56.4	54.4 58.4	2,621	56.6	54.4 58.7	2,338
	Exercise During Pregnancy	*	* * *	*	*	* * *	*	*	* * *	*
	Stressful Life Events	*	* * *	*	*	* * *	*	*	* * *	*
	Physical Abuse	4.4	3.6 5.3	2,682	4.7	3.9 5.7	2,627	4.4	3.5 5.4	2,352
	Smoking	9.3	8.2 10.6	2,679	8.4	7.2 9.7	2,622	8.4	7.2 9.7	2,341
	Alcohol Use	3.6	2.9 4.4	2,668	2.8	2.2 3.6	2,625	3.4	2.7 4.4	2,347
	Breastfeeding Initiation	89.0	87.6 90.2	2,690	88.5	87.1 89.7	2,629	91.6	90.3 92.7	2,348
	Breastfeeding Duration (6 mos +)	52.3	50.1 54.4	2,379	50.5	48.3 52.7	2,343	54.0	51.8 56.2	2,206
	Postpartum Depression	20.9	19.3 22.6	2,686	17.7	16.2 19.4	2,624	18.1	16.5 19.9	2,340
	No Well-Baby Care	3.8	3.1 4.7	2,481	3.3	2.6 4.2	2,463	3.0	2.3 3.9	2,127
	Hearing Screening	91.5	90.4 92.5	2,688	92.3	91.1 93.3	2,627	92.6	91.4 93.7	2,349
	Immun. Info By PNC Provider	46.6	44.7 48.6	2,699	47.2	45.2 49.2	2,637	45.0	42.9 47.2	2,357
	Baby Sleeps on Back	73.3	71.5 75.0	2,634	74.9	73.1 76.6	2,573	75.6	73.7 77.4	2,301
Day Care	*	* * *	*	*	* * *	*	*	* * *	*	
District 1		%	95% CI	n	%	95% CI	n	%	95% CI	n
	Unintended Pregnancy	42.9	38.2 47.7	358	40.5	35.6 45.6	327	39.3	34.1 44.6	295
	No Birth Control Use	62.1	55.5 68.2	182	58.4	51.1 65.3	155	61.8	54.1 68.9	142
	No Health Insurance	46.4	41.7 51.2	365	45.7	40.8 50.8	331	46.6	41.4 52.0	300
	Medicaid Utilization	46.9	42.2 51.8	355	47.3	42.2 52.4	322	39.2	34.1 44.5	296
	Overweight/Obesity	40.4	35.8 45.1	354	43.0	38.0 48.0	322	38.8	33.8 44.2	293
	First Trimester Prenatal Care	86.8	83.0 89.8	347	88.2	84.4 91.1	319	87.1	82.9 90.3	291
	WIC	43.3	38.6 48.0	365	41.2	36.4 46.3	329	43.4	38.2 48.7	300
	Multivitamin Use	79.5	75.3 83.2	362	82.7	78.6 86.3	329	82.5	78.1 86.2	298
	HIV Testing	57.4	52.7 62.0	362	59.5	54.5 64.3	327	56.7	51.4 61.9	300
	No Dental Care	62.0	57.3 66.4	361	63.0	58.0 67.7	327	58.1	52.8 63.2	300
	Exercise During Pregnancy	*	* * *	*	*	* * *	*	*	* * *	*
	Stressful Life Events	*	* * *	*	*	* * *	*	*	* * *	*
	Physical Abuse	2.9	1.7 5.1	363	5.8	3.7 9.0	331	5.0	3.1 8.0	300
	Smoking	11.2	8.4 14.8	364	11.1	8.1 15.0	328	12.5	9.3 16.6	299
	Alcohol Use	4.5	3.0 6.9	362	3.6	2.2 5.9	331	4.7	2.9 7.5	300
	Breastfeeding Initiation	89.9	86.5 92.5	365	90.4	86.8 93.1	329	94.2	91.2 96.3	299
	Breastfeeding Duration (6 mos +)	54.3	49.3 59.3	323	52.2	46.9 57.5	292	57.4	51.9 62.8	278
	Postpartum Depression	19.1	15.6 23.2	362	15.9	12.5 19.9	328	18.1	14.3 22.7	297
	No Well-Baby Care	2.8	1.6 4.9	334	2.0	0.9 4.0	304	1.9	0.8 4.2	263
	Hearing Screening	90.7	87.7 93.1	365	92.2	89.3 94.4	331	89.9	86.2 92.7	300
	Immun. Info By PNC Provider	49.6	44.9 54.3	365	55.0	50.0 59.9	331	54.0	48.7 59.2	300
	Baby Sleeps on Back	72.8	68.3 76.8	354	77.7	73.2 81.6	325	76.8	72.0 81.0	292
Day Care	*	* * *	*	*	* * *	*	*	* * *	*	

*Indicates data are not available for that year

Note: Confidence Intervals (95% CI) may differ from previous publications due to revisions in standard error calculation.

Appendix A: Maternal and Infant Health Trends

By Health District, PRATS 2004-2010

	Year											
	2007			2008			2009			2010		
Idaho	%	95% CI	n									
Unintended Pregnancy	36.3	34.1 38.5	2,294	34.7	32.6 36.8	2,316	35.2	33.2 37.3	2,396	33.2	31.2 35.3	2,357
No Birth Control Use	58.5	54.9 62.0	905	59.5	56.1 62.9	927	55.3	52.0 58.5	973	54.9	51.5 58.3	911
No Health Insurance	37.3	35.2 39.5	2,313	35.6	33.5 37.7	2,341	36.0	34.1 38.1	2,419	37.4	35.4 39.5	2,374
Medicaid Utilization	38.4	36.2 40.5	2,265	38.6	36.5 40.7	2,283	43.3	41.2 45.4	2,386	43.9	41.8 46.0	2,363
Overweight/Obesity	41.2	39.0 43.5	2,229	40.6	38.5 42.8	2,255	40.6	38.5 42.7	2,340	44.4	42.3 46.6	2,313
First Trimester Prenatal Care	86.5	84.9 87.9	2,247	88.9	87.4 90.2	2,254	89.4	88.0 90.7	2,362	90.9	89.5 92.0	2,352
WIC	34.9	32.8 37.0	2,302	36.4	34.3 38.5	2,303	40.9	38.8 42.9	2,411	39.9	37.8 42.0	2,366
Multivitamin Use	79.8	78.0 81.5	2,299	81.5	79.8 83.2	2,307	83.6	82.0 85.1	2,411	83.1	81.5 84.7	2,367
HIV Testing	48.6	46.4 50.8	2,288	45.2	43.1 47.4	2,268	48.3	46.2 50.4	2,392	45.5	43.4 47.6	2,332
No Dental Care	54.5	52.3 56.7	2,303	49.6	47.4 51.7	2,300	46.1	44.0 48.1	2,415	48.9	46.8 51.0	2,367
Exercise During Pregnancy	*	* *	*	57.9	55.8 60.1	2,311	57.8	55.7 59.9	2,405	57.3	55.2 59.4	2,353
Stressful Life Events	*	* *	*	25.5	23.6 27.5	2,332	28.3	26.4 30.2	2,418	27.4	25.5 29.4	2,370
Physical Abuse	4.1	3.2 5.2	2,313	4.3	3.5 5.4	2,328	5.2	4.3 6.2	2,401	4.2	3.4 5.2	2,364
Smoking	8.0	6.8 9.5	2,304	8.1	6.9 9.4	2,326	8.5	7.4 9.7	2,411	8.0	6.9 9.3	2,362
Alcohol Use	3.3	2.6 4.2	2,311	2.9	2.2 3.7	2,323	4.1	3.3 5.1	2,405	3.8	3.0 4.8	2,359
Breastfeeding Initiation	89.9	88.5 91.2	2,311	91.5	90.2 92.6	2,331	90.6	89.4 91.8	2,407	91.5	90.2 92.6	2,370
Breastfeeding Duration (6 mos +)	50.5	48.2 52.8	2,123	55.2	52.9 57.4	2,096	55.4	53.2 57.5	2,215	53.0	50.8 55.2	2,208
Postpartum Depression	17.4	15.7 19.2	2,306	16.4	14.9 18.0	2,327	18.0	16.4 19.7	2,404	16.8	15.3 18.5	2,361
No Well-Baby Care	3.1	2.4 4.1	2,060	2.9	2.2 3.7	2,042	1.9	1.4 2.6	2,276	2.1	1.6 2.8	2,262
Hearing Screening	94.2	93.1 95.1	2,312	93.7	92.6 94.6	2,328	94.8	93.8 95.6	2,402	95.1	94.1 95.9	2,366
Immun. Info By PNC Provider	45.3	43.1 47.5	2,312	66.9	64.7 69.0	2,133	71.0	69.1 73.0	2,253	70.7	68.6 72.6	2,190
Baby Sleeps on Back	77.7	75.8 79.5	2,261	78.5	76.6 80.3	2,308	77.0	75.2 78.7	2,373	80.8	79.0 82.4	2,333
Day Care	*	* *	*	40.5	37.4 43.8	1,057	42.6	39.5 45.6	1,128	36.4	33.5 39.4	1,133
District 1	%	95% CI	n									
Unintended Pregnancy	38.0	32.3 43.9	277	42.5	37.3 47.9	301	43.7	38.3 49.1	295	34.2	29.3 39.4	321
No Birth Control Use	66.6	57.5 74.7	111	57.3	49.3 64.9	136	59.6	51.6 67.1	137	62.1	54.0 69.6	134
No Health Insurance	46.2	40.4 52.0	279	43.4	38.2 48.7	305	45.1	39.8 50.6	296	40.3	35.2 45.6	322
Medicaid Utilization	40.8	35.0 46.8	273	42.5	37.3 48.0	297	48.9	43.5 54.3	295	43.9	38.7 49.2	319
Overweight/Obesity	40.6	35.1 46.4	277	39.1	34.0 44.4	297	42.3	37.1 47.8	290	47.1	41.9 52.3	320
First Trimester Prenatal Care	84.6	79.7 88.5	271	86.5	82.4 89.8	293	88.5	84.4 91.6	290	89.6	85.7 92.5	319
WIC	34.8	29.4 40.7	278	38.2	33.1 43.6	299	45.6	40.2 51.0	292	42.9	37.8 48.2	321
Multivitamin Use	79.8	74.7 84.1	278	83.7	79.3 87.2	300	91.2	87.6 93.9	293	83.8	79.6 87.3	321
HIV Testing	53.8	47.9 59.5	277	53.7	48.3 59.0	294	58.2	52.7 63.4	287	51.2	45.9 56.4	316
No Dental Care	60.2	54.4 65.6	278	57.4	52.1 62.6	298	49.2	43.8 54.6	294	49.5	44.3 54.7	321
Exercise During Pregnancy	*	* *	*	63.9	58.6 68.8	302	59.7	54.2 64.8	294	59.3	54.1 64.3	318
Stressful Life Events	*	* *	*	33.5	28.6 38.8	305	39.4	34.2 44.9	296	27.9	23.4 33.0	321
Physical Abuse	2.5	1.1 5.4	279	5.7	3.6 8.9	303	5.8	3.7 9.0	293	6.9	4.6 10.4	320
Smoking	12.1	8.6 16.8	278	11.9	8.7 16.1	304	14.9	11.3 19.3	295	10.7	7.7 14.7	317
Alcohol Use	4.6	2.7 7.7	279	4.0	2.4 6.7	303	6.1	4.0 9.1	295	2.9	1.6 5.1	319
Breastfeeding Initiation	91.5	88.0 94.1	278	90.3	86.6 93.1	304	92.2	88.8 94.7	294	91.0	87.4 93.6	321
Breastfeeding Duration (6 mos +)	54.3	48.2 60.3	256	58.7	53.1 64.1	274	55.9	50.3 61.4	279	56.3	50.8 61.7	293
Postpartum Depression	14.9	11.0 19.8	278	18.2	14.4 22.7	304	19.4	15.4 24.2	293	12.7	9.5 16.8	320
No Well-Baby Care	4.2	2.4 7.3	244	4.3	2.5 7.3	256	1.0	0.3 2.7	278	2.2	1.2 4.2	308
Hearing Screening	92.1	88.6 94.6	278	89.5	85.8 92.3	303	93.1	89.9 95.4	293	93.7	90.8 95.8	319
Immun. Info By PNC Provider	51.7	45.8 57.4	279	74.9	69.8 79.4	276	75.0	69.9 79.5	280	78.4	73.7 82.4	302
Baby Sleeps on Back	81.7	76.7 85.8	276	79.3	74.7 83.3	302	79.4	74.6 83.4	292	80.4	75.9 84.3	317
Day Care	*	* *	*	39.3	31.9 47.1	140	40.5	33.3 48.1	151	36.2	29.7 43.4	169

*Indicates data are not available for that year

Note: Confidence Intervals (95% CI) may differ from previous publications due to revisions in standard error calculation.

Appendix A: Maternal and Infant Health Trends

By Health District, PRATS 2004-2010

		Year									
		2004			2005			2006			
		%	95% CI	n	%	95% CI	n	%	95% CI	n	
District 2											
	Unintended Pregnancy	34.9	31.2 38.9	383	33.1	29.5 37.0	394	35.3	30.8 40.2	314	
	No Birth Control Use	49.1	43.0 55.2	162	59.5	53.3 65.5	155	56.5	48.9 63.8	125	
	No Health Insurance	34.1	30.4 38.0	388	32.1	28.5 36.0	396	35.7	31.2 40.4	318	
	Medicaid Utilization	44.3	40.3 48.3	385	38.1	34.3 42.1	390	43.7	39.0 48.5	315	
	Overweight/Obesity	38.9	35.1 42.8	381	42.5	38.6 46.4	387	43.3	38.7 48.1	313	
	First Trimester Prenatal Care	83.8	80.6 86.6	377	87.4	84.4 89.8	386	83.7	79.7 87.1	310	
	WIC	44.1	40.2 48.1	387	33.8	30.1 37.7	396	38.6	34.0 43.4	317	
	Multivitamin Use	87.1	84.1 89.6	386	87.5	84.7 89.9	395	84.0	80.1 87.2	317	
	HIV Testing	45.3	41.4 49.3	386	48.5	44.6 52.4	395	50.7	46.0 55.4	316	
	No Dental Care	60.2	56.3 64.0	387	55.2	51.3 59.1	396	52.2	47.5 56.9	318	
	Exercise During Pregnancy	*	* * *	*	*	* * *	*	*	* * *	*	
	Stressful Life Events	*	* * *	*	*	* * *	*	*	* * *	*	
	Physical Abuse	4.2	2.8 6.2	385	2.2	1.3 3.8	396	4.9	3.0 7.7	317	
	Smoking	12.0	9.5 15.1	386	9.1	6.9 11.8	395	13.0	9.8 16.9	318	
	Alcohol Use	4.2	2.8 6.1	384	5.6	4.0 7.7	395	5.4	3.8 7.7	318	
	Breastfeeding Initiation	89.7	87.0 92.0	388	92.5	90.2 94.4	395	91.9	88.8 94.3	317	
	Breastfeeding Duration (6 mos +)	56.1	51.9 60.2	350	53.3	49.1 57.4	351	54.6	49.7 59.4	300	
	Postpartum Depression	18.5	15.5 21.8	385	12.0	9.7 14.8	394	14.5	11.4 18.2	317	
	No Well-Baby Care	2.3	1.3 3.9	347	3.7	2.5 5.6	370	3.4	1.9 5.8	296	
	Hearing Screening	80.7	77.4 83.7	386	88.1	85.2 90.4	395	85.9	82.3 88.8	317	
Immun. Info By PNC Provider	51.5	47.6 55.5	388	58.2	54.3 62.0	396	52.6	47.9 57.3	318		
Baby Sleeps on Back	70.5	66.7 74.1	379	77.9	74.4 81.0	384	73.3	68.9 77.3	309		
Day Care	*	* * *	*	*	* * *	*	*	* * *	*		
District 3											
	Unintended Pregnancy	35.6	30.8 40.7	345	35.0	30.2 40.1	357	38.6	33.2 44.2	297	
	No Birth Control Use	58.2	49.9 66.1	132	58.4	50.6 65.9	150	61.2	52.9 68.9	134	
	No Health Insurance	44.6	39.5 49.8	346	42.8	37.8 47.9	362	46.5	41.1 52.1	301	
	Medicaid Utilization	46.0	40.9 51.3	340	40.2	35.2 45.5	354	39.0	33.6 44.6	294	
	Overweight/Obesity	51.4	46.1 56.7	327	48.3	43.2 53.5	339	40.8	35.4 46.5	279	
	First Trimester Prenatal Care	88.0	84.0 91.0	339	85.9	81.8 89.3	350	82.8	78.0 86.7	292	
	WIC	41.9	36.9 47.2	347	37.1	32.2 42.3	358	34.2	29.1 39.7	300	
	Multivitamin Use	81.3	76.8 85.0	346	82.2	78.0 85.8	361	76.3	71.2 80.7	300	
	HIV Testing	48.2	43.1 53.4	345	49.4	44.3 54.4	359	52.9	47.4 58.4	298	
	No Dental Care	59.7	54.7 64.6	346	61.9	57.0 66.6	360	64.4	58.9 69.5	298	
	Exercise During Pregnancy	*	* * *	*	*	* * *	*	*	* * *	*	
	Stressful Life Events	*	* * *	*	*	* * *	*	*	* * *	*	
	Physical Abuse	5.0	3.1 8.1	346	4.9	2.9 8.0	360	4.7	2.8 7.8	302	
	Smoking	7.9	5.4 11.5	346	8.9	6.4 12.5	360	8.7	5.9 12.7	301	
	Alcohol Use	1.7	0.8 3.7	346	2.7	1.5 4.8	359	3.0	1.6 5.6	300	
	Breastfeeding Initiation	85.1	80.8 88.6	345	85.5	81.4 88.8	361	91.7	87.9 94.4	299	
	Breastfeeding Duration (6 mos+)	47.6	42.2 53.1	302	51.2	45.9 56.4	327	48.5	42.8 54.2	278	
	Postpartum Depression	18.2	14.5 22.7	346	15.7	12.3 19.9	360	14.8	11.2 19.3	298	
	No Well-Baby Care	5.2	3.2 8.3	316	6.5	4.2 9.9	344	5.8	3.5 9.5	264	
	Hearing Screening	88.6	84.8 91.6	346	87.5	83.5 90.7	361	87.9	83.7 91.2	300	
Immun. Info By PNC Provider	44.3	39.3 49.4	348	43.0	38.1 48.0	363	40.7	35.5 46.3	302		
Baby Sleeps on Back	70.1	65.0 74.7	337	73.3	68.5 77.6	357	68.1	62.7 73.1	292		
Day Care	*	* * *	*	*	* * *	*	*	* * *	*		

*Indicates data are not available for that year

- Figure not reliable by PRATS standards (n<30)

Note: Confidence Intervals (95% CI) may differ from previous publications due to revisions in standard error calculation.

Appendix A: Maternal and Infant Health Trends

By Health District, PRATS 2004-2010

	Year											
	2007			2008			2009			2010		
District 2	%	95% CI	n									
Unintended Pregnancy	34.7	30.4 39.2	329	36.7	32.5 41.2	342	36.2	32.0 40.7	349	31.9	27.8 36.3	340
No Birth Control Use	58.5	51.2 65.3	133	54.8	48.0 61.5	145	59.2	52.3 65.8	143	56.5	49.3 63.4	131
No Health Insurance	33.6	29.4 38.1	335	34.0	29.9 38.3	342	30.1	26.1 34.4	355	34.7	30.5 39.1	344
Medicaid Utilization	38.3	33.8 42.9	329	38.3	34.0 42.8	331	39.7	35.4 44.2	344	43.1	38.8 47.6	343
Overweight/Obesity	37.1	32.8 41.5	331	40.5	36.2 45.0	332	41.3	37.0 45.6	348	45.4	41.1 49.8	341
First Trimester Prenatal Care	87.3	83.8 90.2	328	88.8	85.5 91.3	328	89.4	86.4 91.8	338	87.9	84.6 90.5	341
WIC	35.1	30.8 39.6	333	39.1	34.8 43.5	339	35.8	31.7 40.2	355	42.2	37.8 46.6	345
Multivitamin Use	83.4	79.7 86.6	333	87.1	83.8 89.8	340	86.1	82.8 88.8	354	85.2	81.7 88.1	345
HIV Testing	52.4	47.9 56.9	331	51.5	47.0 56.0	334	44.6	40.4 49.0	353	44.4	40.0 48.8	340
No Dental Care	54.1	49.5 58.5	333	51.3	46.9 55.8	338	51.2	46.9 55.5	354	49.6	45.3 54.0	345
Exercise During Pregnancy	*	* * *	*	61.4	57.0 65.7	339	61.2	56.8 65.4	350	59.7	55.3 64.0	342
Stressful Life Events	*	* * *	*	25.2	21.4 29.3	341	27.1	23.3 31.4	355	24.9	21.1 29.0	344
Physical Abuse	4.5	2.9 7.2	335	6.1	4.2 8.7	340	5.7	3.9 8.2	350	4.4	2.8 6.9	343
Smoking	7.9	5.6 11.0	333	13.0	10.2 16.4	341	11.1	8.5 14.5	354	9.8	7.3 13.0	341
Alcohol Use	4.9	3.3 7.2	336	4.5	3.0 6.7	339	7.6	5.6 10.2	352	5.3	3.7 7.6	342
Breastfeeding Initiation	92.8	90.0 94.9	336	93.5	90.9 95.4	338	92.8	90.1 94.7	351	91.9	89.0 94.1	344
Breastfeeding Duration (6 mos +)	54.2	49.4 58.8	307	54.7	50.0 59.4	300	60.9	56.3 65.3	321	57.6	53.0 62.0	325
Postpartum Depression	13.2	10.4 16.7	335	20.7	17.2 24.6	339	15.3	12.3 18.8	351	17.1	13.9 20.8	342
No Well-Baby Care	1.9	1.0 3.6	294	2.3	1.2 4.2	300	2.7	1.6 4.5	329	1.4	0.7 2.8	327
Hearing Screening	88.4	85.2 91.0	334	89.2	86.1 91.7	338	91.7	88.9 93.8	349	91.8	89.1 93.9	343
Immun. Info By PNC Provider	55.5	50.9 59.9	335	75.7	71.5 79.4	316	79.9	76.1 83.2	334	76.7	72.7 80.2	327
Baby Sleeps on Back	77.5	73.5 81.0	330	80.6	76.7 83.9	334	74.6	70.6 78.2	344	83.2	79.6 86.3	340
Day Care	*	* * *	*	48.4	42.7 54.1	203	53.2	47.3 59.0	195	53.6	47.5 59.6	180
District 3	%	95% CI	n									
Unintended Pregnancy	42.1	36.8 47.6	310	33.8	28.6 39.4	289	41.6	36.5 47.0	314	33.0	27.6 38.9	272
No Birth Control Use	61.6	53.6 69.0	146	62.3	53.7 70.2	123	60.1	52.5 67.2	157	52.9	43.7 62.0	109
No Health Insurance	44.3	39.0 49.8	311	42.3	36.9 47.9	293	49.0	43.7 54.3	316	45.2	39.5 51.2	272
Medicaid Utilization	41.3	35.9 46.8	304	44.4	38.9 50.1	288	50.7	45.4 56.0	312	49.1	43.3 55.0	271
Overweight/Obesity	47.8	42.3 53.3	295	46.4	40.7 52.1	277	44.9	39.5 50.3	301	52.3	46.3 58.3	256
First Trimester Prenatal Care	82.1	77.4 86.0	300	86.0	81.6 89.5	283	88.6	84.7 91.7	308	90.2	85.8 93.3	271
WIC	35.2	30.1 40.6	310	41.4	36.0 47.1	287	47.4	42.1 52.8	315	42.9	37.2 48.9	269
Multivitamin Use	77.3	72.4 81.6	308	79.8	74.8 84.0	285	79.3	74.6 83.3	316	80.8	75.7 85.0	269
HIV Testing	49.3	43.9 54.7	308	42.7	37.1 48.4	281	48.2	42.9 53.5	313	45.9	40.1 51.8	270
No Dental Care	60.5	55.2 65.7	310	52.8	47.2 58.4	285	52.9	47.5 58.1	315	53.1	47.3 58.9	272
Exercise During Pregnancy	*	* * *	*	55.7	50.1 61.2	291	54.3	49.0 59.6	315	52.9	47.1 58.7	269
Stressful Life Events	*	* * *	*	28.8	23.9 34.3	293	35.3	30.4 40.6	315	31.5	26.2 37.3	271
Physical Abuse	5.1	3.1 8.3	311	5.4	3.3 8.8	293	5.4	3.4 8.6	313	4.5	2.6 7.7	273
Smoking	8.4	5.7 12.2	311	10.5	7.4 14.8	292	10.1	7.3 14.0	315	10.2	7.0 14.6	272
Alcohol Use	2.4	1.2 4.5	310	2.5	1.2 5.0	291	1.9	0.9 4.0	313	3.6	1.9 6.6	269
Breastfeeding Initiation	84.9	80.5 88.5	312	90.0	86.0 92.9	294	89.5	85.7 92.5	314	91.4	87.4 94.2	273
Breastfeeding Duration (6 mos +)	41.4	35.9 47.1	281	45.5	39.7 51.4	262	47.3	41.8 52.8	288	49.3	43.3 55.3	255
Postpartum Depression	21.3	17.2 26.1	311	15.5	11.8 20.0	294	15.9	12.4 20.3	313	16.2	12.2 21.2	272
No Well-Baby Care	4.5	2.6 7.6	268	5.2	3.1 8.6	254	2.1	1.0 4.5	293	1.4	0.6 3.6	251
Hearing Screening	92.4	89.1 94.8	312	90.1	86.2 93.0	291	92.5	89.1 94.9	314	91.6	87.6 94.4	273
Immun. Info By PNC Provider	46.4	41.1 51.8	312	61.5	55.6 67.0	264	71.0	65.8 75.8	293	70.5	64.7 75.6	252
Baby Sleeps on Back	75.7	70.7 80.1	303	75.3	70.1 79.9	289	77.8	73.0 81.9	311	74.2	68.6 79.1	266
Day Care	*	* * *	*	44.0	35.8 52.7	124	53.4	45.3 61.4	135	33.8	26.3 42.3	127

*Indicates data are not available for that year

– Figure not reliable by PRATS standards (n<30)

Note: Confidence Intervals (95% CI) may differ from previous publications due to revisions in standard error calculation.

Appendix A: Maternal and Infant Health Trends

By Health District, PRATS 2004-2010

		Year								
		2004			2005			2006		
District 4		%	95% CI	n	%	95% CI	n	%	95% CI	n
	Unintended Pregnancy	33.6	29.1 38.4	400	31.7	27.1 36.8	367	30.7	26.1 35.8	349
	No Birth Control Use	58.4	50.9 65.5	167	60.7	52.1 68.7	128	60.2	51.5 68.3	126
	No Health Insurance	29.5	25.2 34.3	404	23.6	19.3 28.5	369	28.1	23.5 33.2	352
	Medicaid Utilization	32.4	27.9 37.2	402	26.4	21.9 31.4	364	25.4	20.9 30.4	348
	Overweight/Obesity	37.1	32.5 42.0	391	31.4	26.9 36.3	362	35.1	30.3 40.2	346
	First Trimester Prenatal Care	91.1	87.7 93.6	396	90.5	86.9 93.2	363	91.0	87.3 93.6	345
	WIC	27.1	22.9 31.8	405	25.8	21.4 30.7	368	24.9	20.5 30.0	346
	Multivitamin Use	86.3	82.6 89.3	404	86.9	83.0 90.0	368	88.7	85.0 91.6	345
	HIV Testing	57.0	52.2 61.6	405	51.8	46.8 56.9	365	53.4	48.2 58.6	342
	No Dental Care	53.3	48.5 58.0	400	42.9	37.9 48.0	367	45.1	40.0 50.3	347
	Exercise During Pregnancy	*	* *	*	*	* *	*	*	* *	*
	Stressful Life Events	*	* *	*	*	* *	*	*	* *	*
	Physical Abuse	3.0	1.7 5.4	401	3.3	1.8 5.9	367	4.4	2.6 7.3	350
	Smoking	8.4	5.9 11.7	401	7.8	5.2 11.4	368	7.1	4.7 10.6	345
	Alcohol Use	5.9	4.0 8.5	398	3.4	2.0 5.9	368	5.8	3.9 8.7	351
	Breastfeeding Initiation	90.0	86.7 92.5	405	91.6	88.3 94.0	369	93.8	90.7 95.9	352
	Breastfeeding Duration (6 mos +)	53.7	48.6 58.7	359	52.5	47.2 57.8	329	55.0	49.7 60.2	333
	Postpartum Depression	21.1	17.4 25.3	405	15.1	11.7 19.3	369	20.2	16.4 24.7	352
	No Well-Baby Care	2.4	1.2 4.7	374	0.9	0.3 2.7	356	1.9	0.9 4.2	313
	Hearing Screening	96.0	93.7 97.5	404	95.9	93.2 97.5	369	96.2	93.6 97.8	352
Immun. Info By PNC Provider	51.0	46.2 55.8	405	44.7	39.8 49.7	369	46.0	40.9 51.1	352	
Baby Sleeps on Back	77.8	73.5 81.6	395	81.6	77.3 85.2	362	79.5	74.9 83.4	345	
Day Care	*	* *	*	*	* *	*	*	* *	*	
District 5		%	95% CI	n	%	95% CI	n	%	95% CI	n
	Unintended Pregnancy	35.2	30.6 40.2	363	41.4	36.6 46.5	357	38.3	33.4 43.5	339
	No Birth Control Use	61.7	54.3 68.6	162	63.5	56.7 69.8	185	66.3	59.0 72.9	161
	No Health Insurance	52.7	47.8 57.5	367	53.6	48.7 58.5	360	51.2	46.2 56.2	348
	Medicaid Utilization	56.6	51.7 61.4	355	49.8	44.8 54.8	356	47.3	42.3 52.4	345
	Overweight/Obesity	40.5	35.6 45.6	338	42.7	37.6 47.9	328	39.4	34.4 44.6	321
	First Trimester Prenatal Care	84.8	80.6 88.1	348	79.8	75.3 83.6	353	80.4	75.9 84.2	339
	WIC	46.1	41.2 51.1	366	43.6	38.7 48.6	357	44.0	39.0 49.1	344
	Multivitamin Use	77.9	73.4 81.8	367	80.4	76.1 84.1	354	77.4	72.7 81.4	344
	HIV Testing	59.0	54.1 63.7	364	56.3	51.3 61.1	358	58.1	53.1 63.0	343
	No Dental Care	73.5	69.0 77.5	362	66.8	62.1 71.2	360	65.0	60.0 69.6	344
	Exercise During Pregnancy	*	* *	*	*	* *	*	*	* *	*
	Stressful Life Events	*	* *	*	*	* *	*	*	* *	*
	Physical Abuse	6.0	3.9 9.0	364	6.5	4.3 9.7	359	6.5	4.3 9.7	346
	Smoking	13.0	9.9 17.0	361	8.7	6.1 12.3	358	10.2	7.5 13.8	345
	Alcohol Use	4.5	2.8 7.1	358	4.2	2.6 6.8	358	2.8	1.5 4.9	346
	Breastfeeding Initiation	89.6	86.0 92.3	363	85.2	81.2 88.5	359	86.3	82.2 89.5	346
	Breastfeeding Duration (6 mos +)	47.9	42.7 53.2	324	43.9	38.7 49.1	317	46.6	41.5 51.9	321
	Postpartum Depression	22.2	18.4 26.5	362	20.2	16.5 24.6	360	15.5	12.1 19.5	344
	No Well-Baby Care	7.0	4.8 10.2	342	6.7	4.4 10.2	334	5.4	3.3 8.5	315
	Hearing Screening	90.5	87.0 93.2	365	92.4	89.0 94.8	357	96.8	94.2 98.2	346
Immun. Info By PNC Provider	44.6	39.8 49.5	367	48.4	43.5 53.4	361	42.1	37.3 47.2	348	
Baby Sleeps on Back	63.5	58.5 68.1	359	66.8	61.8 71.4	347	69.8	64.9 74.3	338	
Day Care	*	* *	*	*	* *	*	*	* *	*	

*Indicates data are not available for that year

Note: Confidence Intervals (95% CI) may differ from previous publications due to revisions in standard error calculation.

Appendix A: Maternal and Infant Health Trends

By Health District, PRATS 2004-2010

	Year											
	2007			2008			2009			2010		
District 4	%	95% CI	n									
Unintended Pregnancy	36.2	31.0 41.8	343	33.6	28.5 39.1	319	33.5	28.8 38.5	371	34.6	29.9 39.7	363
No Birth Control Use	54.2	45.3 62.9	128	59.2	49.6 68.1	105	53.8	45.4 61.9	139	51.4	43.2 59.5	138
No Health Insurance	27.5	22.8 32.9	345	26.5	21.8 31.7	322	26.5	22.1 31.4	373	29.8	25.3 34.7	367
Medicaid Utilization	25.4	20.6 30.8	339	24.1	19.6 29.4	312	32.3	27.6 37.4	367	32.1	27.5 37.2	363
Overweight/Obesity	39.5	34.4 44.9	339	34.7	29.7 40.1	312	37.3	32.5 42.4	360	40.9	36.0 45.9	362
First Trimester Prenatal Care	89.7	85.8 92.6	336	89.9	85.8 92.9	307	89.5	85.9 92.3	366	92.8	89.7 95.0	361
WIC	24.5	19.8 29.9	340	23.7	19.2 28.9	319	30.3	25.8 35.3	372	29.4	25.0 34.3	364
Multivitamin Use	85.2	81.0 88.7	342	82.2	77.5 86.0	319	88.1	84.3 91.1	371	87.1	83.3 90.1	365
HIV Testing	50.7	45.3 56.0	341	37.3	32.1 42.7	316	50.4	45.3 55.4	370	47.6	42.6 52.6	361
No Dental Care	44.3	39.0 49.8	342	38.5	33.3 43.9	318	34.2	29.6 39.1	373	40.0	35.1 45.0	363
Exercise During Pregnancy	*	* *	*	56.4	50.9 61.7	315	60.8	55.8 65.5	372	58.0	52.9 62.8	364
Stressful Life Events	*	* *	*	23.1	18.7 28.1	321	25.0	20.8 29.8	374	24.9	20.7 29.6	366
Physical Abuse	4.1	2.2 7.5	345	3.5	1.9 6.4	320	3.9	2.3 6.5	371	4.9	3.1 7.8	364
Smoking	8.1	5.3 12.3	340	6.4	4.0 9.9	319	6.2	4.1 9.3	374	8.3	5.8 11.8	366
Alcohol Use	4.6	2.8 7.5	344	3.5	2.0 6.2	320	6.7	4.5 9.7	373	5.9	3.9 8.9	366
Breastfeeding Initiation	92.0	88.3 94.6	344	91.8	88.3 94.3	321	93.6	90.6 95.7	373	92.0	88.8 94.4	365
Breastfeeding Duration (6 mos +)	53.1	47.5 58.6	318	58.2	52.5 63.7	295	58.6	53.4 63.7	349	55.8	50.6 60.9	343
Postpartum Depression	17.1	13.3 21.8	342	12.9	9.7 16.9	318	16.8	13.4 20.8	373	17.5	14.0 21.7	364
No Well-Baby Care	2.0	0.8 4.8	315	1.9	0.8 4.3	279	1.6	0.7 3.4	348	2.1	1.0 4.3	354
Hearing Screening	96.4	93.5 98.0	344	95.8	92.9 97.5	320	96.9	94.7 98.2	371	97.1	94.9 98.4	365
Immun. Info By PNC Provider	40.7	35.5 46.1	344	64.6	59.0 69.8	289	72.2	67.4 76.5	353	72.2	67.3 76.7	330
Baby Sleeps on Back	78.1	73.4 82.2	335	79.3	74.5 83.5	318	78.5	74.0 82.4	365	84.9	80.9 88.2	362
Day Care	*	* *	*	44.6	36.3 53.3	128	42.4	35.8 49.3	199	39.6	33.2 46.4	200
District 5	%	95% CI	n									
Unintended Pregnancy	38.1	33.0 43.5	309	40.0	34.9 45.3	320	37.0	32.1 42.3	315	38.5	33.5 43.9	335
No Birth Control Use	53.9	45.5 62.0	129	58.3	50.5 65.6	149	46.8	39.2 54.6	143	54.1	46.1 62.0	142
No Health Insurance	50.0	44.7 55.3	316	44.6	39.5 49.9	326	45.8	40.7 51.0	322	47.9	42.7 53.1	340
Medicaid Utilization	46.4	41.1 51.8	310	45.5	40.3 50.8	322	48.8	43.6 54.1	314	54.5	49.4 59.6	339
Overweight/Obesity	43.5	38.1 49.1	286	50.5	45.1 55.8	306	44.0	38.8 49.4	301	50.6	45.3 55.9	324
First Trimester Prenatal Care	84.3	79.8 87.9	305	87.1	82.9 90.3	316	84.3	80.1 87.9	312	89.5	85.8 92.4	335
WIC	38.7	33.6 44.0	315	38.4	33.4 43.7	320	40.7	35.7 45.9	321	46.8	41.6 52.1	338
Multivitamin Use	75.1	70.2 79.4	313	84.6	80.4 88.0	321	83.1	78.9 86.7	321	79.9	75.5 83.8	338
HIV Testing	51.4	46.0 56.6	312	58.8	53.6 63.9	315	53.1	47.9 58.3	318	53.0	47.8 58.3	329
No Dental Care	62.5	57.3 67.5	316	56.7	51.4 61.8	320	56.2	51.0 61.3	322	55.7	50.5 60.7	339
Exercise During Pregnancy	*	* *	*	61.6	56.3 66.5	320	56.1	50.9 61.2	321	59.7	54.5 64.7	337
Stressful Life Events	*	* *	*	26.4	21.9 31.3	324	23.8	19.6 28.6	322	32.2	27.4 37.5	338
Physical Abuse	5.8	3.7 9.0	315	4.8	3.0 7.6	323	7.0	4.7 10.3	322	4.7	2.9 7.5	339
Smoking	8.8	6.1 12.6	316	5.8	3.7 8.9	323	8.0	5.5 11.5	318	6.3	4.0 9.8	337
Alcohol Use	4.2	2.5 6.9	316	3.9	2.4 6.5	323	4.8	3.0 7.5	318	4.7	2.8 7.8	335
Breastfeeding Initiation	87.9	83.8 91.1	312	91.2	87.8 93.7	324	85.8	81.7 89.1	321	89.6	86.0 92.4	339
Breastfeeding Duration (6 mos +)	48.0	42.5 53.5	290	53.6	48.1 59.0	294	48.3	42.9 53.7	293	46.2	40.9 51.6	314
Postpartum Depression	17.6	13.8 22.0	311	15.9	12.4 20.1	323	18.4	14.7 22.8	321	19.5	15.7 24.1	336
No Well-Baby Care	3.8	2.1 6.8	289	3.0	1.6 5.6	285	3.7	2.1 6.4	298	2.2	1.1 4.1	319
Hearing Screening	96.3	93.7 97.8	316	97.1	94.4 98.5	324	93.8	90.8 95.9	323	94.1	91.0 96.2	339
Immun. Info By PNC Provider	40.7	35.6 46.0	315	76.0	71.1 80.3	294	70.4	65.2 75.1	295	71.6	66.5 76.2	308
Baby Sleeps on Back	65.5	60.1 70.4	304	72.4	67.4 76.9	318	68.0	63.0 72.7	315	76.9	72.2 81.1	330
Day Care	*	* *	*	44.3	36.8 52.0	149	47.7	40.1 55.4	145	39.3	32.3 46.7	162

*Indicates data are not available for that year

Note: Confidence Intervals (95% CI) may differ from previous publications due to revisions in standard error calculation.

Appendix A: Maternal and Infant Health Trends

By Health District, PRATS 2004-2010

		Year								
		2004			2005			2006		
District 6		%	95% CI	n	%	95% CI	n	%	95% CI	n
	Unintended Pregnancy	35.4	31.1 40.0	399	36.7	32.3 41.4	400	37.3	32.6 42.3	368
	No Birth Control Use	53.8	46.5 60.9	161	58.5	51.1 65.4	166	48.6	40.6 56.6	139
	No Health Insurance	45.2	40.6 49.8	403	44.0	39.4 48.7	403	42.8	38.0 47.8	368
	Medicaid Utilization	54.1	49.5 58.6	398	53.6	49.0 58.2	400	48.7	43.8 53.7	361
	Overweight/Obesity	40.1	35.7 44.8	392	39.5	35.0 44.2	389	36.6	31.9 41.5	359
	First Trimester Prenatal Care	88.3	85.0 91.0	392	87.8	84.2 90.7	390	84.0	79.9 87.5	356
	WIC	50.9	46.3 55.5	402	51.3	46.7 55.9	401	44.7	39.8 49.6	367
	Multivitamin Use	75.4	71.2 79.1	400	83.4	79.7 86.5	402	80.8	76.7 84.3	367
	HIV Testing	61.4	56.9 65.8	395	60.5	55.9 65.0	398	54.6	49.7 59.4	364
	No Dental Care	62.6	58.1 66.9	402	61.7	57.2 66.0	402	56.4	51.5 61.1	366
	Exercise During Pregnancy	*	* *	*	*	* *	*	*	* *	*
	Stressful Life Events	*	* *	*	*	* *	*	*	* *	*
	Physical Abuse	6.7	4.6 9.7	401	5.1	3.2 7.9	401	4.1	2.4 6.8	369
	Smoking	9.8	7.3 13.0	401	10.7	7.8 14.4	402	8.4	5.8 12.1	366
	Alcohol Use	2.9	1.7 5.0	401	1.2	0.4 3.1	402	1.4	0.6 3.6	367
	Breastfeeding Initiation	87.6	84.0 90.5	403	87.8	84.2 90.6	404	88.5	84.7 91.4	368
	Breastfeeding Duration (6 mos +)	54.3	49.3 59.2	345	46.9	42.1 51.8	362	55.2	50.1 60.1	353
	Postpartum Depression	21.0	17.4 25.1	403	22.8	19.1 27.0	401	20.6	16.8 24.9	367
	No Well-Baby Care	3.5	2.0 6.1	374	1.7	0.7 3.8	372	0.8	0.3 2.3	334
	Hearing Screening	86.8	83.4 89.6	402	91.6	88.6 93.9	404	95.0	92.1 96.9	368
Immun. Info By PNC Provider	44.4	39.9 49.0	403	47.9	43.4 52.6	404	44.7	39.9 49.6	369	
Baby Sleeps on Back	79.8	75.7 83.3	398	75.0	70.6 78.9	392	81.7	77.5 85.3	361	
Day Care	*	* *	*	*	* *	*	*	* *	*	
District 7		%	95% CI	n	%	95% CI	n	%	95% CI	n
	Unintended Pregnancy	29.7	25.6 34.2	419	30.7	26.4 35.4	407	28.3	24.1 33.0	365
	No Birth Control Use	57.1	49.0 64.9	136	59.9	51.9 67.5	141	58.4	50.6 65.9	143
	No Health Insurance	35.2	30.8 39.8	423	37.3	32.8 42.1	413	32.3	27.8 37.0	368
	Medicaid Utilization	47.8	43.2 52.4	416	51.4	46.8 56.1	409	47.5	42.6 52.4	367
	Overweight/Obesity	38.1	33.7 42.7	407	38.2	33.6 42.9	386	34.9	30.3 39.9	350
	First Trimester Prenatal Care	89.5	86.2 92.0	415	86.3	82.3 89.4	404	87.1	83.4 90.1	365
	WIC	41.5	37.1 46.1	423	44.8	40.2 49.6	411	44.8	40.0 49.7	367
	Multivitamin Use	77.9	73.8 81.5	421	76.3	72.1 80.1	408	69.5	64.8 73.8	365
	HIV Testing	49.8	45.3 54.3	423	49.9	45.2 54.6	407	43.4	38.6 48.3	363
	No Dental Care	62.3	57.8 66.5	417	57.6	52.9 62.1	409	61.5	56.7 66.1	365
	Exercise During Pregnancy	*	* *	*	*	* *	*	*	* *	*
	Stressful Life Events	*	* *	*	*	* *	*	*	* *	*
	Physical Abuse	4.0	2.5 6.4	422	5.4	3.6 8.1	413	2.1	1.0 4.2	368
	Smoking	7.1	5.0 10.1	420	5.4	3.5 8.1	411	5.0	3.2 7.7	367
	Alcohol Use	0.5	0.1 1.9	419	0.9	0.3 2.5	412	0.5	0.1 2.0	365
	Breastfeeding Initiation	90.7	87.7 93.1	421	87.5	84.0 90.4	412	92.5	89.5 94.7	367
	Breastfeeding Duration (6 mos +)	54.1	49.3 58.9	376	52.8	47.9 57.8	365	60.6	55.5 65.4	343
	Postpartum Depression	24.0	20.3 28.1	423	20.5	16.9 24.6	412	19.5	15.9 23.7	365
	No Well-Baby Care	3.3	1.9 5.6	394	3.1	1.7 5.6	383	1.9	0.9 4.0	342
	Hearing Screening	94.3	91.9 96.0	420	93.0	90.1 95.1	410	90.3	87.0 92.9	366
Immun. Info By PNC Provider	41.8	37.4 46.4	423	47.1	42.5 51.7	413	43.2	38.5 48.1	368	
Baby Sleeps on Back	72.9	68.6 76.8	412	69.7	65.1 73.9	406	76.9	72.5 80.7	364	
Day Care	*	* *	*	*	* *	*	*	* *	*	

*Indicates data are not available for that year

Note: Confidence Intervals (95% CI) may differ from previous publications due to revisions in standard error calculation.

Appendix A: Maternal and Infant Health Trends

By Health District, PRATS 2004-2010

	Year											
	2007			2008			2009			2010		
District 6	%	95% CI	n									
Unintended Pregnancy	37.9	33.1 43.0	358	33.7	29.1 38.6	350	32.4	27.8 37.4	370	32.6	28.0 37.6	354
No Birth Control Use	56.4	48.3 64.2	139	58.7	50.6 66.3	133	49.7	41.2 58.3	122	53.8	45.8 61.7	137
No Health Insurance	38.9	34.1 43.9	357	33.7	29.1 38.6	355	30.9	26.4 35.9	372	39.3	34.5 44.3	356
Medicaid Utilization	46.8	41.8 51.9	348	42.1	37.2 47.2	343	44.4	39.5 49.3	371	48.4	43.5 53.4	356
Overweight/Obesity	43.1	38.2 48.2	347	41.5	36.7 46.5	344	43.7	38.8 48.6	361	43.0	38.1 48.0	346
First Trimester Prenatal Care	86.0	82.1 89.1	348	89.3	85.6 92.1	339	89.8	86.5 92.4	369	90.8	87.4 93.4	354
WIC	48.1	43.2 53.1	358	43.7	38.9 48.7	348	51.3	46.4 56.1	372	45.0	40.1 50.0	357
Multivitamin Use	81.3	77.2 84.9	357	79.1	74.7 82.9	350	79.9	75.6 83.6	372	84.2	80.3 87.4	357
HIV Testing	48.7	43.7 53.7	355	49.4	44.4 54.4	341	44.5	39.7 49.4	371	40.2	35.4 45.1	351
No Dental Care	54.3	49.3 59.2	357	52.4	47.5 57.3	349	46.3	41.5 51.2	373	48.3	43.4 53.2	357
Exercise During Pregnancy	*	* *	*	55.8	50.9 60.6	352	57.3	52.4 62.0	370	55.0	50.0 59.9	353
Stressful Life Events	*	* *	*	25.7	21.6 30.4	355	32.6	28.0 37.6	371	28.4	24.1 33.2	357
Physical Abuse	4.1	2.4 7.0	359	4.6	2.8 7.5	353	4.7	2.9 7.5	371	3.3	1.8 5.8	354
Smoking	6.3	4.1 9.6	359	9.5	6.8 13.2	355	10.2	7.4 14.1	370	8.7	6.1 12.2	356
Alcohol Use	1.5	0.6 3.7	359	2.1	1.0 4.1	355	2.1	1.0 4.3	369	2.3	1.1 4.7	356
Breastfeeding Initiation	87.5	83.6 90.6	359	89.5	86.0 92.2	354	88.8	85.3 91.5	373	90.8	87.3 93.4	357
Breastfeeding Duration (6 mos +)	49.8	44.7 55.0	329	52.9	47.7 58.1	315	53.5	48.3 58.6	337	53.0	47.8 58.1	327
Postpartum Depression	17.5	14.0 21.7	359	20.0	16.3 24.4	354	22.4	18.4 26.9	373	17.6	14.1 21.8	356
No Well-Baby Care	2.5	1.4 4.6	325	1.8	0.8 4.0	313	2.4	1.2 4.6	361	2.0	0.9 4.2	341
Hearing Screening	95.6	92.9 97.3	359	94.0	91.2 96.0	356	98.1	96.4 99.0	371	97.4	95.1 98.7	356
Immun. Info By PNC Provider	48.1	43.2 53.1	358	70.8	66.0 75.2	323	68.6	63.8 73.1	342	66.6	61.6 71.2	327
Baby Sleeps on Back	86.6	82.8 89.7	351	80.0	75.8 83.7	352	78.7	74.4 82.5	368	83.3	79.1 86.8	348
Day Care	*	* *	*	32.1	25.7 39.2	164	35.6	29.0 42.8	171	31.4	25.0 38.6	164
District 7	%	95% CI	n									
Unintended Pregnancy	27.4	23.2 32.1	368	29.2	25.1 33.6	395	27.1	23.0 31.7	382	27.9	23.6 32.7	372
No Birth Control Use	62.8	54.2 70.7	119	61.8	53.8 69.2	136	58.6	50.3 66.4	132	59.0	50.2 67.2	120
No Health Insurance	31.3	26.9 36.1	370	33.8	29.5 38.3	398	30.9	26.5 35.6	385	31.2	26.7 36.1	373
Medicaid Utilization	43.6	38.7 48.5	362	45.4	40.8 50.2	390	45.6	40.9 50.5	383	45.1	40.2 50.0	372
Overweight/Obesity	35.9	31.3 40.9	354	37.5	33.0 42.2	387	35.7	31.2 40.5	379	37.4	32.7 42.3	364
First Trimester Prenatal Care	88.6	85.0 91.4	359	92.5	89.6 94.7	388	94.1	91.3 96.0	379	91.2	87.9 93.7	371
WIC	39.4	34.7 44.3	368	43.0	38.4 47.7	391	42.3	37.6 47.1	384	41.0	36.2 46.0	372
Multivitamin Use	75.3	70.8 79.3	368	79.0	74.9 82.6	392	78.7	74.5 82.4	384	80.5	76.3 84.0	372
HIV Testing	38.5	33.8 43.4	364	41.1	36.5 45.8	387	39.8	35.2 44.7	380	37.4	32.7 42.4	365
No Dental Care	55.6	50.7 60.4	367	51.6	46.9 56.3	392	46.6	41.9 51.4	384	52.7	47.8 57.6	370
Exercise During Pregnancy	*	* *	*	57.1	52.4 61.7	392	56.4	51.5 61.1	383	58.5	53.6 63.3	370
Stressful Life Events	*	* *	*	20.9	17.3 24.9	393	21.0	17.2 25.4	385	23.4	19.3 28.0	373
Physical Abuse	2.4	1.2 4.5	369	2.8	1.6 4.9	396	5.5	3.5 8.4	381	1.8	0.8 3.9	371
Smoking	5.9	3.9 8.9	367	5.5	3.7 8.1	392	5.1	3.2 8.0	385	4.5	2.7 7.3	373
Alcohol Use	1.5	0.6 3.4	367	1.0	0.4 2.6	392	1.1	0.4 2.9	385	1.4	0.6 3.3	372
Breastfeeding Initiation	93.2	90.3 95.4	370	94.1	91.5 96.0	396	90.7	87.3 93.2	381	92.7	89.7 94.9	371
Breastfeeding Duration (6 mos +)	55.0	50.0 60.0	342	60.4	55.5 65.2	356	62.8	57.7 67.6	348	54.3	49.3 59.3	351
Postpartum Depression	16.0	12.7 19.9	370	18.3	14.9 22.2	395	18.4	14.9 22.6	380	16.2	12.9 20.2	371
No Well-Baby Care	3.2	1.8 5.7	325	2.3	1.2 4.3	355	1.0	0.4 2.6	369	2.9	1.6 5.3	362
Hearing Screening	92.6	89.7 94.8	369	94.9	92.4 96.6	396	94.2	91.4 96.1	381	96.0	93.7 97.6	371
Immun. Info By PNC Provider	46.7	41.9 51.6	369	59.4	54.6 64.1	371	66.6	61.7 71.1	356	64.8	59.8 69.5	344
Baby Sleeps on Back	80.1	75.9 83.7	362	82.6	78.7 85.8	395	79.0	74.8 82.6	378	81.7	77.5 85.2	370
Day Care	*	* *	*	31.0	24.4 38.5	149	29.2	22.2 37.4	132	27.7	20.8 35.9	131

*Indicates data are not available for that year

Note: Confidence Intervals (95% CI) may differ from previous publications due to revisions in standard error calculation.

Appendix B: PRATS Questionnaire and Results, 2010

In the following pages, the 2010 Idaho Pregnancy Risk Assessment Tracking System (PRATS) is presented in its entirety. In addition to the text of each question, the results for most of the questions are included. The percentage of each response category can be found in parentheses to the right of the corresponding response category. Also note that the percentages are weighted results (see the *Weighting the Data* section on page 2 for further explanation), and percentages may not always sum to 100 percent due to rounding. The results have not been provided for all questions.

NOTE: Care should be exercised when interpreting these results. In order to interpret the results correctly, it is important to understand what group has been asked the question. For example, question 10 on page 91 indicates 54.9 percent of respondents reported that they were not using any kind of birth control when they got pregnant. However, this question is only asked of those who responded “No” to question 9, or those who were not trying to get pregnant when they got pregnant. The exclusion of certain respondents is indicated by the italicized text in brackets at the beginning of question 10. These are called “skip patterns” and are used extensively throughout the survey. In order to properly identify what group has been asked a specific question, the skip pattern must be considered. In addition, there may be some confusion when trying to compare results from Appendix B to those in previous sections. The risk factors reported in the main body of the report are often constructed from the responses of multiple questions and may not be comparable to the results of one particular question.

Missing data were excluded from the n’s, percentages, and means calculations. Other responses may also be excluded and are indicated by showing the response without any percentage. Also, the results of some questions, such as question 1 about the survey date, are suppressed due to the length of resulting lists.

For help with interpretation of these results, contact the Idaho PRATS Program Director at (208) 332-7366.

Appendix B: PRATS Questionnaire and Results, 2010

When answering the following questions, please refer to the time when you were pregnant with the baby you gave birth to in 2010. If you delivered twins or triplets during your pregnancy, please refer to your experience with the first- born (of the twins or triplets) when answering the questions.

First, please tell us...

1. What is today's date?

____/____/____

Next, we would like to ask a few questions about you and the time before you became pregnant with your baby. Please check the box next to your answer.

2. **Just before** you got pregnant, did you have health insurance? (Don't count Medicaid.)

No (37.4%)

Yes (62.6%)

3. **Just before** you got pregnant, were you on Medicaid?

No (95.5%)

Yes (4.6%)

4. In the **month before** you got pregnant with your baby, how many times a week did you take a multiple vitamin (a pill that contains many different vitamins and minerals)?

Check one answer.

I didn't take a multiple vitamin at all (46.7%)

1 to 3 times a week (12.0%)

4 to 6 times a week (9.6%)

Every day of the week (31.7%)

5. Have you ever heard or read that taking the vitamin folic acid can help prevent some birth defects?

No (15.8%)

Yes (84.2%)

6. **Just before** you got pregnant, how much did you weigh?

_____ Pounds

Mean (153.4 lbs.)

Median (144.5 lbs.)

Minimum (86.0 lbs.)

Maximum (350.0 lbs.)

7. **How tall are you without shoes?**

_____Feet _____Inches

Mean (5'5")

Median (5'4")

Minimum (4'7")

Maximum (6'7")

8. **Thinking back to just before you got pregnant, how did you feel about becoming pregnant? Check one answer.**

I wanted to be pregnant sooner (17.0%)

I wanted to be pregnant later (25.1%)

I wanted to be pregnant then (49.8%)

I didn't want to be pregnant then or at any time in the future (8.1%)

9. **When you got pregnant with your baby, were you trying to become pregnant?**

No (42.0%) **Go to Question 10**

Yes (58.0%) **Go to Question 12**

10. ***[Only asked of mothers who were not trying to get pregnant]* When you got pregnant with your baby, were you or your partner using any kind of birth control?**

Birth Control includes oral contraceptive ("the pill"), condom, skin patch, cervical ring, intrauterine device (IUD), diaphragm, spermicide, implants, (Norplant®), shots (DepoProvera®), or other methods such as not having sex at certain times (rhythm method), having your tubes tied, or your partner having a vasectomy.

No (54.9%) **Go to Question 11**

Yes (45.1%) **Go to Question 12**

11. ***[Only asked of mothers who were not trying to get pregnant]* Why were you or your partner not using any birth control?**

For each item, check "Yes" if it applied to you or "No" if it did not. "Yes" is reported.

I didn't mind if I got pregnant (64.3%)

I didn't think I could get pregnant (35.6%)

I had been having side effects from the birth control I used (26.1%)

I didn't want to use birth control (45.0%)

My partner didn't want to use birth control (30.5%)

I didn't think I was going to have sex (9.8%)

I couldn't afford birth control (27.2%)

Other → Please tell us (38.6%)

12. **During the 3 months before you got pregnant with your baby, did you have any of the following health problems?**

For each item, check "Yes" if you had the problem or "No" if you did not. "Yes" is reported.

Asthma (6.8%)

High blood pressure (hypertension) (1.8%)

High blood sugar (diabetes) (0.7%)

Anemia (poor blood, low iron) (4.8%)

Heart problems (0.8%)

The next questions are about the prenatal care you received during your pregnancy. Prenatal care includes visits to a doctor, nurse, or other health care provider, before your baby was born, to get checkups and advice about pregnancy.

13. How many weeks pregnant were you when you were sure you were pregnant?

For example, you had a pregnancy test or a doctor or nurse said you were pregnant.

_____ Weeks

Mean (5.8 weeks)

Median (4.7 weeks)

Minimum (1.0 week)

Maximum (40.0 weeks)

14. How many weeks pregnant were you when you had your first visit for prenatal care?

_____ Weeks **Go to Question 15**

Mean (9.2 weeks)

Median (7.9 weeks)

Minimum (0.0 weeks)

Maximum (32.0 weeks)

I didn't go for prenatal care (0.3%) **Go to Question 21**

15. *[Only asked of mothers who received prenatal care]* Did you get prenatal care as early in your pregnancy as you wanted?

No (11.8%) **Go to Question 16**

Yes (88.2%) **Go to Question 17**

16. *[Only asked of mothers who received prenatal care, but not as early as wanted]* Which of these things kept you from getting prenatal care as early as you wanted?

For each item, check "Yes" if it applied to you or "No" if it did not. "Yes" is reported.

I didn't have my Medicaid card (33.6%)

I couldn't get an appointment (29.9%)

I didn't have enough money or insurance to pay for my visits (44.5%)

I didn't know that I was pregnant (37.1%)

I had no way to get to the clinic or doctor's office (10.2%)

The doctor would not start care earlier (23.9%)

I had a hard time finding a doctor or health care provider who would take me as a patient (8.2%)

I had too many other things going on (13.2%)

Other → Please tell us (23.4%)

17. **[Only asked of mothers who received prenatal care] Who did you go to most of the time for your prenatal care visits?**

Don't include visits for WIC.

Check one answer.

Physician/Medical Doctor (MD) such as OB-GYN or family doctor (86.7%)

Physician Assistant (PA), Nurse Practitioner (NP), or Registered Nurse (RN) (3.5%)

Certified Nurse-Midwife (CNM) (6.4%)

Other Midwife (Certified Professional Midwife, Certified Midwife, lay or direct entry midwife) (3.1%)

Other → Please tell us (0.4%)

18. **[Only asked of mothers who received prenatal care] During any of your prenatal care visits, did a doctor, nurse, or other health care provider give you information about any of the issues listed below?**

For each item, check "Yes" if you were given information or "No" if you were not given information. "Yes" is reported.

What you should eat during your pregnancy (83.0%)

How eating fish containing high levels of mercury could affect your baby (71.4%)

How much weight you should gain during pregnancy (82.7%)

The importance of taking vitamin supplements during your pregnancy (95.8%)

The importance of getting regular dental care during your pregnancy (57.2%)

Which kinds of medicine are safe to take during your pregnancy (91.4%)

How drinking alcohol during pregnancy could affect your baby (87.3%)

How smoking during pregnancy could affect your baby (87.7%)

How using illegal drugs during pregnancy could affect your baby (85.5%)

Doing tests to screen for birth defects or diseases that run in your family (87.4%)

Getting your blood tested for HIV (the virus that causes AIDS) (79.1%)

What happens if a baby is shaken (73.1%)

Physical abuse to women by their partners (60.7%)

The benefits of breastfeeding your baby (89.5%)

19. **[Only asked of mothers who received prenatal care] How was your prenatal care paid for?**

Check all that apply.

Medicaid (38.9%)

Personal income (cash, check, or credit card) (45.0)

Health insurance or HMO (54.8%)

The military, CHAMPUS, or TriCare (3.2%)

Other → Please tell us (3.1%)

20. **[Only asked of mothers who received prenatal care] During any of your prenatal care visits, did a doctor, nurse, or other health care provider ask you...?**

For each item, check "Yes" if you were asked, or "No" if you were not asked. "Yes" is reported.

If you were drinking alcohol during your pregnancy (82.0%)

If you were smoking cigarettes during your pregnancy (84.1%)

If you were using illegal drugs during your pregnancy (79.9%)

The next questions are about your pregnancy and the things that may have happened during your pregnancy.

21. Did you have any of these problems during your pregnancy?

For each item, check "Yes" if you had the problem or "No" if you did not. "Yes" is reported.

High blood sugar (gestational diabetes) that started during this pregnancy (8.6%)

Vaginal bleeding (14.5%)

Kidney or bladder (urinary tract) infection (15.5%)

Severe nausea, vomiting, or dehydration (22.5%)

Cervix had to be sewn shut (incompetent cervix) (1.2%)

High blood pressure, hypertension, (including pregnancy-induced hypertension [PIH], preeclampsia, or toxemia) (12.4%)

Problems with the placenta (such as abruption placentae or placenta previa) (5.2%)

Labor pains more than 3 weeks before my baby was due (preterm or early labor) (18.2%)

Water broke more than 3 weeks before my baby was due (premature rupture of membranes [PROM]) (4.4%)

22. Did you try to get Medicaid coverage during this pregnancy?

No (57.5%) Go to Question 24

Yes (42.5%) Go to Question 23

23. *[Only asked of mothers who applied for Medicaid during pregnancy]* Did any of these things happen when you tried to get Medicaid during your pregnancy?

For each item, check "Yes" if it happened to you or "No" if it did not. "Yes" is reported.

I had a hard time getting help from the Medicaid staff (15.2%)

I was told I was not eligible for Medicaid services because my household made too much money (15.3%)

I was told I was not eligible for Medicaid services because I am from another country (2.8%)

I did not understand how to use my Medicaid card or what was covered (7.5%)

I did not get all of the Medicaid services that I needed (10.1%)

I had a problem finding a doctor who would accept me as a Medicaid patient (2.3%)

I was assigned to a doctor that I did not choose (3.0%)

Other → Please tell us (15.7%)

24. At any time during your pregnancy, did you have a blood test for HIV (the virus that causes AIDS)?

No (29.1%)

Yes (45.5%)

I don't know (25.5%)

25. During your pregnancy, did you get the H1N1 flu vaccine? (H1N1 flu is also referred to as the swine flu, but is different than seasonal flu.)

No (50.3%)

Yes (49.7%)

26. **During your pregnancy**, did you participate in the WIC Program (the Supplemental Nutrition Program for Women, Infants, and Children)?

No (60.1%)

Yes (39.9%)

27. **During the last 3 months** of your pregnancy, how many times a week did you take a multiple vitamin (a pill that contains many different vitamins and minerals)?

Check one answer.

I didn't take a multiple vitamin at all (7.7%)

1 to 3 times a week (9.2%)

4 to 6 times a week (13.6%)

Every day of the week (69.6%)

The next questions are about the care of your teeth and gums before and during your pregnancy.

28. **During your pregnancy**, did you go to a dentist or dental clinic for routine care such as teeth cleaning or regular check-up?

No (48.9%) Go to Question 29

Yes (51.1%) Go to Question 30

29. *[Only asked of mothers who did not receive dental care during pregnancy]* Why did you **not** go to a dentist or dental clinic for routine care during your pregnancy?

Check all that apply.

It had been less than 1 year since my last dental visit (17.4%)

I did not have enough money or insurance to pay for the visit (46.3%)

I did not feel that I needed to go to a dentist (26.6%)

I did not feel that it would be safe for my baby (9.2%)

I had too many other things going on (25.2%)

Other → Please tell us (13.3%)

30. **During your pregnancy**, did you notice any of the following conditions in your mouth?

For each item, check "Yes" if you had the condition or "No" if you did not. **"Yes" is reported.**

Bleeding gums when brushing (47.5%)

Swollen, puffy gums (22.6%)

Loose teeth (4.0%)

Receding gums or teeth that appear longer (9.0%)

Pain in teeth or gums (24.9%)

Bad breath that does not go away with normal brushing (8.3%)

Spaces between the teeth that were not there before (3.2%)

Inability to chew or swallow properly (2.4%)

Teeth with obvious decay (11.3%)

31. How long has it been since you had your teeth cleaned by a dentist or dental hygienist?

Less than 1 year (52.2%)

1 to 2 years (23.6%)

3 or more years (19.8%)

I have never had my teeth cleaned by a dentist or dental hygienist (1.9%)

I don't know (2.5%)

The next questions are about smoking cigarettes and drinking alcohol.

32. Have you smoked at least 100 cigarettes (5 packs) in your entire life?

A pack has 20 cigarettes.

No (71.2%)

Yes (28.8%)

33. *[Only asked of mothers who smoked at least 100 cigarettes in their entire life]* In the 3 months before you got pregnant, how many cigarettes did you smoke on an average day?

None (0 cigarettes) (39.3%)

Less than 1 cigarette a day (5.3%)

1 to 5 cigarettes (18.3%)

6 to 10 cigarettes (23.5%)

11 to 20 cigarettes (12.7%)

21 to 40 cigarettes (0.8%)

41 cigarettes or more (0.2%)

34. *[Only asked of mothers who smoked at least 100 cigarettes in their entire life]* In the last 3 months of your pregnancy, how many cigarettes did you smoke on an average day?

None (0 cigarettes) (72.2%)

Less than 1 cigarette a day (5.0%)

1 to 5 cigarettes (15.8%)

6 to 10 cigarettes (5.4%)

11 to 20 cigarettes (1.4%)

21 to 40 cigarettes (0.0%)

41 cigarettes or more (0.2%)

35. *[Only asked of mothers who smoked at least 100 cigarettes in their entire life]* How many cigarettes do you smoke on an average day now?

None (0 cigarettes) (57.2%)

Less than 1 cigarette a day (3.7%)

1 to 5 cigarettes (16.5%)

6 to 10 cigarettes (15.8%)

11 to 20 cigarettes (6.1%)

21 to 40 cigarettes (0.5%)

41 cigarettes or more (0.2%)

36. Have you had any alcoholic drinks in the past 2 years?

A drink is equal to a 12 ounce beer or malt beverage, 5 ounce glass of wine, or drink with one shot (1.5 ounces) of liquor.

No (46.5%) **Go to Question 39**

Yes (53.5%) **Go to Question 37**

37. [*Only asked of mothers who had alcohol in the past 2 years*] During the 3 months before you got pregnant, how many alcoholic drinks did you have in an average week?

I didn't drink then (21.3%)

Less than 1 drink a week (39.9%)

1 to 3 drinks a week (26.7%)

4 to 6 drinks a week (7.6%)

7 or more drinks a week (4.5%)

38. [*Only asked of mothers who had alcohol in the past 2 years*] During the last 3 months of your pregnancy, how many alcoholic drinks did you have in an average week?

I didn't drink then (92.9%)

Less than 1 drink a week (6.1%)

1 to 3 drinks a week (0.8%)

4 to 6 drinks a week (0.0%)

7 or more drinks a week (0.1%)

The next questions are about physical activity or exercise before and during your pregnancy.

39. During the 3 months before you got pregnant with your baby, how often did you participate in any physical activities or exercise for 30 minutes or more? (For example: walking for exercise, swimming, cycling, dancing, or gardening.)

Do not count exercise you may have done as part of your regular job.

Never (7.4%)

Less than 1 day per week (17.9%)

1 to 4 days per week (56.9%)

5 or more days per week (17.9%)

I was told by a doctor, nurse, or health care provider not to exercise (0.0%)

40. During the last 3 months of your pregnancy, how often did you participate in any physical activities or exercise for 30 minutes or more?

Do not count exercise you may have done as part of your regular job.

Never (14.0%)

Less than 1 day per week (25.3%)

1 to 4 days per week (45.7%)

5 or more days per week (11.6%)

I was told by a doctor, nurse, or health care provider not to exercise (3.4%)

Pregnancy can be a difficult time for some women. The next questions are about things that may have happened before and during your pregnancy.

41. This question is about things that may have happened during the 12 months before your baby was born.

For each item, check "Yes" if it happened to you or check "No" if it did not. "Yes" is reported.

- A close family member was very sick and had to go into the hospital (20.3%)
- I got separated or divorced from my husband or partner (7.1%)
- I moved to a new address (35.7%)
- I was homeless (1.9%)
- My husband or partner lost his job (15.9%)
- I lost my job even though I wanted to go on working (9.1%)
- I argued with my husband or partner more than usual (21.6%)
- My husband or partner said he didn't want me to be pregnant (6.6%)
- I had a lot of bills I couldn't pay (27.3%)
- I was in a physical fight (2.3%)
- I or my husband or partner went to jail (3.0%)
- Someone very close to me had a bad problem with drinking or drugs (10.8%)
- Someone very close to me died (14.6%)

42. During your pregnancy, would you have had the kinds of help listed below if you needed them?

For each one, check "Yes" if you would have had it or check "No" if not. "Yes" is reported.

- Someone to loan me \$50 (85.1%)
- Someone to help me if I were sick and needed to be in bed (87.8%)
- Someone to take me to the clinic or doctor's office if I needed a ride (90.9%)
- Someone to talk with about my problems (89.2%)

43. During your pregnancy, did anyone push, hit, slap, kick, choke, or physically hurt you in any way?

- No (95.8%)
- Yes (4.2%)

The next questions are about your labor and delivery.

44. How much did you weigh at delivery?

_____ Pounds

- Mean (184.3 lbs.)
- Median (178.0 lbs.)
- Minimum (110.0 lbs.)
- Maximum (400.0 lbs.)

45. **After the birth of your baby, how long did you stay in the hospital or birthing center?**

- Less than 24 hours (Less than 1 day) (10.5%)
- 24-48 hours (1-2 days) (61.1%)
- 3 days (18.7%)
- 4 days (5.1%)
- 5 days (2.1%)
- 6 days or more (1.1%)
- I didn't have my baby in a hospital or birthing center (1.5%)

46. **After your baby was born, how long did your baby stay in the hospital or birthing center?**

- My baby is still in the hospital (0.1%)
- Less than 24 hours (Less than 1 day) (11.0%)
- 24-48 hours (1-2 days) (57.1%)
- 3 days (16.7%)
- 4 days (4.5%)
- 5 days (2.2%)
- 6 days or more (6.7%)
- I didn't have my baby in a hospital or birthing center (1.7%)

47. **After your baby was born, was his or her hearing tested?**

- No (1.2%)
- Yes (95.1%)
- I don't know (3.7%)

48. **How was your delivery paid for?**

- Medicaid (43.0%)
- Personal income (cash, check, or credit card) (38.3%)
- Health insurance or HMO (53.1%)
- The military, CHAMPUS, or TriCare (2.9%)
- Other → Please tell us (3.8%)

The next questions are about the time since your baby was born.

49. **During the 6 weeks after your baby was born, did a doctor, nurse, or other health care worker give you information about using birth control?**

- No (11.8%)
- Yes (88.2%)

50. **During the 3 months after your delivery, how often would you say you felt down, depressed, or hopeless?**

Check one answer.

- Always (3.6%)
- Often (14.3%)
- Sometimes (28.5%)
- Rarely (31.4%)
- Never (22.2%)

51. **During the 3 months after your delivery, how often would you say you had little interest or little pleasure in doing things?**

Check one answer.

Always (3.2%)
Often (11.3%)
Sometimes (21.9%)
Rarely (35.8%)
Never (27.8%)

52. **During the 3 months after your delivery, would you say that you were...**

Check one answer.

Not depressed at all (42.9%)	Go to Question 54
A little depressed (40.3%)	Go to Question 53
Moderately depressed (12.0%)	Go to Question 53
Very depressed (4.9%)	Go to Question 53

53. **Since the birth of your baby, have any of the following things happened?**

For each item, check "Yes" if it has happened to you or "No" if it has not. **"Yes" is reported.**
Professional may include a psychiatrist, psychologist, counselor, family practice doctor, obstetrician, gynecologist, nurse, etc...

I wanted to see a professional about my depression (28.1%)
I went to see a professional about my depression (20.8%)
I participated in a support group for depression (1.2%)
I was prescribed medication to treat my depression (22.9%)

The next questions are about breastfeeding your baby.

54. **Did you ever breastfeed or pump breast milk to feed your baby?**

No (8.5%)	Go to Question 59
Yes (91.5%)	Go to Question 55

55. ***[Only asked of mothers who ever breastfed]* Are you still breastfeeding or pumping breast milk to feed your baby?**

No (68.1%)	Go to Question 56
Yes (31.9%)	Go to Question 58

56. ***[Only asked of mothers who were breastfeeding but stopped]* How many weeks or months did you breastfeed or pump milk to feed your baby?**

_____ Weeks **OR** _____ Months

Mean (21.4 weeks)
Median (16.7 weeks)
Minimum (0 weeks)
Maximum (73.7 weeks)

Less than 1 week (3.1%)

57. **[Only asked of mothers who were breastfeeding but stopped] Why did you stop breastfeeding?**

Check all that apply.

- My baby lost interest in breastfeeding (21.6%)
- My baby had difficulty nursing (18.9%)
- Breast milk alone did not satisfy my baby (37.0%)
- I thought my baby was not gaining enough weight (9.7%)
- I had nipple or breast problems (12.9%)
- My baby's teeth made it painful to continue (6.3%)
- I felt it was the right time to stop (25.2%)
- I didn't have enough milk (45.9%)
- I went back to work or school (23.1%)
- It was inconvenient to continue (11.5%)
- Other → Please tell us (19.4%)

58. **[Only asked of mothers who ever breastfed] How many weeks or months old was your baby when you started feeding him/her formula?**

_____ Weeks **OR** _____ Months

- Mean (13.4 weeks)
- Median (8.4 weeks)
- Minimum (0 weeks)
- Maximum (69.3 weeks)

- Less than 1 week (14.8%)
- My baby has never been fed formula (24.7 %)

59. **How many weeks or months old was your baby when you started feeding him/her foods other than breast milk or formula (for example: baby cereal, baby food, or solid food)?**

_____ Weeks **OR** _____ Months

- Mean (22.7 weeks)
- Median (21.3 weeks)
- Minimum (0 weeks)
- Maximum (52.0 weeks)

- Less than 1 week (0.2%)
- My baby has never been fed foods other than breast milk or formula (1.7 %)

The next questions are about the care of your baby.

60. **How do you usually lay your baby down to sleep?**

Check one answer.

- On his or her side (12.3%)
- On his or her back (80.8%)
- On his or her stomach (6.9%)

61. **About how many hours a day, on average, is your baby in the same room with someone who is smoking?**

_____ Hours

1 or more hours a day (0.7%)

Less than 1 hour a day (2.5%)

My baby is never in the same room with someone who is smoking (96.9%)

62. **About how many times has your baby been to a doctor or nurse for routine well-baby check-ups?**

Routine well-baby check-ups may include height/weight measurements, physical examination, developmental and behavioral assessment, sleep position and nutrition counseling, immunizations, and other information or referrals as necessary. Don't count when you took your baby for care when he or she was sick or visits only for immunizations.

_____ Times

1 or more routine well-baby visits (97.9%)

My baby hasn't been for routine well-baby care (2.1%)

63. **What, if anything, prevented your baby from having routine well-baby check-ups? Check all that apply.**

I didn't have enough money or insurance to pay for it (3.1%)

I couldn't get an appointment (1.9%)

I had no way to get the baby to the clinic or doctor's office (1.5%)

I didn't have anyone to take care of my other children (2.0%)

I didn't know where to go (0.6%)

I didn't know about routine well-baby check-ups (1.4%)

Other → Please tell us (3.9%)

Nothing prevented my baby from having routine well-baby check-ups (81.1%)

64. **Is your baby enrolled in the Immunization Reminder Information System (IRIS)?**

No (9.9%)

Yes (71.7%)

I don't know (18.4%)

65. **Were you given information about the importance of immunizing your baby by any of the following?**

For each item, check "Yes" if you were given information or "No" if you were not. "Yes" is reported.

Prenatal care provider (70.7%)

Childbirth preparation or parenting classes (35.6%)

Hospital or birthing center (82.3%)

Pediatrician, family doctor, or other health care provider (94.3%)

Other → Please tell us (19.5%)

66. **Are your baby's immunizations (shots) up to date according to the immunization schedule?**

No (11.1%)

Go to Question 67

Yes (88.4%)

Go to Question 68

I don't know (0.6%)

Go to Question 67

67. *[Only asked of mothers whose baby's immunization were not up to date]* If no or don't know, why are your baby's immunizations (shots) not up to date?
Check all that apply.

My baby was sick or had medical problems (23.1%)
I wanted to wait until my baby was older (45.2%)
I didn't have enough money or insurance (9.9%)
I had no way to get the baby to the clinic or office (5.1%)
The clinic was not open when it was convenient for me (6.2%)
I don't know what is recommended by the schedule (2.9%)
I had too many other things going on (6.4%)
I don't think immunizations are safe (28.9%)
Other → Please tell us (29.7%)

68. Did your baby receive the Hepatitis B vaccine before leaving the hospital or birthing center?

No (11.9%) Go to Question 69
Yes (79.9%) Go to Question 70
I don't know (7.0%) Go to Question 70
I did not have my baby in a hospital or birthing center (1.1%) Go to Question 70

69. *[Only asked of mothers whose baby did not receive the Hepatitis B vaccine]* Why didn't your baby receive the Hepatitis B vaccine before leaving the hospital or birthing center?
Check all that apply.

I was not offered the Hepatitis B vaccine for my baby at that time (12.4%)
I felt my baby was too young for this vaccination (60.0%)
I felt my baby was not at risk for Hepatitis B (39.7%)
I was worried that the vaccine would harm my baby (27.7%)
Other → Please tell us (25.3%)

70. What type of health insurance is your baby covered by?
Check all that apply.

Medicaid or CHIP (45.8%)
Health Insurance or HMO (50.5%)
The military, CHAMPUS, or TriCare (2.9%)
Other → Please tell us (1.9%)
My baby is not covered by any type of health insurance (4.5%)

The next questions are about you and your family.

71. Are you currently in school or working outside of the home?

No (51.0%) Go to Question 74
Yes (49.1%) Go to Question 72

72. [Only asked of mothers who are currently in school or are working outside the home] When you go to work or school, do you take your baby to day care?

Check one answer.

No (63.6%)

Go to Question 74

Yes, and it is licensed (27.5%)

Go to Question 73

Yes, and it is not licensed (7.7%)

Go to Question 73

Yes, and I don't know if it is licensed (1.2%)

Go to Question 73

73. [Only asked of mothers who take their baby to day care while they are in school or who work outside the home] When you leave your baby at day care, how often do you feel that he or she is safe and well cared for?

Check one answer.

Always (87.6%)

Often (10.2%)

Sometimes (1.3%)

Rarely (0.0%)

Never (0.8%)

74. **In the 12 months before your delivery, what was your total household income from all sources, before taxes?**

Include your income, your husband's or partner's income, and any other income you may have used.

Check one answer.

Less than \$10,000 (14.9%)

\$10,000 to \$14,999 (9.5%)

\$15,000 to \$19,999 (8.2%)

\$20,000 to \$24,999 (9.4%)

\$25,000 to \$29,999 (6.9%)

\$30,000 to \$34,999 (7.9%)

\$35,000 to \$49,999 (13.8%)

\$50,000 to \$74,999 (15.5%)

\$75,000 or more (14.0%)

Thanks for answering our questions! Your answers will help us work to make Idaho mothers and babies healthier.

Appendix C: Technical Notes

Sampling Fractions

Birth records were sampled using the following sampling fractions:

Sampling Stratum	Sampling Frame	Sample	Approximate Sampling Fraction
<i>TOTAL</i>	21,144	4,200	1 in 5
Health District 1	2,089	600	1 in 3
Health District 2	1,017	600	1 in 2
Health District 3	3,497	600	1 in 6
Health District 4	5,394	600	1 in 9
Health District 5	2,812	600	1 in 5
Health District 6	2,520	600	1 in 4
Health District 7	3,815	600	1 in 6

Response and Refusal Rates

The overall response rate was 56.6 percent, computed as the number of completed surveys (2,377) divided by the total sample (4,200). Response rates varied depending upon health district (see the following table). District 7 had the highest response rate at 62.2 percent and District 3 had the lowest response rate at 45.5 percent.

Sampling Stratum	Sample	Number Completed	Percent Completed
<i>TOTAL</i>	4,200	2377	56.6%
Health District 1	600	322	53.7%
Health District 2	600	345	57.5%
Health District 3	600	273	45.5%
Health District 4	600	367	61.2%
Health District 5	600	340	56.7%
Health District 6	600	357	59.5%
Health District 7	600	373	62.2%

The overall refusal rate was 6.8 percent, computed as the number of women who refused the survey anytime during the mail or telephone phase (286) divided by the total sample (4,200).

Completion Rates by Survey Phase

Of the 2,377 completed surveys, 90.2 percent were completed by mail and 9.8 percent were completed by telephone (see the following table). The first mailing had the highest return, accounting for 67.2 percent of completed mail surveys. Returns from the second mailing accounted for 15.6 percent of completed mail surveys, and the third mailing accounted for 7.4 percent of completed mail surveys. The telephone phase, accounting for 9.8 percent of all completed surveys, was an important tool for

reaching women with low education, low income, younger age, and/or Spanish speakers.

Survey Phase	Number Completed	Percent Completed
<i>TOTAL</i>	<i>2,377</i>	<i>100.0%</i>
Mailing Phase	2,144	90.2%
Mailing 1	1,598	67.2%
Mailing 2	370	15.6%
Mailing 3	176	7.4%
Telephone Phase	233	9.8%

Costs associated with this publication are available from the
Idaho Department of Health and Welfare

HW-1206

December 2013